

D 39

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
3 May 2001 (03.05.2001)

PCT

(10) International Publication Number  
WO 01/31014 A2

(51) International Patent Classification<sup>7</sup>: C12N 15/12,  
15/63, C07K 14/72, 16/28, 16/42, C12N 5/20, G01N  
33/542, 33/68, A61P 25/00, C12Q 1/68

US 09/454,399 (CIP)  
Filed on 3 December 1999 (03.12.1999)  
US 09/481,794 (CIP)  
Filed on 12 January 2000 (12.01.2000)

(21) International Application Number: PCT/US00/29601

(22) International Filing Date: 27 October 2000 (27.10.2000)

(71) Applicant (for all designated States except US): PHAR-  
MACIA & UPJOHN COMPANY [US/US]; 301 Henri-  
etta Street, Kalamazoo, MI 49001 (US).

(25) Filing Language: English

(26) Publication Language: English

(72) Inventors; and  
(75) Inventors/Applicants (for US only): VOGELI, Gabriel  
[US/US]; 2576 Ninth Avenue West, Seattle, WA 98119  
(US). WOOD, Linda, S. [US/US]; 10193 FoxHol-  
low, Portage, MI 49024 (US). MERCHANT, Kalpana  
[US/US]; 5015 Glencove Lane, Portage, MI 49024 (US).

(30) Priority Data:  
09/427,653 27 October 1999 (27.10.1999) US  
09/427,859 27 October 1999 (27.10.1999) US  
09/428,020 27 October 1999 (27.10.1999) US  
09/428,114 27 October 1999 (27.10.1999) US  
09/429,695 28 October 1999 (28.10.1999) US  
09/429,676 28 October 1999 (28.10.1999) US  
09/429,555 28 October 1999 (28.10.1999) US  
09/429,517 28 October 1999 (28.10.1999) US  
09/454,399 3 December 1999 (03.12.1999) US  
09/481,794 12 January 2000 (12.01.2000) US

(74) Agent: GASS, David, A. Marshall, O'Toole, Gerstein,  
Murray & Borun; 6300 Sears Tower, 233 South Wacker  
Drive, Chicago, IL 60606-6402 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:  
Without international search report and to be republished  
upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(63) Related by continuation (CON) or continuation-in-part  
(CIP) to earlier applications:

US 09/427,653 (CIP)  
Filed on 27 October 1999 (27.10.1999)  
US 09/427,859 (CIP)  
Filed on 27 October 1999 (27.10.1999)  
US 09/428,020 (CIP)  
Filed on 27 October 1999 (27.10.1999)  
US 09/428,114 (CIP)  
Filed on 27 October 1999 (27.10.1999)  
US 09/429,695 (CIP)  
Filed on 28 October 1999 (28.10.1999)  
US 09/429,676 (CIP)  
Filed on 28 October 1999 (28.10.1999)  
US 09/429,555 (CIP)  
Filed on 28 October 1999 (28.10.1999)  
US 09/429,517 (CIP)  
Filed on 28 October 1999 (28.10.1999)

(54) Title: G PROTEIN-COUPLED RECEPTORS EXPRESSED IN BRAIN

(57) Abstract: The present invention provides genes encoding heretofore unknown G protein-coupled receptors, constructs and recombinant host cells incorporating the genes; the GPCR polypeptides encoded by the genes; antibodies to the polypeptides; and methods of making and using all of the foregoing.



WO 01/31014 A2

- 1. -

## G PROTEIN-COUPLED RECEPTORS EXPRESSED IN BRAIN

### RELATED APPLICATIONS

This patent application is a continuation-in-part of the following U.S. patent applications: Serial No. 09/481,794 filed January 12, 2000; Serial No. 09/454,399 filed December 3, 1999; Serial Nos. 09/429,517, 09/429,555, 09/429,676, 09/429,695 filed October 28, 1999; and Serial Nos. 09/428,114, 09/428,020, 09/427,859 and 09/427,653 filed October 27, 1999. All these application are incorporated herein by reference.

### FIELD OF THE INVENTION

The present invention relates generally to the fields of genetics and cellular and molecular biology. More particularly, the invention relates to a novel G protein-coupled seven transmembrane receptor polynucleotide and polypeptide sequences that are expressed in the brain.

### DESCRIPTION OF RELATED ART

Humans and other life forms are comprised of living cells. Among the mechanisms through which the cells of an organism communicate with each other and obtain information and stimuli from their environment is through cell membrane receptor molecules expressed on the cell surface. Many such receptors have been identified, characterized, and sometimes classified into major receptor superfamilies based on structural motifs and signal transduction features. Such families include (but are not limited to) ligand-gated ion channel receptors, voltage-dependent ion channel receptors, receptor tyrosine kinases, receptor protein tyrosine phosphatases, and G protein-coupled receptors. The receptors are a first essential link for translating an extracellular signal into a cellular physiological response.

The G protein-coupled receptors (GPCR) form a vast superfamily of cell surface receptors which are characterized by an amino-terminal extracellular domain, a carboxyl-terminal intracellular domain, and a serpentine structure that passes through the cell membrane seven times. Hence, such receptors are sometimes also referred to as seven transmembrane (7TM) receptors. These seven

- 2 -

transmembrane domains define three extracellular loops and three intracellular loops, in addition to the amino- and carboxyl-terminal domains. The extracellular portions of the receptor have a role in recognizing and binding one or more extracellular binding partners (ligands), whereas the intracellular portions have a role in recognizing and communicating with downstream effector molecules.

The G protein-coupled receptors bind a variety of ligands including calcium ions, hormones, chemokines, neuropeptides, neurotransmitters, nucleotides, lipids, odorants, and even photons, and are important in the normal (and sometimes the aberrant) function of many cell types. [See generally A.D. Strosberg, *Eur. J. Biochem.*, 196: 1-10 (1991) and S. K. Bohm *et al.*, *Biochem J.*, 322: 1-18 (1997).]

When a specific ligand binds to its corresponding receptor, the ligand stimulates the receptor to activate a specific heterotrimeric guanine-nucleotide-binding regulatory protein (G-protein) that is coupled to the intracellular portion of the receptor. The G protein in turn transmits a signal to an effector molecule within the cell, by either stimulating or inhibiting the activity of that effector molecule. These effector molecules include adenylate cyclase, phospholipases, and ion channels. Adenylate cyclase and phospholipases are enzymes that are involved in the production of the second messenger molecules cAMP, inositol triphosphate and diacylglycerol. It is through this sequence of events that an extracellular ligand stimuli exerts intracellular changes through a G protein-coupled receptor. Each such receptor has its own characteristic primary structure, expression pattern, ligand-binding profile, and intracellular effector system.

Because of the vital role of G protein-coupled receptors in the communication between cells and their environment, such receptors are attractive targets for therapeutic intervention, and many drugs have been registered which are directed towards activating or antagonizing such receptors. For receptors having a known ligand, the identification of agonists or antagonists may be sought specifically for enhancing or inhibiting the action of the ligand. Some G protein-coupled receptors have roles in disease pathogenesis (e.g., certain chemokine receptors that act as HIV co-receptors and may have a role in AIDS pathogenesis), and are attractive targets for therapeutic intervention even in the absence of knowledge of the natural

- 3 -

ligand of the receptor. Other receptors are attractive targets for therapeutic intervention by virtue of their expression pattern in tissues or cell types that are attractive targets for therapeutic intervention. Examples of this latter category of receptors include receptors expressed in immune cells, for targeting to enhance  
5 immune responses to fight pathogens or cancer or inhibit autoimmune responses; and receptors expressed in the brain or other neurons, for targeting to treat schizophrenia, depression, bipolar disease, or other neurological disorders. This latter category of receptor is also useful as a marker for identifying and/or purifying (*e.g.*, via fluorescence activated cell sorting) cellular subtypes that express the receptor.  
10 Unfortunately, only a limited number of G protein receptors from the central nervous system (CNS) are known. A need exists for identifying the existence and structure of such G protein-coupled receptors.

### SUMMARY OF THE INVENTION

15 The present invention addresses one or more of the needs identified above in that it provides purified polynucleotides encoding heretofore unknown G protein-coupled receptors (GPCR); constructs and recombinant host cells incorporating the polynucleotides; GPCR polypeptides encoded by the polynucleotides; antibodies to the polypeptides; and methods of making and using all  
20 of the foregoing. As set forth in detail herein, the GPCR polypeptides described herein are expressed in the brain, providing a therapeutic indication for GPCR polypeptides and binding partners to treat diseases associated with this tissue.

The invention provides purified and isolated GPCR seven transmembrane receptor polypeptides comprising any one of the amino acid  
25 sequences set forth in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18 or 20, or a fragment thereof comprising an epitope specific to the seven transmembrane receptor. By "epitope specific to" is meant a portion of the receptor that is recognizable by an antibody that is specific for that seven transmembrane receptor, as defined in detail below.

30 One preferred embodiment comprises a purified and isolated polypeptide designated CON193, comprising the complete amino acid sequence set

- 4 -

forth in SEQ ID NO: 2. This amino acid sequence was deduced from a polynucleotide sequence encoding CON193 (SEQ ID NO:1), as set forth below:

```

ntgggtgttg gaccattaaa atgcattatg gaatttttaa aagttggggg agagggagac 60
agtaaaaata acctatattt tctcttggtt tttttttttt aactctagga aagcccagac 120
5 aaattttgag ctatttcata acctaccaga cttatc atg cta aca ctg aat aaa 174
                                     Met Leu Thr Leu Asn Lys
                                     1           5
aca gac cta ata cca gct tca ttt att ctg aat gga gtc cca gga ctg 222
Thr Asp Leu Ile Pro Ala Ser Phe Ile Leu Asn Gly Val Pro Gly Leu
10          10          15          20
gaa gac aca caa ctc tgg att tcc ttc cca ttc tgc tct atg tat gtt 270
Glu Asp Thr Gln Leu Trp Ile Ser Phe Pro Phe Cys Ser Met Tyr Val
          25          30          35
gtg gct atg gta ggg aat tgt gga ctc ctc tac ctc att cac tat gag 318
15 Val Ala Met Val Gly Asn Cys Gly Leu Leu Tyr Leu Ile His Tyr Glu
          40          45          50
gat gcc ctg cac aaa ccc atg tac tac ttc ttg gcc atg ctt tcc ttt 366
Asp Ala Leu His Lys Pro Met Tyr Tyr Phe Leu Ala Met Leu Ser Phe
          55          60          65          70
20 act gac ctt gtt atg tgc tct agt aca atc cct aaa gcc ctc tgc atc 414
Thr Asp Leu Val Met Cys Ser Ser Thr Ile Pro Lys Ala Leu Cys Ile
          75          80          85
ttc tgg ttt cat ctc aag gac att gga ttt gat gaa tgc ctt gtc cag 462
Phe Trp Phe His Leu Lys Asp Ile Gly Phe Asp Glu Cys Leu Val Gln
25          90          95          100
atg ttc ttc atc cac acc ttc aca ggg atg gag tct ggg gtg ctt atg 510
Met Phe Phe Ile His Thr Phe Thr Gly Met Glu Ser Gly Val Leu Met
          105          110          115
30 ctt atg gcc ctg gat cgc tat gtg gcc atc tgc tac ccc tta cgc tat 558
Leu Met Ala Leu Asp Arg Tyr Val Ala Ile Cys Tyr Pro Leu Arg Tyr
          120          125          130
tca act atc ctc acc aat cct gta att gca aag gtt ggg act gcc acc 606
Ser Thr Ile Leu Thr Asn Pro Val Ile Ala Lys Val Gly Thr Ala Thr
135          140          145          150
35 ttc ctg aga ggg gta tta ctc att att ccc ttt act ttc ctc acc aag 654
Phe Leu Arg Gly Val Leu Leu Ile Ile Pro Phe Thr Phe Leu Thr Lys
          155          160          165
cgc ctg ccc tcc tgc aga ggc aat ata ctt ccc cat acc tac tgt gac 702
Arg Leu Pro Ser Cys Arg Gly Asn Ile Leu Pro His Thr Tyr Cys Asp
40          170          175          180

```

- 5 -

```

cac atg tct gta gcc aaa ttg tcc tgt ggt aat gtc aag gtc aat gcc 750
His Met Ser Val Ala Lys Leu Ser Cys Gly Asn Val Lys Val Asn Ala
      185              190              195

atc tat ggt ctg atg gtt gcc ctc ctg att ggg ggc ttt gac ata ctg 798
5 Ile Tyr Gly Leu Met Val Ala Leu Leu Ile Gly Gly Phe Asp Ile Leu
      200              205              210

tgt atc acc atc tcc tat acc atg att ctc cgg gca gtg gtc agc ctc 846
Cys Ile Thr Ile Ser Tyr Thr Met Ile Leu Arg Ala Val Val Ser Leu
      215              220              225              230

10 tcc tca gca gat gct cgg cag aag gcc ttt aat acc tgc act gcc cac 894
Ser Ser Ala Asp Ala Arg Gln Lys Ala Phe Asn Thr Cys Thr Ala His
      235              240              245

att tgt gcc att gtt ttc tcc tat act cca gct ttc ttc tcc ttc ttt 942
Ile Cys Ala Ile Val Phe Ser Tyr Thr Pro Ala Phe Phe Ser Phe Phe
      250              255              260

15 tcc cac cgc ttt ggg gaa cac ata atc ccc cct tct tgc cac atc att 990
Ser His Arg Phe Gly Glu His Ile Ile Pro Pro Ser Cys His Ile Ile
      265              270              275

gta gcc aat att tat ctg ctc cta cca ccc act atg aac cct att gtc 1038
20 Val Ala Asn Ile Tyr Leu Leu Leu Pro Pro Thr Met Asn Pro Ile Val
      280              285              290

tat ggg gtg aaa acc aaa cag ata cga gac tgt gtc ata agg atc ctt 1086
Tyr Gly Val Lys Thr Lys Gln Ile Arg Asp Cys Val Ile Arg Ile Leu
      295              300              305              310

25 tca ggt tct aag gat acc aaa tcc tac agc atg tga atgaacactt 1132
Ser Gly Ser Lys Asp Thr Lys Ser Tyr Ser Met
      315              320

gccaggagtg agaagagaag gaaagaatta cttctatattg cctcttatgc aggagttcat1192
aaaatctttc tggaagtact gtattgatca caaatggag tttgntgact ggtgcattc 1252
30 caataagtac cttgggaatc tnacatcact ggaaggccca ccacatttct ataaat 1308

```

Another preferred embodiment comprises a purified and isolated polypeptide designated CON166, comprising the complete amino acid sequence set forth in SEQ ID NO: 4. This amino acid sequence was deduced from a polynucleotide sequence encoding CON166 (SEQ ID NO: 3), as set forth below:

```

35 atg gat gaa aca gga aat ctg aca gta tct tct gcc aca tgc cat gac 48
Met Asp Glu Thr Gly Asn Leu Thr Val Ser Ser Ala Thr Cys His Asp
      1              5              10              15

act att gat gac ttc cgc aat caa gtg tat tcc acc ttg tac tct atg 96
Thr Ile Asp Asp Phe Arg Asn Gln Val Tyr Ser Thr Leu Tyr Ser Met
40              20              25              30

```

- 6 -

	atc tct gtt gta ggc ttc ttt ggc aat ggc ttt gtg ctc tat gtc ctc	144
	Ile Ser Val Val Gly Phe Phe Gly Asn Gly Phe Val Leu Tyr Val Leu	
	35 40 45	
5	ata aaa acc tat cac aag aag tca gcc ttc caa gta tac atg att aat	192
	Ile Lys Thr Tyr His Lys Lys Ser Ala Phe Gln Val Tyr Met Ile Asn	
	50 55 60	
	tta gca gta gca gat cta ctt tgt gtg tgc aca ctg cct ctc cgt gtg	240
	Leu Ala Val Ala Asp Leu Leu Cys Val Cys Thr Leu Pro Leu Arg Val	
	65 70 75 80	
10	gtc tat tat gtt cac aaa ggc att tgg ctc ttt ggt gac ttc ttg tgc	288
	Val Tyr Tyr Val His Lys Gly Ile Trp Leu Phe Gly Asp Phe Leu Cys	
	85 90 95	
	cgc ctc agc acc tat gct ttg tat gtc aac ctc tat tgt agc atc ttc	336
	Arg Leu Ser Thr Tyr Ala Leu Tyr Val Asn Leu Tyr Cys Ser Ile Phe	
15	100 105 110	
	ttt atg aca gcc atg agc ttt ttc cgg tgc att gca att gtt ttt cca	384
	Phe Met Thr Ala Met Ser Phe Phe Arg Cys Ile Ala Ile Val Phe Pro	
	115 120 125	
20	gtc cag aac att aat ttg gtt aca cag aaa aaa gcc agg ttt gtg tgt	432
	Val Gln Asn Ile Asn Leu Val Thr Gln Lys Lys Ala Arg Phe Val Cys	
	130 135 140	
	gta ggt att tgg att ttt gtg att ttg acc agt tct cca ttt cta atg	480
	Val Gly Ile Trp Ile Phe Val Ile Leu Thr Ser Ser Pro Phe Leu Met	
	145 150 155 160	
25	gcc aaa cca caa aaa gat gag aaa aat aat acc aag tgc ttt gag ccc	528
	Ala Lys Pro Gln Lys Asp Glu Lys Asn Asn Thr Lys Cys Phe Glu Pro	
	165 170 175	
	cca caa gac aat caa act aaa aat cat gtt ttg gtc ttg cat tat gtg	576
	Pro Gln Asp Asn Gln Thr Lys Asn His Val Leu Val Leu His Tyr Val	
30	180 185 190	
	tca ttg ttt gtt ggc ttt atc atc cct ttt gtt att ata att gtc tgt	624
	Ser Leu Phe Val Gly Phe Ile Ile Pro Phe Val Ile Ile Ile Val Cys	
	195 200 205	
35	tac aca atg atc att ttg acc tta cta aaa aaa tca atg aaa aaa aat	672
	Tyr Thr Met Ile Ile Leu Thr Leu Leu Lys Lys Ser Met Lys Lys Asn	
	210 215 220	
	ctg tca agt cat aaa aag gct ata gga atg atc atg gtc gtg acc gct	720
	Leu Ser Ser His Lys Lys Ala Ile Gly Met Ile Met Val Val Thr Ala	
	225 230 235 240	
40	gcc ttt tta gtc agt ttc atg cca tat cat att caa cgt acc att cac	768
	Ala Phe Leu Val Ser Phe Met Pro Tyr His Ile Gln Arg Thr Ile His	
	245 250 255	

- 7 -

```

      ctt cat ttt tta cac aat gaa act aaa ccc tgt gat tct gtc ctt aga   816
      Leu His Phe Leu His Asn Glu Thr Lys Pro Cys Asp Ser Val Leu Arg
                260                      265                      270
5      atg cag aag tcc gtg gtc ata acc ttg tct ctg gct gca tcc aat tgt   864
      Met Gln Lys Ser Val Val Ile Thr Leu Ser Leu Ala Ala Ser Asn Cys
                275                      280                      285
      tgc ttt gac cct ctc cta tat ttc ttt tct ggg ggt aac ttt agg aaa   912
      Cys Phe Asp Pro Leu Leu Tyr Phe Phe Ser Gly Gly Asn Phe Arg Lys
                290                      295                      300
10     agg ctg tct aca ttt aga aag cat tct ttg tcc agc gtg act tat gta   960
      Arg Leu Ser Thr Phe Arg Lys His Ser Leu Ser Ser Val Thr Tyr Val
      305                      310                      315                      320
      ccc aga aag aag gcc tct ttg cca gaa aaa gga gaa gaa ata tgt aaa 1008
      Pro Arg Lys Lys Ala Ser Leu Pro Glu Lys Gly Glu Glu Ile Cys Lys
15     325                      330                      335
      gta tag
      Val
                                           1014

```

Still another preferred embodiment comprises a purified and isolated polypeptide designated CON103, comprising the complete amino acid sequence set forth in SEQ ID NO: 6. This amino acid sequence was deduced from a polynucleotide sequence encoding CON103 (SEQ ID NO: 5), as set forth below:

```

20     ggggcctact tcaccgtgta cccggacttg ggaccatcac agacttcaga accatcagga 60
      acctgggagc aactgaaagc tgaactacag tgggctttca gacacacagc aggtcgcgga 120
      gcacaaatag gactgggtcc ctccaggcca ccagcagggc ggtggagggtc ttactgact 180
25     ccctgcctac ctctcaggac aatgtccttt tggtccaca gtccctgaag ccagagctgg 240
      tgggggcagg gaggcagcca ccagcctcta tatgtagtgg aggaggggggt gtccaggagg 300
      ggctgcatga tctgagagc cccacctca cccggctgga ctatcctccc acttcagggt 360
      ttctctgggc ttccatcttg cccctgctga gccctgcttc ctctctacc agcagcaca 420
      cccccaggct gggctcagag acctcatgtg gtgggatcac tcagtacccc gaggcggagg 480
30     gaaggaggga gggctgcagg gttccccctg gcctgcaaac aggaacacag ggtgtttctc 540
      agtggctgcg agaatgctga tgaaaacccc aggatgttgt gtcaccgtgg tggccagctg 600
      atagtccaa tcatccact ttgccctgag cactcctgca ggggtagaag actccagaac 660
      cttctctcag gcccatggcc caagcagccc atg gaa ctt cat aac ctg agc tct 714
                                           Met Glu Leu His Asn Leu Ser Ser
35     1                      5
      cca tct ccc tct ctc tcc tcc tct gtt ctc cct ccc tcc ttc tct ccc 762
      Pro Ser Pro Ser Leu Ser Ser Ser Val Leu Pro Pro Ser Phe Ser Pro
                10                      15                      20
      tca ccc tcc tct gct ccc tct gcc ttt acc act gtg ggg ggg tcc tct 810
40     Ser Pro Ser Ser Ala Pro Ser Ala Phe Thr Thr Val Gly Gly Ser Ser
      25                      30                      35                      40

```

- 8 -

	gga ggg ccc tgc cac ccc acc tct tcc tgc ctg gtg tct gcc ttc ctg	858
	Gly Gly Pro Cys His Pro Thr Ser Ser Ser Leu Val Ser Ala Phe Leu	
	45 50 55	
5	gca cca atc ctg gcc ctg gag ttt gtc ctg ggc ctg gtg ggg aac agt	906
	Ala Pro Ile Leu Ala Leu Glu Phe Val Leu Gly Leu Val Gly Asn Ser	
	60 65 70	
	ttg gcc ctc ttc atc ttc tgc atc cac acg cgg ccc tgg acc tcc aac	954
	Leu Ala Leu Phe Ile Phe Cys Ile His Thr Arg Pro Trp Thr Ser Asn	
	75 80 85	
10	acg gtg ttc ctg gtc agc ctg gtg gcc gct gac ttc ctc ctg atc agc	1002
	Thr Val Phe Leu Val Ser Leu Val Ala Ala Asp Phe Leu Leu Ile Ser	
	90 95 100	
	aac ctg ccc ctc cgc gtg gac tac tac ctc ctc cat gag acc tgg cgc	1050
	Asn Leu Pro Leu Arg Val Asp Tyr Leu Leu His Glu Thr Trp Arg	
15	105 110 115 120	
	ttt ggg gct gct gcc tgc aaa gtc aac ctc ttc atg ctg tcc acc aac	1098
	Phe Gly Ala Ala Ala Cys Lys Val Asn Leu Phe Met Leu Ser Thr Asn	
	125 130 135	
	cgc acg gcc agc gtt gtc ttc ctc aca gcc atc gca ctc aac cgc tac	1146
20	Arg Thr Ala Ser Val Val Phe Leu Thr Ala Ile Ala Leu Asn Arg Tyr	
	140 145 150	
	ctg aag gtg gtg cag ccc cac cac gtg ctg agc cgt gct tcc gtg ggg	1194
	Leu Lys Val Val Gln Pro His His Val Leu Ser Arg Ala Ser Val Gly	
	155 160 165	
25	gca gct gcc cgg gtg gcc ggg gga ctc tgg gtg ggc atc ctg ctc ctc	1242
	Ala Ala Ala Arg Val Ala Gly Gly Leu Trp Val Gly Ile Leu Leu Leu	
	170 175 180	
	aac ggg cac ctg ctc ctg agc acc ttc tcc ggc ccc tcc tgc ctc agc	1290
	Asn Gly His Leu Leu Leu Ser Thr Phe Ser Gly Pro Ser Cys Leu Ser	
30	185 190 195 200	
	tac agg gtg ggc acg aag ccc tgc gcc tgc ctc cgc tgg cac cag gca	1338
	Tyr Arg Val Gly Thr Lys Pro Ser Ala Ser Leu Arg Trp His Gln Ala	
	205 210 215	
	ctg tac ctg ctg gag ttc ttc ctg cca ctg gcg ctc atc ctc ttt gct	1386
35	Leu Tyr Leu Leu Glu Phe Phe Leu Pro Leu Ala Leu Ile Leu Phe Ala	
	220 225 230	
	att gtg agc att ggg ctc acc atc cgg aac cgt ggt ctg ggc ggg cag	1434
	Ile Val Ser Ile Gly Leu Thr Ile Arg Asn Arg Gly Leu Gly Gly Gln	
	235 240 245	
40	gca ggc ccg cag agg gcc atg cgt gtg ctg gcc atg gtg gtg gcc gtc	1482
	Ala Gly Pro Gln Arg Ala Met Arg Val Leu Ala Met Val Val Ala Val	
	250 255 260	

- 9 -

tac acc atc tgc ttc ttg ccc agc atc atc ttt ggc atg gct tcc atg 1530  
 Tyr Thr Ile Cys Phe Leu Pro Ser Ile Ile Phe Gly Met Ala Ser Met  
 265 270 275 280  
 gtg gct ttc tgg ctg tcc gcc tgc cga tcc ctg gac ctc tgc aca cag 1578  
 5 Val Ala Phe Trp Leu Ser Ala Cys Arg Ser Leu Asp Leu Cys Thr Gln  
 285 290 295  
 ctc ttc cat ggc tcc ctg gcc ttc acc tac ctc aac agt gtc ctg gac 1626  
 Leu Phe His Gly Ser Leu Ala Phe Thr Tyr Leu Asn Ser Val Leu Asp  
 300 305 310  
 10 ccc gtg ctc tac tgc ttc tct agc ccc aac ttc ctc cac cag agc cgg 1674  
 Pro Val Leu Tyr Cys Phe Ser Ser Pro Asn Phe Leu His Gln Ser Arg  
 315 320 325  
 gcc ttg ctg ggc ctc acg cgg ggc cgg cag ggc cca gtg agc gac gag 1722  
 Ala Leu Leu Gly Leu Thr Arg Gly Arg Gln Gly Pro Val Ser Asp Glu  
 15 330 335 340  
 agc tcc tac caa ccc tcc agg cag tgg cgc tac cgg gag gcc tct agg 1770  
 Ser Ser Tyr Gln Pro Ser Arg Gln Trp Arg Tyr Arg Glu Ala Ser Arg  
 345 350 355 360  
 aag gcg gag gcc ata ggg aag ctg aaa gtg cag ggc gag gtc tct ctg 1818  
 20 Lys Ala Glu Ala Ile Gly Lys Leu Lys Val Gln Gly Glu Val Ser Leu  
 365 370 375  
 gaa aag gaa ggc tcc tcc cag ggc tga gggccagctg cagggctgca 1865  
 Glu Lys Glu Gly Ser Ser Gln Gly  
 380 385  
 25 gcgctgtggg ggtaagggct gccgcgtctt ggcttgagg gacaaggcca gcacacgggtg1925  
 cctcaaccaa ctggacaagg gatggcggca gaccaggggc caggccaaag cactggcagg1985  
 actcatgtgg gtggcaggga gagaaacca cctaggcctc tcagtgtgtc caggatggca2045  
 ttcccagaat gcaggggaga gcaggatgcc ggggtggagga gacaggcaag gtgccgttgg2105  
 cacaccagct cagacagggg cctgcgcagc tgcaggggac agacgccaat cactgtcaca2165  
 30 gcagagtcac cttagaaatt ggacagctgc atgttctgtg ctctccagtt tgtcccttcc2225  
 aatattaata aacttccctt ttaaataat tttttgcag accaatatct gtctttaatt2285  
 ctaacctggg actgtcagta ggcgtcaaag tgagcgcccc agtgaaggaa ccttgagag2345  
 agtgggagca ttccagcct tccaggggga ctcgtcttcc agactttgga gcccgcagt2405  
 ctgaagcaga ctctttcttg gtag 2429

35 Another preferred embodiment comprises a purified and isolated  
 polypeptide designated CON203, comprising the complete amino acid sequence set  
 forth in SEQ ID NO: 8. This amino acid sequence was deduced from a  
 polynucleotide sequence encoding CON203 (SEQ ID NO: 7), as set forth below:  
 40 ttgaatttag gtgacactat agaagagcta tgacgtcgca tgcacgcgta cgtaagctcg 60  
 gaattcggct cgagctgaac taatgactgc cgccataaga agacagagag aactgagtat 120

- 10 -

	cctcccaaag	gtgacactgg	aagca	atg	aac	acc	aca	gtg	atg	caa	ggc	ttc	172	
				Met	Asn	Thr	Thr	Val	Met	Gln	Gly	Phe		
				1				5						
5	aac	aga	tct	gag	cgg	tgc	ccc	aga	gac	act	cgg	ata	gta	220
	Asn	Arg	Ser	Glu	Arg	Cys	Pro	Arg	Asp	Thr	Arg	Ile	Val	
	10				15				20				25	
	ttc	cca	gcc	ctc	tac	aca	gtg	gtt	ttc	ttg	acc	ggc	atc	268
	Phe	Pro	Ala	Leu	Tyr	Thr	Val	Val	Phe	Leu	Thr	Gly	Ile	
					30				35				40	
10	act	ttg	gct	ctg	tgg	gtg	ttt	gtt	cac	atc	ccc	agc	tcc	316
	Thr	Leu	Ala	Leu	Trp	Val	Phe	Val	His	Ile	Pro	Ser	Ser	
					45				50				55	
	atc	atc	tac	ctc	aaa	aac	act	ttg	gtg	gcc	gac	ttg	ata	364
	Ile	Ile	Tyr	Leu	Lys	Asn	Thr	Leu	Val	Ala	Asp	Leu	Ile	
15					60				65				70	
	atg	ctt	cct	ttc	aaa	atc	ctc	tct	gac	tca	cac	ctg	gca	412
	Met	Leu	Pro	Phe	Lys	Ile	Leu	Ser	Asp	Ser	His	Leu	Ala	
					75				80				85	
	ctc	aga	gct	ttt	gtg	tgt	cgt	ttt	tct	tcg	gtg	ata	ttt	460
20	Leu	Arg	Ala	Phe	Val	Cys	Arg	Phe	Ser	Ser	Val	Ile	Phe	
					90				95				100	
	atg	tat	gtg	ggc	atc	gtg	ctg	tta	ggg	ctc	ata	gcc	ttt	508
	Met	Tyr	Val	Gly	Ile	Val	Leu	Leu	Gly	Leu	Ile	Ala	Phe	
					110				115				120	
25	ctc	aag	atc	atc	aga	cct	ttg	aga	aat	att	ttt	cta	aaa	556
	Leu	Lys	Ile	Ile	Arg	Pro	Leu	Arg	Asn	Ile	Phe	Leu	Lys	
					125				130				135	
	ttt	gca	aaa	acg	gtc	tca	atc	ttc	atc	tgg	gtc	ttt	ttg	604
	Phe	Ala	Lys	Thr	Val	Ser	Ile	Phe	Ile	Trp	Val	Phe	Leu	
30					140				145				150	
	tcc	ctg	cca	aat	atg	atc	ttg	agc	aac	aag	gaa	gca	aca	652
	Ser	Leu	Pro	Asn	Met	Ile	Leu	Ser	Asn	Lys	Glu	Ala	Thr	
					155				160				165	
	gtg	aaa	aag	tgt	gct	tcc	tta	aag	ggg	cct	ctg	ggg	ctg	700
35	Val	Lys	Lys	Cys	Ala	Ser	Leu	Lys	Gly	Pro	Leu	Gly	Leu	
					170				175				180	
	caa	atg	gta	aat	aac	ata	tgc	cag	ttt	att	ttc	tgg	act	748
	Gln	Met	Val	Asn	Asn	Ile	Cys	Gln	Phe	Ile	Phe	Trp	Thr	
					190				195				200	
40	cta	atg	ctt	gtg	ttt	tat	gtg	gtt	att	gca	aaa	aaa	gta	796
	Leu	Met	Leu	Val	Phe	Tyr	Val	Val	Ile	Ala	Lys	Lys	Val	
					205				210				215	

- 11 -

tat aga aag tcc aaa agt aag gac aga aaa aac aac aaa aag ctg gaa 844  
 Tyr Arg Lys Ser Lys Ser Lys Asp Arg Lys Asn Asn Lys Lys Leu Glu  
 220 225 230  
 ggc aaa gta ttt gtt gtc gtg gct gtc ttc ttt gtg tgt ttt gct cca 892  
 5 Gly Lys Val Phe Val Val Val Ala Val Phe Phe Val Cys Phe Ala Pro  
 235 240 245  
 ttt cat ttt gcc aga gtt cca tat act cac agt caa acc aac aat aag 940  
 Phe His Phe Ala Arg Val Pro Tyr Thr His Ser Gln Thr Asn Asn Lys  
 250 255 260 265  
 10 act gac tgt aga ctg caa aat caa ctg ttt att gct aaa gaa aca act 988  
 Thr Asp Cys Arg Leu Gln Asn Gln Leu Phe Ile Ala Lys Glu Thr Thr  
 270 275 280  
 ctc ttt ttg gca gca act aac att tgt atg gat ccc tta ata tac ata 1036  
 Leu Phe Leu Ala Ala Thr Asn Ile Cys Met Asp Pro Leu Ile Tyr Ile  
 15 285 290 295  
 ttc tta tgt aaa aaa ttc aca gaa aag cta cca tgt atg caa ggg aga 1084  
 Phe Leu Cys Lys Lys Phe Thr Glu Lys Leu Pro Cys Met Gln Gly Arg  
 300 305 310  
 aag acc aca gca tca agc caa gaa aat cat agc agt cag aca gac aac 1132  
 20 Lys Thr Thr Ala Ser Ser Gln Glu Asn His Ser Ser Gln Thr Asp Asn  
 315 320 325  
 ata acc tta ggc tga caactgtaca taggggtaac ttctatttat tgatgagact 1187  
 Ile Thr Leu Gly  
 330  
 25 tccgtagata atgtggaaat caaatttaac caagaaaaaa agattggaac aaatgctctc1247  
 ttacatttta tttatcctgg tgtccaggaa aagattatat taaatttaaa tccacataga1307  
 tctattcata agctgaatga accattacct aagagaatgc aacaggatac caatggccac1367  
 tagaggcata ttctttcttc tttttttttt gttaaatttc aagagcattc actttacatt1427  
 tggaaagact aagggggaacg gttatcctac aaacctccct tcaacacctt ttacatt 1484  
 30 Another preferred embodiment comprises a purified and isolated  
 polypeptide designated CON198, comprising the complete amino acid sequence set  
 forth in SEQ ID NO: 10. This amino acid sequence was deduced from a  
 polynucleotide sequence encoding CON198 (SEQ ID NO: 9), as set forth below:  
 35 atg atg gtg gat ccc aat ggc aat gaa tcc agt gct aca tac ttc atc 48  
 Met Met Val Asp Pro Asn Gly Asn Glu Ser Ser Ala Thr Tyr Phe Ile  
 1 5 10 15  
 cta ata ggc ctc cct ggt tta gaa gag gct cag ttc tgg ttg gcc ttc 96  
 40 Leu Ile Gly Leu Pro Gly Leu Glu Glu Ala Gln Phe Trp Leu Ala Phe  
 20 25 30  
 cca ttg tgc tcc ctc tac ctt att gct gtg cta ggt aac ttg aca atc 144  
 Pro Leu Cys Ser Leu Tyr Leu Ile Ala Val Leu Gly Asn Leu Thr Ile  
 35 40 45  
 45

- 12 -

	atc tac att gtg cgg act gag cac agc ctg cat gag ccc atg tat ata	192
	Ile Tyr Ile Val Arg Thr Glu His Ser Leu His Glu Pro Met Tyr Ile	
	50 55 60	
5	ttt ctt tgc atg ctt tca ggc att gac atc ctc atc tcc acc tca tcc	240
	Phe Leu Cys Met Leu Ser Gly Ile Asp Ile Leu Ile Ser Thr Ser Ser	
	65 70 75 80	
10	atg ccc aaa atg ctg gcc atc ttc tgg ttc aat tcc act acc atc cag	288
	Met Pro Lys Met Leu Ala Ile Phe Trp Phe Asn Ser Thr Thr Ile Gln	
	85 90 95	
15	ttt gat gct tgt ctg cta cag atg ttt gcc atc cac tcc tta tct ggc	336
	Phe Asp Ala Cys Leu Leu Gln Met Phe Ala Ile His Ser Leu Ser Gly	
	100 105 110	
20	atg gaa tcc aca gtg ctg ctg gcc atg gct ttt gac cgc tat gtg gcc	384
	Met Glu Ser Thr Val Leu Leu Ala Met Ala Phe Asp Arg Tyr Val Ala	
	115 120 125	
25	atc tgt cac cca ctg cgc cat gcc aca gta ctt acg ttg cct cgt gtc	432
	Ile Cys His Pro Leu Arg His Ala Thr Val Leu Thr Leu Pro Arg Val	
	130 135 140	
30	acc aaa att ggt gtg gct gct gtg gtg cgg ggg gct gca ctg atg gca	480
	Thr Lys Ile Gly Val Ala Val Val Arg Gly Ala Ala Leu Met Ala	
	145 150 155 160	
35	ccc ctt cct gtc ttc atc aag cag ctg ccc ttc tgc cgc tcc aat atc	528
	Pro Leu Pro Val Phe Ile Lys Gln Leu Pro Phe Cys Arg Ser Asn Ile	
	165 170 175	
40	ctt tcc cat tcc tac tgc cta cac caa gat gtc atg aag ctg gcc tgt	576
	Leu Ser His Ser Tyr Cys Leu His Gln Asp Val Met Lys Leu Ala Cys	
	180 185 190	
45	gat gat atc cgg gtc aat gtc gtc tat ggc ctt atc gtc atc atc tcc	624
	Asp Asp Ile Arg Val Asn Val Val Tyr Gly Leu Ile Val Ile Ile Ser	
	195 200 205	
50	ggc att ggc ctg gac tca ctt ctc atc tcc ttc tca tat ctg ctt att	672
	Ala Ile Gly Leu Asp Ser Leu Leu Ile Ser Phe Ser Tyr Leu Leu Ile	
	210 215 220	
55	ctt aag act gtg ttg ggc ttg aca cgt gaa gcc cag gcc aag gca ttt	720
	Leu Lys Thr Val Leu Gly Leu Thr Arg Glu Ala Gln Ala Lys Ala Phe	
	225 230 235 240	
60	ggc act tgc gtc tct cat gtg tgt gct gtg ttc ata ttc tat gta cct	768
	Gly Thr Cys Val Ser His Val Cys Ala Val Phe Ile Phe Tyr Val Pro	
	245 250 255	
65	ttc att gga ttg tcc atg gtg cat cgc ttt agc aag cgg cgt gac tct	816
	Phe Ile Gly Leu Ser Met Val His Arg Phe Ser Lys Arg Arg Asp Ser	
	260 265 270	
70	cgg ctg ccc gtc atc ttg gcc aat atc tat ctg ctg gtt cct cct gtg	864
	Pro Leu Pro Val Ile Leu Ala Asn Ile Tyr Leu Leu Val Pro Pro Val	
	275 280 285	
75	ctc aac cca att gtc tat gga gtg aag aca aag gag att cga cag cgc	912
	Leu Asn Pro Ile Val Tyr Gly Val Lys Thr Lys Glu Ile Arg Gln Arg	
	290 295 300	

- 13 -

atc ctt cga ctt ttc cat gtg gcc aca cac gct tca gag ccc tag 957  
 Ile Leu Arg Leu Phe His Val Ala Thr His Ala Ser Glu Pro  
 305 310 315

It will be appreciated that SEQ ID NO: 10 contains methionine residues at positions 1 and 2. Translation of the relevant mRNA sequences may occur beginning from either or both methionines, which can be determined for a particular cell source by purifying expressed CON198 protein and performing amino-terminal sequencing thereon. CON198 polypeptides beginning at either Met<sub>1</sub> or Met<sub>2</sub> of SEQ ID NO: 10 are intended a polypeptides of the invention.

Another preferred embodiment comprises a purified and isolated polypeptide designated CON197, comprising the complete amino acid sequence set forth in SEQ ID NO: 12. This amino acid sequence was deduced from a polynucleotide sequence encoding CON197 (SEQ ID NO: 11), as set forth below:

1  
 15 ATGGAAGCGAGAACAGAAGAGTGATAAGAGAATTCATCCTCCTGGTCTGACCCAGTCTCAAGATATT  
 M E S E N R R V I R E F I L L G L T Q S Q D I

70  
 20 CAGCTCCTGGTCTTTGTGCTAGTTTTAATATTCTACTTCATCATCCTCCCTGGAAATTTCTCATTATT  
 Q L L V F V L V L I F Y F I I L P G N F L I I

139  
 25 TTCACCATAAAGTCAGACCCTGGGCTCACAGCCCCCTCTATTCTTTCTGGGCAACTTGGCCTTCCTG  
 F T I K S D P G L T A P L Y F F L G N L A F L

208  
 30 GATGCATCCTACTCCTTCATTGTGGCTCCCCGGATGTTGGTGGACTTCTCTCTGCGAAGAAGATAATC  
 D A S Y S F I V A P R M L V D F L S A K K I I

277  
 35 TCCTACAGAGGCTGCATCACTCAGCTCTTTTCTTGCACTTCCTTGGAGGAGGGGAGGGATTACTCCTT  
 S Y R G C I T Q L F F L H F L G G G E G L L L

346  
 40 GTTGTGATGGCCTTTGACCGCTACATCGCCATCTGCCGGCCTCTGCACTATCCTACTGTGATGAACCT  
 V V M A F D R Y I A I C R P L H Y P T V M N P

415  
 45 AGAACCTGCTATGCAATGATGTTGGCTCTGTGGCTTGGGGGTTTGTCCACTCCATTATCCAGGTGGTC  
 R T C Y A M M L A L W L G G F V H S I I Q V V

484  
 50 CTCATCCTCCGCTTGCCTTTTGTGGCCCAACCAGCTGGACAACCTTCTTCTGTGATGTCCACAGGTC  
 L I L R L P F C G P N Q L D N F F C D V P Q V

553  
 ATCAAGCTGGCCTGCACCGACACATTGTGGTGGAGCTTCTGATGGTCTTCAACAGTGGCCTGATGACA  
 I K L A C T D T F V V E L L M V F N S G L M T

622  
 CTCCTGTGCTTTCTGGGGCTTCTGGCCTCCTATGCAGTCATTCTTTGTGCGATACGAGGGTCTTCTTCT  
 L L C F L G L L A S Y A V I L C R I R G S S S

- 14 -

691  
GAGGCAAAAAACAAGGCCATGTCCACGTGCATCACCCATATCATTTGTTATATTCTTCATGTTTGGACCT  
E A K N K A M S T C I T H I I V I F F M F G P

5 760  
GGCATCTTCATCTACACGCGCCCTTCAGGGCTTTCCCAGCTGACAAGGTGGTTTCTCTCTTCCACACA  
G I F I Y T R P F R A F P A D K V V S L F H T

829  
10 GTGATTTTTCCTTTGTTGAATCCTGTCAATTTATACCTTCGCAACCAGGAAGTGAAAGCTTCCATGAAA  
V I F P L L N P V I Y T L R N Q E V K A S M K

898  
15 AAGGTGTTTAATAAGCACATAGCCTGAAAAAGGGCGCAAAAAAAAAAGAATAAAATAGACTGTAGAA  
K V F N K H I A \*

967  
TTTTTAAAAAAAAAAAAAAAAAAAAA

Another preferred embodiment comprises a purified and isolated

20 polypeptide designated CON202, comprising the complete amino acid sequence set  
forth in SEQ ID NO: 14. This amino acid sequence was deduced from a  
polynucleotide sequence encoding CON202 (SEQ ID NO: 13), as set forth below:

1  
25 TGCTTCCCCATAAGGTAACAGCTTTGTTAGCNCCTGTCTGACATCATTGCTTGTNACTTAAGAACTGAT  
70  
AGGTNTTTTTTTTTTTTTTTTTTCAGATATTCTGATGGCAAAACAAGTGGAAGAAAAGAGGAAGCATGA

139  
30 CTGCAGATCAGATCAGTTCTCTTTGTGGATTATATTTTCAGTAAATGTATGGATCTATCTTTTCCTTG  
208  
TTCTTATATCTAGATCATGAGACTTGACTGAGGCTGTATCCTTATCCTCCATCCATCTATGGCGAACTA  
M A N Y

35 277  
TAGCCATGCAGCTGACAACATTTTGCAAAATCTCTCGCCTCTAACAGCCTTTCTGAAACTGACTTCCTT  
S H A A D N I L Q N L S P L T A F L K L T S L

40 346  
GGGTTTCATAATAGGAGTCAGCGTGGTGGGCAACCTCCTGATCTCCATTTTGCTAGTGAAAGATAAGAC  
G F I I G V S V V G N L L I S I L L V K D K T

415  
45 CTTGCATAGAGCACCTTACTACTTCCTGTTGGATCTTTGCTGTTTCAGATATCCTCAGATCTGCAATTTG  
L H R A P Y Y F L L D L C C S D I L R S A I C

484  
50 TTTCCCATTTGTGTTCAACTCTGTCAAAAATGGTTCTACCTGGACTTATGGGACTCTGACTTGCAAAGT  
F P F V F N S V K N G S T W T Y G T L T C K V

553  
GATTGCCTTTCTGGGGGTTTTGTCTGTTTCCACACTGCTTTTCATGCTCTTCTGCATCAGTGTCAACAG  
I A F L G V L S C F H T A F M L F C I S V T R

622  
ATATTTAGCTATCGCCCATCACCGCTTCTATACAAAGAGGCTGACCTTTTGGACGTGTCTGGCTGTGAT  
Y L A I A H H R F Y T K R L T F W T C L A V I

- 15 -

691  
 CTGTATGGTGTGGACTCTGTCTGTGGCCATGGCATTTCCTCCCGGTTTTAGACGTGGGCACTTACTCATT  
 C M V W T L S V A M A F P P V L D V G T Y S F

5 760  
 CATTAGGGAGGAAGATCAATGCACCTTCCAACACCGCTCCTTCAGGGCTAATGATTTCCTAGAAATTTAT  
 I R E E D Q C T F Q H R S F R A N D S L G F M

10 829  
 GCTGCTTCTGTCTCTCATCCTCCTAGCCACACAGCTTGTCTACCTCAAGCTGATATTTTTTCGTCCACGA  
 L L L A L I L L A T Q L V Y L K L I F F V H D

15 898  
 TCGAAGAAAAATGAAGCCAGTCCAGTTTGTAGCAGCAGTCAGCCAGAAGTGGACTTTTCATGGTCTCTGG  
 R R K M K P V Q F V A A V S Q N W T F H G P G

20 967  
 AGCCAGTGGCCAGGAGCTGCCAATTGGCTAGCAGGATTTGGAAGGGGTCCCACACCACCCACCTTGCT  
 A S G Q A A A N W L A G F G R G P T P P T L L

25 1036  
 GGGCATCAGGCAAAATGCAAACACCACAGGCAGAAGAAGGCTATTGGTCTTAGACGAGTTCAAATGGA  
 G I R Q N A N T T G R R R L L V L D E F K M E

30 1105  
 GAAAAGATCAGCAGAATGTTCTATATAATGACTTTTCTGTTTCTAACCTTGTGGGGCCCTACCTGGT  
 K R I S R M F Y I M T F L F L T L W G P Y L V

35 1174  
 GGCCTGTTATTGGAGAGTTTTTGCAAGAGGGCCTGTAGTACCAGGGGGATTCTAACAGCTGCTGTCTG  
 A C Y W R V F A R G P V V P G G F L T A A V W

40 1243  
 GATGAGTTTGGCCCAAGCAGGAATCAATCCTTTTGTCTGCATTTTCTCAAACAGGGAGCTGAGGCGCTG  
 M S F A Q A G I N P F V C I F S N R E L R R C

45 1312  
 TTTCAGCACAAACCTTCTTTACTGCAGAAAATCCAGGTTACCAAGGGAACCTTACTGTGTTATATGAGG  
 F S T T L L Y C R K S R L P R E P Y C V I

Still another preferred embodiment comprises a purified and isolated  
 polypeptide designated CON222, comprising the complete amino acid sequence set  
 forth in SEQ ID NO: 16. This amino acid sequence was deduced from a  
 polynucleotide sequence encoding CON222 (SEQ ID NO: 15), as set forth below:

1 ATGTTTAGACCTCTGTGAATCTCTCTCACATATATTTTAAGAAATCCAGTACTGTGGGTATGCA  
 M F R P L V N L S H I Y F K K F Q Y C G Y A

67 CCACATGTTGCGAGCTGTAAACCAAACTGATGGAATTTTCATCTCTAGAGAATCTCTGGCAAGC  
 P H V R S C K P N T D G I S S L E N L L A S

133 ATTATTACAGAGATATTGTCTGGGTTGTATCTGCAGTTACCTGCTTTGGAAACATTTTGTTCATT  
 I I Q R V F V W V V S A V T C F G N I F V I

199 TGCATGCGACCTTATATCAGGTCTGAGAACAAGCTGTATGCCATGTCAATCATTTCTCTCTGCTGT  
 C M R P Y I R S E N K L Y A M S I I S L C C

265 GCCGACTGCTTAATGGGAATATATTTATTTCGTGATCGGAGGCTTTGACCTAAAGTTTCGTGGAGAA  
 A D C L M G I Y L F V I G G F D L K F R G E

331 TACAATAAGCATGCGAGCTGTGGATGGAGAGTACTCATTGTGCAGCTGTAGATCTTTGGCCATT  
Y N K H A Q L W M E S T H C Q L V G S L A I  
397 CTGTCCACAGAAGTATCAGTTTTACTGTTAACATTTCTGACATTGGAAAAATACATCTGCATTGTC  
L S T E V S V L L L T F L T L E K Y I C I V  
5 463 TATCCTTTTATAGATGTGTGAGACCTGGAAAAATGCAGAACATTACAGTTCTGATTCTCATTGGATT  
Y P F R C V R P G K C R T I T V L I L I W I  
529 ACTGGTTTTATAGTGGCTTTCATTCCATTGAGCAATAAGGAATTTTCAA AAACTACTATGGCACC  
T G F I V A F I P L S N K E F F K N Y Y G T  
595 AATGGAGTATGCTTCCCTCTTCATTGAGAAGATACAGAAAGTATTGGAGCCCAGATTATTTCAGTG  
N G V C F P L H S E D T E S I G A Q I Y S V  
10 661 GCAATTTTTCTTGGTATTAATTGGCCGCATTTATCATCATAGTTTTTTCCTATGGAAGCATGTTT  
A I F L G I N L A A F I I I V F S Y G S M F  
727 TATAGTGTTCATCAAAGTGCCATAACAGCAACTGAAATACGGAATCAAGTTAAAAAGAGATGATC  
Y S V H Q S A I T A T E I R N Q V K K E M I  
15 793 CTTGCCAAACGTTTTTCTTTATAGTATTTACTGATGCATTATGCTGGATACCCATTTTTGTAGTG  
L A K R F F F I V F T D A L C W I P I F V V  
859 AAATTTCTTTCACTGCTTCAGGTAGAAATACCAGGTACCATAACCTCTGGGTAGTGATTTTTATT  
K F L S L L Q V E I P G T I T S W V V I F I  
20 925 CTGCCCATTAACAGTGCTTTGAACCCAATTCTCTATACTCTGACCACAAGACCATTTAAAGAAATG  
L P I N S A L N P I L Y T L T T R P F K E M  
991 ATTCATCGGTTTTTGGTATAACTACAGACAAAGAAAATCTATGGACAGCAAAGGTCAGAAAACATAT  
I H R F W Y N Y R Q R K S M D S K G Q K T Y  
1057 GCTCCATCATTCATCTGGGTGGAATGTGGCCACTGCAGGAGATGCCACCTGAGTTAATGAAGCCG  
A P S F I W V E M W P L Q E M P P E L M K P 1123  
25 GACCTTTTCACATACCCCTGTGAAATGTCACTGATTTCTCAATCAACGAGACTCAATTCCATTCA  
D L F T Y P C E M S L I S Q S T R L N S Y S  
1189 TGA 1191

Another preferred embodiment comprises a purified and isolated

30 polypeptide designated CON215, comprising the complete amino acid sequence set forth in SEQ ID NO: 18. This amino acid sequence was deduced from a polynucleotide sequence encoding CON215 (SEQ ID NO: 17), as set forth below:

35	atg Met 1	ggg Gly	ttc Phe	aac Asn	ttg Leu 5	acg Thr	ctt Leu	gca Ala	aaa Lys	tta Leu 10	cca Pro	aat Asn	aac Asn	gag Glu	ctg Leu 15	cac His	48
40	ggc Gly	caa Gln	gag Glu	agt Ser 20	cac His	aat Asn	tca Ser	ggc Gly	aac Asn 25	agg Arg	agc Ser	gac Asp	ggg Gly	cca Pro 30	gga Gly	aag Lys	96
	aac Asn	acc Thr	acc Thr 35	ctt Leu	cac His	aat Asn	gaa Glu	ttt Phe 40	gac Asp	aca Thr	att Ile	gtc Val	ttg Leu 45	cca Pro	gtg Val	ctt Leu	144
45	tat Tyr 50	ctc Leu	att Ile	ata Ile	ttt Phe	gtg Val	gca Ala 55	agc Ser	atc Ile	ttg Leu	ctg Leu	aat Asn 60	ggg Gly	tta Leu	gca Ala	gtg Val	192

- 17 -

	tgg atc ttc ttc cac att agg aat aaa acc agc ttc ata ttc tat ctc	240
	Trp Ile Phe Phe His Ile Arg Asn Lys Thr Ser Phe Ile Phe Tyr Leu	
	65 70 75 80	
5	aaa aac ata gtg gtt gca gac ctc ata atg acg ctg aca ttt cca ttt	288
	Lys Asn Ile Val Val Ala Asp Leu Ile Met Thr Leu Thr Phe Pro Phe	
	85 90 95	
10	cga ata gtc cat gat gca gga ttt gga cct tgg tac ttc aag ttt att	336
	Arg Ile Val His Asp Ala Gly Phe Gly Pro Trp Tyr Phe Lys Phe Ile	
	100 105 110	
15	ctc tgc aga tac act tca gtt ttg ttt tat gca aac atg tat act tcc	384
	Leu Cys Arg Tyr Thr Ser Val Leu Phe Tyr Ala Asn Met Tyr Thr Ser	
	115 120 125	
20	atc gtg ttc ctt ggg ctg ata agc att gat cgc tat ctg aag gtg gtc	432
	Ile Val Phe Leu Gly Leu Ile Ser Ile Asp Arg Tyr Leu Lys Val Val	
	130 135 140	
	aag cca ttt ggg gac tct cgg atg tac agc ata acc ttc acg aag gtt	480
	Lys Pro Phe Gly Asp Ser Arg Met Tyr Ser Ile Thr Phe Thr Lys Val	
	145 150 155 160	
25	tta tct gtt tgt gtt tgg gtg atc atg gct gtt ttg tct ttg cca aac	528
	Leu Ser Val Cys Val Trp Val Ile Met Ala Val Leu Ser Leu Pro Asn	
	165 170 175	
30	atc atc ctg aca aat ggt cag cca aca gag gac aat atc cat gac tgc	576
	Ile Ile Leu Thr Asn Gly Gln Pro Thr Glu Asp Asn Ile His Asp Cys	
	180 185 190	
35	tca aaa ctt aaa agt cct ttg ggg gtc aaa tgg cat acg gca gtc acc	624
	Ser Lys Leu Lys Ser Pro Leu Gly Val Lys Trp His Thr Ala Val Thr	
	195 200 205	
40	tat gtg aac agc tgc ttg ttt gtg gcc gtg ctg gtg att ctg atc gga	672
	Tyr Val Asn Ser Cys Leu Phe Val Ala Val Leu Val Ile Leu Ile Gly	
	210 215 220	
	tgt tac ata gcc ata tcc agg tac atc cac aaa tcc agc agg caa ttc	720
	Cys Tyr Ile Ala Ile Ser Arg Tyr Ile His Lys Ser Ser Arg Gln Phe	
	225 230 235 240	
45	ata agt cag tca agc cga aag cga aaa cat aac cag agc atc agg gtt	768
	Ile Ser Gln Ser Ser Arg Lys Arg Lys His Asn Gln Ser Ile Arg Val	
	245 250 255	
50	gtt gtg gct gtg ttt ttt acc tgc ttt cta cca tat cac ttg tgc aga	816
	Val Val Ala Val Phe Phe Thr Cys Phe Leu Pro Tyr His Leu Cys Arg	
	260 265 270	
55	att cct ttt act ttt agt cac tta gac agg ctt tta gat gaa tct gca	864
	Ile Pro Phe Thr Phe Ser His Leu Asp Arg Leu Leu Asp Glu Ser Ala	
	275 280 285	
60	caa aaa atc cta tat tac tgc aaa gaa att aca ctt ttc ttg tct gcg	912
	Gln Lys Ile Leu Tyr Tyr Cys Lys Glu Ile Thr Leu Phe Leu Ser Ala	
	290 295 300	

- 18 -

```

          tgt aat gtt tgc ctg gat cca ata att tac ttt ttc atg tgt agg tca   960
          Cys Asn Val Cys Leu Asp Pro Ile Ile Tyr Phe Phe Met Cys Arg Ser
          305                      310                      315                      320

5      ttt tca aga agg ctg ttc aaa aaa tca aat atc aga acc agg agt gaa   1008
          Phe Ser Arg Arg Leu Phe Lys Lys Ser Asn Ile Arg Thr Arg Ser Glu
                      325                      330                      335

          agc atc aga tca ctg caa agt gtg aga aga tcg gaa gtt ctc ata tat   1056
          Ser Ile Arg Ser Leu Gln Ser Val Arg Arg Ser Glu Val Leu Ile Tyr
                      340                      345                      350

          tat gat tat act gat gtg tag   1077
          Tyr Asp Tyr Thr Asp Val
          355

```

Another preferred embodiment comprises a purified and isolated polypeptide designated CON217, comprising the complete amino acid sequence set forth in SEQ ID NO: 20. This amino acid sequence was deduced from a polynucleotide sequence encoding CON217 (SEQ ID NO: 19), as set forth below:

```

20      -41                      C ATGGCATCCC CAGCCTAGCT CCCAATCCCA CTTTGGCAGC
          1 ATGTTAGCCAACAGCTCCTCAACCAACAGTTCTGTCTCTCCCGTGTCTGACTACCGACCTACCCAC
            M L A N S S S T N S S V L P C P D Y R P T H
          67 CGCCTGCACTTGGTGGTCTACAGCTTGGTGTGGCTGCCGGGCTCCCCCTCAACGCGCTAGCCCTC
            R L H L V V Y S L V L A A G L P L N A L A L
25      133 TGGGTCTTCTGCGCGCGCTGCGCGTCACTCGGTGGTGAGCGTGACATGTGTAACCTGGCGGCC
            W V F L R A L R V H S V V S V Y M C N L A A
          199 AGCGACCTGCTCTTACCCTCTCGCTGCCCGTTCGTCTCTCTACTACGCACTGCACCACTGGCCCC
            S D L L F T L S L P V R L S Y Y A L H H W P
          265 TTCCCCGACCTCCTGTGCCAGACGACGGCGCCATCTTCCAGATGAACATGTACGGCAGCTGCATC
            F P D L L C Q T T G A I F Q M N M Y G S C I
30      331 TTCCTGATGCTCATCAACGTGGACCGCTACGCCGCCATCGTGCACCCGCTGCGACTGCGCCACCTG
            F L M L I N V D R Y A A I V H P L R L R H L
          397 CGGCGGCCCCGCGTGGCGCGGCTGCTCTGCTGGGCGTGTGGGCGCTCATCCTGGTGTGTTGCCGTG
            R R P R V A R L L C L G V W A L I L V F A V
35      463 CCGCGCGCCCGCGTGCACAGGCCCTCGCGTTGCCGCTACCGGGACCTCGAGGTGCGCCTATGCTTC
            P A A R V H R P S R C R Y R D L E V R L C F
          529 GAGAGCTTCAGCGACGAGCTGTGAAAGGCAGGCTGCTGCCCTCGTGCTGCTGGCCGAGGCGCTG
            E S F S D E L W K G R L L P L V L L A E A L
          595 GGCTTCCTGCTGCCCCTGGCGGCGGTGCTACTCGTCGGGCCGAGTCTTCTGGACGCTGGCGCGC
            G F L L P L A A V V Y S S G R V F W T L A R
40      661 CCCGACGCCACGCAGAGCCAGCGGCGGGAAGACCGTGCGCCTCCTGCTGGCTAACCTCGTCATC
            P D A T Q S Q R R R K T V R L L L A N L V I
          727 TTCCTGCTGTGCTTCGTGCCCTACAACAGCACGCTGGCGGTCTACGGGCTGCTGCGGAGCAAGCTG
            F L L C F V P Y N S T L A V Y G L L R S K L
45      793 GTGGCGGCCAGCGTGCTGCCCGCATCGCGTGCGCGGGGTGCTGATGGTGATGGTGCTGCTGGCC
            V A A S V P A R D R V R G V L M V M V L L A

```

- 19 -

859 GGCGCCAACTGCGTGCTGGACCCGCTGGTGTACTACTTTAGCCCGAGGGCTTCCGCAACACCCTG  
 G A N C V L D P L V Y Y F S A E G F R N T L  
 925 CGCGGCTGGGCACTCCGCACCGGGCCAGGACCTCGGCCACCAACGGGACGCGGGCGGCGCTCGCG  
 R G L G T P H R A R T S A T N G T R A A L A  
 5 991 CAATCCGAAAGGTCCGCCGTCAACACGACGCCACCAGGCCGGATGCCGCCAGTCAGGGGCTGCTC  
 Q S E R S A V T T D A T R P D A A S Q G L L  
 1057 CGACCCTCCGACTCCCACTCTCTGTCTTCCTTCACACAGTGTCCCCAGGATTCGCCCTCTGAACA  
 R P S D S H S L S S F T Q C P Q D S A L \*  
 1123 CACATGCCAT TGCCTGTCC GTGCCCAGCT CCCAACGCCT CTCGTTCTGG GAGGCTTACA  
 10 1183 GGGTGTACAC ACAAGAAGGT GGGCTGGGCA CTTGGACCTT TGGGTGGCAA TTCCAGCTTA  
 1243 GCAACGCAGA AGAGTACAAA GTGTGGAAGC CAGGGCCCAG GGAAGGCAGT GCTGCTGGAA  
 1303 ATGGCTTCTT TAAACTGTGA GCACGCAGAG CACCCCTTCT CCAGCGGTGG GAAGTGATGC  
 1363 AGAGAGCCCA CCCGTGCAGA GGGCAGAAGA GGACGAAATG CCTTTGGGTG GGCAGGGCAT  
 1423 TAAACTGCTA AAAGCTGGTT AGATGGAACA GAAATGGGC ATTCTGGATC TAAACCCCA  
 15 1483 CAGGGGCTG AGAGCTGAAG AGCACCAGGT TTGGTGGACA AAGCTACTGA GATGCTGTGT  
 1543 CATCTGTGA CTTCTGTCTA GGCTCATGGA TGCCACCCCC TTTCATTTCG GCCTAGGCTT  
 1603 CCCCTGTCTA CCACTGAGGC CTAATACAAG AGTTCTATG GACAGAACTA CATTCTTCT  
 1663 CGCATAGTGA CTTGTGACAA TTTAGACTTG GCATCCAGCA TGGGATAGTT GGGGCAAGGC  
 1723 AAAACTAACT TAGAGTTTCC CCCTCAACAA CATCCAAGTC CAAACCCCTT TTAGGTTATC  
 20 1783 CTTTCTTCCA TCACATCCCC TTTTCCAGGC CTCCTCCATT TTAGGTCCTT AATATTCTTT  
 1843 CTTTTTCTCT CTCTCTCGTT TCTCTTCTCT CTCTCTCTC CTCTCTCTC TCTTCTCTC  
 1903 TTCTCTCTCT CTCCCTCTCT CTCCTTTGTC CAGAGTAAGG ATAAAATTCT TTCTACTAAA  
 1963 GCACTGGTTC TCAAACTTTT TGGTCTCAGA CCCCACTCTT AGAAATTGAG GATCTCAAAG  
 2023 AGCTTTGCTT ATATTTTGTT CTTTGTGATC TTACCATACT AGAAATTAAA GCGAATACAT  
 25 2083 TTTTAAAATA AATACACATG CACACATTAC ATTAGCCATG GGAGCAATAA TGTCACCACA  
 2143 CCACTTCAT GAAGCCTCTG GAAACTCTA CAGTATACTT GTGAGAGAAT GAGAGTGAAA  
 2203 GGGACAAATA ACATCTGTGT AGCAGTATTA TGAAAATAGC TTGACCTGTG GCACTTCCTC  
 2263 AGAGGGTGG TCCCTGGATC ACACTTTGTG AACCATACTT GTCCTGAAGT ATTGGAGTTC  
 2323 ATGTCTAACT TCTTCCAGG GCATTATGTA CAGTGCTTTT TATTACTGTG GGGAGAGGGC  
 30 2383 AGTGCTAAAT AAATTAATCA CTAAGTATAA AAAAAAAAAA AAAAAAAAAA AAAAAA

Although SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, and 20 provide  
 for particular human sequences, the invention is intended to include within its scope  
 other human allelic variants; non-human mammalian forms of GPCR polypeptides,  
 and other vertebrate forms of GPCR polypeptides.

It will be appreciated that extracellular epitopes are particularly useful  
 for generating and screening for antibodies and other binding compounds that bind to  
 receptors such as GPCR polypeptides. Thus, in another preferred embodiment, the  
 invention provides a purified and isolated polypeptide comprising at least one  
 extracellular domain of a GPCR polypeptide of the invention. By "extracellular  
 domain", is it meant the amino terminal extracellular domain or an extracellular loop  
 that spans two membrane domains.

- 20 -

A purified and isolated polypeptide comprising the N-terminal extracellular domain of GPCR polypeptides of the invention is highly preferred. Also preferred is a purified and isolated polypeptide comprising a GPCR seven transmembrane receptor fragment selected from the group consisting of the N-terminal extracellular domain of GPCR polypeptides of the invention, transmembrane domains of GPCR polypeptides of the invention, extracellular loops connecting transmembrane domains of GPCR polypeptides of the invention, intracellular loops connecting transmembrane domains of GPCR polypeptides of the invention, the C-terminal cytoplasmic domain of GPCR polypeptides, and fusions thereof. Such fragments may be continuous portions of the native receptor. However, it will also be appreciated that knowledge of the GPCR gene and protein sequences as provided herein permits recombining of various domains that are not contiguous in the native protein.

In another embodiment, the invention provides purified and isolated polynucleotides (*e.g.*, cDNA, genomic DNA, synthetic DNA, RNA, or combinations thereof, single or double stranded) that comprise a nucleotide sequence encoding an amino acid sequence of the polypeptides of the invention. Another embodiment provides a purified and isolated polynucleotide encoding the amino acid sequence of the polypeptide of the invention fused to a heterologous tag amino acid sequence. Such polynucleotides are useful for recombinantly expressing the receptor and also for detecting expression of the receptor in cells (*e.g.*, using Northern hybridization and *in situ* hybridization assays, and Western studies). Polynucleotides encoding polypeptides of the invention also are useful to design antisense and other molecules for the suppression of GPCR polypeptides expression in a cultured cell or animal (for therapeutic purposes or to provide a model for diseases characterized by aberrant GPCR polypeptide expression). Such polynucleotides are also useful to design antisense and other molecules for the suppression of GPCR polypeptide expression in a cultured cell or tissue or in an animal, for therapeutic purposes or to provide a model for diseases characterized by aberrant GPCR polypeptide expression. Specifically excluded from the definition of polynucleotides of the invention are entire isolated chromosomes of native host cells. A preferred polynucleotide set forth in any one of

- 21 -

the SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, and 19 corresponds to a naturally occurring GPCR sequence. It will be appreciated that numerous other sequences exist that also encode GPCR polypeptides having the amino acid sequence set out in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18 and 20 due to the well-known degeneracy of the universal genetic code. All such sequences represent polynucleotides of the invention.

The invention also provides a purified and isolated polynucleotide comprising a nucleotide sequence that encodes a mammalian seven transmembrane receptor, wherein the polynucleotide hybridizes to a nucleotide sequence set forth in any one of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, or 19 or the non-coding strand complementary thereto, under the following hybridization conditions:

- (a) hybridization for 16 hours at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% Dextran sulphate; and
- (b) washing 2 times for 30 minutes at 60°C in a wash solution comprising 0.1% SSC, 1% SDS. Polynucleotides that encode a human allelic variant are highly preferred.

A highly preferred polynucleotide of the invention comprises the sequence set forth in SEQ ID NO: 1, which comprises a human CON193 encoding DNA sequence:

```

20 ntgggtgttg gaccattaaa atgcattatg gaatttttaa aagttggggg agagggagac 60
   agtaaaaaa acctatattt tctctgtttt tttttttttt aactctagga aagcccagac 120
   aaattttgag ctatttcata acctaccaga cttatcatgc taacctgaa taaaacagac 180
   ctaataccag cttcatttat tctgaatgga gtcccaggac tggaagacac acaactctgg 240
   atttccttcc cattctgctc tatgtatggt gtggctatgg tagggaattg tggactcctc 300
25 tacctcattc actatgagga tgccctgcac aaacccatgt actacttctt ggccatgctt 360
   tcccttactg accttggtat gtgctctagt acaatcccta aagccctctg catcttctgg 420
   tttcatctca aggacattgg atttgatgaa tgccctgtcc agatgttctt catccacacc 480
   ttcacaggga tggagtctgg ggtgcttatg cttatggccc tggatcgcta tgtggccatc 540
   tgctacctct tacgctatc aactatcctc accaatcctg taattgcaaa ggttggggact 600
30 gccaccttcc tgagaggggt attactcatt attcccttta ctttctcacc caagcgccctg 660
   cccctctgca gaggcaatat acttcccat accactgtg accacatgct tgtagccaaa 720
   ttgtcctgtg gtaatgtcaa ggtcaatgcc atctatggtc tgatggtlgc cctcctgatt 780
   gggggccttg acatactgtg taccaccatc tctatacca tgattctcgg ggcagtgggtc 840
   agcctctcct cagcagatgc tcggcagaag gcctttaata cctgcactgc ccacatttgt 900
35 gccattgttt tctctatac tccagcttcc ttctccttct tttccaccg ctttggggaa 960
   cacataatcc ccccttcttg ccacatcatt gttagcaata tttatctgct cctaccaccc 1020
   actatgaacc ctattgtcta tggggtgaaa accaaacaga tacgagactg tgcataaagg 1080

```

- 22 -

atcctttcag gttctaagga taccaaatcc tacagcatgt gaatgaacac ttgccaggag 1140  
 tgagaagaga aggaaagaat tactttctatt tgccctcttat gcaggagtgc ataaaaatcctt 1200  
 tctggaagta ctgtattgat cacaaaatgg agtttgntga ctggtgcatt ctcaataagt 1260  
 accttgggaa tctnacatca ctggaaggcc caccacattt ctataaat 1308

5 Also preferred is a polynucleotide comprising nucleotides 157-1119 of  
 SEQ ID NO: 1, which represent the portion of SEQ ID NO: 1 that encodes CON193  
 amino acids.

Another highly preferred polynucleotide of the invention comprises the  
 sequence set forth in SEQ ID NO: 3, which comprises a human CON166 encoding  
 10 DNA sequence:

atggatgaaa caggaaatct gacagtatct tctgccacat gccatgacac tattgatgac 60  
 ttccgcaatc aagtgtattc caccctgtac tctatgatct ctggtgtagg cttcttttggc 120  
 aatggctttg tgctctatgt cctcataaaa acctatcaca agaagtcagc cttccaagta 180  
 tacatgatta atttagcagt agcagatcta ctttggtgtg gcacactgcc tctccgtgtg 240  
 15 gtctattatg ttcacaaagg catttggtct tttggtgact tcttggtgcc cctcagcacc 300  
 tatgctttgt atgtcaacct ctattgtagc atcttcttta tgacagccat gagctttttc 360  
 cgggtgcattg caattgtttt tccagtcag aacattaatt tggttacaca gaaaaaagcc 420  
 aggtttgtgt gtgtaggtat ttggattttt gtgattttga ccagttctcc atttctaattg 480  
 gccaaaccac aaaaagatga gaaaaataat accaagtgtc ttgagccccc acaagacaat 540  
 20 caaactaaaa atcatgtttt ggtcttgcac tatgtgtcat tgtttgttgg ctttatcacc 600  
 ccttttgtta ttataattgt ctgttacaca atgacattt tgacctact aaaaaaatca 660  
 atgaaaaaaa atctgtcaag tcataaaaag gctataggaa tgatcatggt cgtgaccgct 720  
 gcctttttag tcagtttcat gccatatcat attcaacgta ccattcacct tcatttttta 780  
 cacaatgaaa ctaaaccttg tgattctgtc cttagaatgc agaagtcctg ggtcataacc 840  
 25 ttgtctctgg ctgcatccaa ttgttgcttt gacctctcc tatatttctt ttctgggggt 900  
 aacttttaga aaaggctgtc tacatttaga aagcattctt tgtccagcgt gacttatgta 960  
 cccagaaaga aggcctcttt gccagaaaaa ggagaagaaa tatgtaaagt atag 1014

The final three nucleotides of this sequence represent a stop codon.

Still another highly preferred polynucleotide of the invention  
 30 comprises the sequence set forth in SEQ ID NO: 5, which comprises a human  
 CON103 encoding DNA sequence:

ggggcctact tcaccgtgta cccggacttg ggaccatcac agacttcaga accatcagga 60  
 acctgggagc aactgaaagc tgaactacag tgggctttca gacacacagc aggctgcgga 120  
 gcacaaatag gactgggtcc ctccaggcca ccagcagggc ggtggaggtc ttcaactgact 180  
 35 cctgcctac ctctcaggac aatgtccttt tggtccaca gtcctgaag ccagagctgg 240  
 tgggggcagg gaggcagcca ccagcctcta tatgtagtgg aggagggggg gtccagggag 300  
 ggctgcatga tctgagagc cccacctca cccggctgga ctatcctccc acttcagggt 360  
 ttctctgggc ttccatcttg cccctgtga gccctgcttc ctcctctacc agcagcacia 420  
 cccccaggct gggctcagag acctcatgtg gtgggacac tcagtacccc gaggcggagg 480  
 40 gaaggaggga gggctgcagg gttcccttg gcctgcaaac aggaacacag ggtgtttctc 540  
 agtggctgcg agaatgtga tgaaaacccc aggatgttgt gtcaccgtgg tggccagctg 600  
 atagtcccaa tcatccact ttgccttgag cactcctgca ggggtagaag actccagaac 660

- 23 -

	cttctctcag gcccatggcc caagcagccc atg gaa ctt cat aac ctg agc tct	714
	cca tet ccc tct ctc tcc tcc tct gtt ctc cct ccc tcc ttc tct ccc	762
	tca ccc tcc tct gct ccc tct gcc ttt acc act gtg ggg ggg tcc tct	810
	gga ggg ccc tgc cac ccc acc tct tcc tgc ctg gtg tct gcc ttc ctg	858
5	gca cca atc ctg gcc ctg gag ttt gtc ctg ggc ctg gtg ggg aac agt	906
	ttg gcc ctc ttc atc ttc tgc atc cac acg cgg ccc tgg acc tcc aac	954
	acg gtg ttc ctg gtc agc ctg gtg gcc gct gac ttc ctc ctg atc agc	1002
	aac ctg ccc ctc cgc gtg gac tac tac ctc ctc cat gag acc tgg cgc	1050
	ttt ggg gct gct gcc tgc aaa gtc aac ctc ttc atg ctg tcc acc aac	1098
10	cgc acg gcc agc gtt gtc ttc ctc aca gcc atc gca ctc aac cgc tac	1146
	ctg aag gtg gtg cag ccc cac cac gtg ctg agc cgt gct tcc gtg ggg	1194
	gca gct gcc cgg gtg gcc ggg gga ctc tgg gtg ggc atc ctg ctc ctc	1242
	aac ggg cac ctg ctc ctg agc acc ttc tcc ggc ccc tcc tgc ctc agc	1290
	tac agg gtg ggc acg aag ccc tgc gcc tgc ctc cgc tgg cac cag gca	1338
15	ctg tac ctg ctg gag ttc ttc ctg cca ctg ggc ctc atc ctc ttt gct	1386
	att gtg agc att ggg ctc acc atc cgg aac cgt ggt ctg ggc ggg cag	1434
	gca ggc ccg cag agg gcc atg cgt gtg ctg gcc atg gtg gtg gcc gtc	1482
	tac acc atc tgc ttc ttg ccc agc atc atc ttt ggc atg gct tcc atg	1530
	gtg gct ttc tgg ctg tcc gcc tgc cga tcc ctg gac ctc tgc aca cag	1578
20	ctc ttc cat ggc tcc ctg gcc ttc acc tac ctc aac agt gtc ctg gac	1626
	ccc gtg ctc tac tgc ttc tct agc ccc aac ttc ctc cac cag agc cgg	1674
	gcc ttg ctg ggc ctc acg cgg ggc cgg cag ggc cca gtg agc gac gag	1722
	agc tcc tac caa ccc tcc agg cag tgg cgc tac cgg gag gcc tct agg	1770
	aag gcg gag gcc ata ggg aag ctg aaa gtg cag ggc gag gtc tct ctg	1818
25	gaa aag gaa ggc tcc tcc cag ggc tga gggcagctg cagggctgca	1865
	gcgctgtggg ggtaagggct gccgcgctct ggcctggagg gacaaggcca gcacacggtg	1925
	cctcaaccaa ctggacaagg gatggcgcca gaccaggggc caggccaaag cactggcagg	1985
	actcatgttg gtggcagggg gagaaaccca cctaggcctc tcagtgtgtc caggatggca	2045
	ttcccagaat gcaggggaga gcaggatgcc ggggtggagga gacaggcaag gtgccgttg	2105
30	cacaccagct cagacagggg cctgcgcagc tgcaggggac agacgccaat cactgtcaca	2165
	gcagagtcac cttagaaatt ggacagctgc atgttctgtg ctctccagtt tgtcccttcc	2225
	aatattaata aacttccctt ttaaataat tttattgcag accaatatct gtctttaatt	2285
	ctaacctggg actgtcagta ggcgtcaaag tgagcgcccc agtgaaggaa ccttgagag	2345
	agtgggagca ttcccagcct tccaggggga ctctcttcc agactttgga gcccgcatgt	2405
35	ctgaagcaga ctctttcttg gtgag	2429

Also preferred is a polynucleotide comprising nucleotides 691-1842 of SEQ ID NO:

5, which represent the portion of SEQ ID NO: 5 that encodes CON103 amino acids.

Nucleotides 1843-1845 represent a stop codon.

Another highly preferred polynucleotide of the invention comprises the

40 sequence set forth in SEQ ID NO: 7, which comprises a CON203-encoding DNA sequence:

```

ttgaatttag gtgacactat agaagagcta tgacgtcgca tgcacgcgta cgtaagctcg 60
gaattcggct cgagctgaac taatgactgc cgccataaga agacagagag aactgagtat 120
cctcccaaaag gtgacactgg aagcaatgaa caccacagtg atgcaaggct tcaacagatc 180

```

- 24 -

5      tgagcgggtgc cccagagaca ctcggatagt acagctggta tccccagccc tctacacagt 240  
       ggttttcttg accggcatcc tgctgaatac tttggctctg tgggtgtttg ttcacatccc 300  
       cagctcctcc accttcatca tctacctcaa aaacactttg gtggccgact tgataatgac 360  
       actcatgctt cctttcaaaa tcttctctga ctcacacctg gcacctggc agctcagagc 420  
 10     ttttgtgtgt cgtttttctt cggatgattt ttatgagacc atgtatgttg gcatcgtgct 480  
       gttagggttc atagcctttg acagattcct caagatcctc agacctttga gaaatatttt 540  
       tctaaaaaaa cctgtttttg caaaaacggt ctcaatcttc atctgggtct ttttgggtct 600  
       catctccctg ccaaataatga tcttgagcaa caagggaagca acaccatcgt ctgtgaaaaa 660  
       gtgtgcttcc ttaaaggggc ctctggggct gaaatggcat caaatggtaa ataacatatg 720  
 15     ccagtttatt ttctggactg gttttatcct aatgcttgtg ttttatgttg ttattgcaaa 780  
       aaaagtatat gattcttata gaaagtcaa aagtaaggac agaaaaaaca acaaaaagct 840  
       ggaaggcaaa gtatttgttg tcgtggctgt cttctttgtg tgttttctc catttcattt 900  
       tgccagagtt ccatatactc acagtcaaac caacaataag actgactgta gactgcaaaa 960  
       tcaactgttt attgctaaag aaacaactct ctttttggca gcaactaaca tttgtatgga 1020  
 15     tcccttaata tacatattct tatgtaaaaa attcacagaa aagctacat gtatgcaagg 1080  
       gagaaagacc acagcatcaa gccaaagaaa tcatagcagt cagacagaca acataacctt 1140  
       aggtgacaaa ctgtacatag ggttaacttc tatttattga tgagacttcc gtagataatg 1200  
       tggaaatcaa atttaaccaa gaaaaaaga ttggaacaaa tgctctctta cattttattt 1260  
       atcctgggtg ccaggaaaag attatattaa atttaaatcc acatagatct attcataagc 1320  
 20     tgaatgaacc attacctaag agaatgaac aggataccaa tggccactag aggcatattc 1380  
       cttctctctt ttttttctt aaatttcaag agcattcact ttacatttgg aaagactaag 1440  
       gggaaagggt atcctacaaa cctcccttca acacctttta catt 1484

Also preferred is a polynucleotide comprising nucleotides 146-1144 of SEQ ID NO:

7, which represent the portion of SEQ ID NO: 7 that encodes CON203 amino acids.

25      Nucleotides 1145-1147 represent a stop codon.

Another highly preferred polynucleotide of the invention comprises the sequence set forth in SEQ ID NO: 9, which comprises a human CON198 encoding DNA sequence:

30      ATGATGGTGG ATCCCAATGG CAATGAATCC AGTGCTACAT ACTTCATCCT AATAGGCCTC 60  
       CCTGGTTTAG AAGAGGCTCA GTTCTGGTTG GCCTTCCCAT TGTGCTCCCT CTACCTTATT 120  
       GCTGTGCTAG GTAACCTGAC AATCATCTAC ATTGTGCGGA CTGAGCACAG CCTGCATGAG 180  
       CCCATGTATA TATTTCTTTG CATGCTTTCA GGCATTGACA TCCTCATCTC CACCTCATCC 240  
       ATGCCCAAAA TGCTGGCCAT CTTCTGGTTC AATTCCACTA CCATCCAGTT TGATGCTTGT 300  
       CTGTACAGA TGTTTGCCAT CCACTCCTTA TCTGGCATGG AATCCACAGT GCTGCTGGCC 360  
 35     ATGGCTTTTG ACCGCTATGT GGCCATCTGT CACCCACTGC GCCATGCCAC AGTACTTACG 420  
       TTGCCTCGTG TCACCAAAAT TGGTGTGGCT GCTGTGGTGC GGGGGGCTGC ACTGATGGCA 480  
       CCCCTTCCTG TCTTCATCAA GCAGCTGCC TTTCTGCCGT CCAATATCCT TTCCCATTC 540  
       TACTGCCTAC ACCAAGATGT CATGAAGCTG GCCTGTGATG ATATCCGGGT CAATGTCGTC 600  
       TATGGCCTTA TCGTCATCAT CTCCGCCATT GGCCTGGACT CACTTCTCAT CTCCTTCTCA 660  
 40     TATCTGCTTA TTCTTAAGAC TGTGTTGGC TTGACACGTG AAGCCCAGGC CAAGGCATTT 720  
       GGCATTGCG TCTCTCATGT GTGTGCTGTG TTCATATTCT ATGTACCTTT CATTGGATTG 780  
       TCCATGGTGC ATCGCTTTAG CAAGCGCGT GACTCTCCGC TGCCCGTCAT CTTGGCCAAT 840  
       ATCTATCTGC TGGTTCCTCC TGTGCTCAAC CCAATTGTCT ATGGAGTGAA GACAAAGGAG 900  
       ATTCGACAGC GCATCCTTCG ACTTTTCCAT GTGGCCACAC ACGCTTCAGA GCCCTAG 957

- 25 -

The last three nucleotides of this sequence represent a stop codon.

Still another A highly preferred polynucleotide of the invention comprises the sequence set forth in SEQ ID NO: 11, which comprises a human CON197 encoding DNA sequence:

```

5  ATGGAAGCG AGAACAGAAG AGTGATAAGA GAATTCATCC TCCTTGGTCT GACCCAGTCT 60
   CAAGATATTC AGCTCCTGGT CTTTGTGCTA GTTTTAATAT TCTACTTCAT CATCTCCCT 120
   GGAAATTTTC TCATTATTTT CACCATAAAG TCAGACCCTG GGCTCACAGC CCCCCTCTAT 180
   TTCTTTCTGG GCAACTTGGC CTTCTGGAT GCATCCTACT CCTTCATTGT GGCTCCCCGG 240
   ATGTTGGTGG ACTTCCTCTC TGCGAAGAAG ATAATCTCCT ACAGAGGCTG CATCACTCAG 300
10 CTCTTTTCT TGCACCTCCT TGGAGGAGGG GAGGGATTAC TCCTTGTGTG GATGGCCTTT 360
   GACCGCTACA TCGCCATCTG CCGGCCTCTG CACTATCCTA CTGTCATGAA CCCTAGAACC 420
   TGCTATGCAA TGATGTTGGC TCTGTGGCTT GGGGGTTTTG TCCACTCCAT TATCCAGGTG 480
   GTCCTCATCC TCCGCTTGCC TTTTGTGGC CCAAACCAGC TGGACAACTT CTTCTGTGAT 540
   GTCCACAGG TCATCAAGCT GGCCTGCACC GACACATTG TGGTGGAGCT TCTGATGGTC 600
15 TTCAACAGTG GCCTGATGAC ACTCCTGTGC TTTCTGGGGC TTCTGGCCTC CTATGCAGTC 660
   ATTCTTTGTC GCATACGAGG GTCTTCTTCT GAGGCAAAAA ACAAGGCCAT GTCCACGTGC 720
   ATCACCATA TCATTGTTAT ATTCTTCATG TTTGGACCTG GCATCTTCAT CTACACGCGC 780
   CCCTTCAGGG CTTTCCCAGC TGACAAGGTG GTTCTCTCT TCCACACAGT GATTTTCTCT 840
   TTGTTGAATC CTGTCAATTA TACCCTTCGC AACCAGGAAG TGAAAGCTTC CATGAAAAAG 900
20 GTGTTTAATA AGCACATAGC CTGA 924

```

The last three nucleotides of this sequence represent a stop codon.

Another highly preferred polynucleotide of the invention comprises the sequence set forth in SEQ ID NO: 13, which comprises a human CON202 encoding DNA sequence:

```

25 1      TGCTTCCCCA TAAGGTAACA GCTTTGTTAG CNCTGTCTGA CATCATTGCT
   51      TGTTWACTTA AGAAGTGATA GGTYYTTTTT TTTTTTTTTT TTCAGATATT
   101     CTGATGGCAA AACAAGTGGA AGAAAAGAGG AAGCATGACT GCAGATCAGA
   151     TCAGTTCTCT TTGTGGATTA TATTTTCAGT AAAATGTATG GATCTATCTT
   201     TTCCTTGTTT TTATATCTAG ATCATGAGAC TTGACTGAGG CTGTATCCTT
30 251     ATCCTCCATC CATCTATGGC GAACTATAGC CATGCAGCTG ACAACATTTT
   301     GCAAAATCTC TCGCCTCTAA CAGCCTTTCT GAAACTGACT TCCTTGGGTT
   351     TCATAATAGG AGTCAGCGTG GTGGGCAACC TCCTGATCTC CATTTTGCTA
   401     GTGAAAGATA AGACCTTGCA TAGAGCACCT TACTACTTCC TGTTGGATCT
   451     TTGCTGTTCA GATATCTCA GATCTGCAAT TTGTTTCCA TTTGTGTTCA
35 501     ACTCTGTCAA AAATGGTTCT ACCTGGACTT ATGGGACTCT GACTTGCAAA
   551     GTGATTGCCT TTCTGGGGGT TTTGTCCTGT TTCCACACTG CTTTCATGCT
   601     CTTCTGCATC AGTGTACCA GATATTTAGC TATCGCCCAT CACCGCTTCT
   651     ATACAAAGAG GCTGACCTTT TGGACGTGTC TGGCTGTGAT CTGTATGGTG
   701     TGGACTCTGT CTGTGGCCAT GGCATTTCCT CCGGTTTTAG ACGTGGGCAC

```

- 26 -

751 TTACTCATTG ATTAGGGAGG AAGATCAATG CACCTTCCAA CACCGCTCCT  
 801 TCAGGGCTAA TGATTCCTTA GGATTTATGC TGC'TTCTTG TCTCATCCTC  
 851 CTAGCCACAC AGCTTGTCTA CCTCAAGCTG ATATTTTTCG TCCACGATCG  
 901 AAGAAAAATG AAGCCAGTCC AGTTTGTAGC AGCAGTCAGC CAGAACTGGA  
 5 951 CTTTTCATGG TCCTGGAGCC AGTGGCCAGG CAGCTGCCAA TTGGCTAGCA  
 1001 GGATTTGGAA GGGGTCCCAC ACCACCCACC TTGCTGGGCA TCAGGCAAAA  
 1051 TGCAAAACACC ACAGGCAGAA GAAGGCTATT GGTCTTAGAC GAGTTCAAAA  
 1101 TGGAGAAAAG AATCAGCAGA ATGTTCTATA TAATGACTTT TCTGTTTCTA  
 1151 ACCTTGTGGG GCCCCTACCT GGTGGCCTGT TATTGGAGAG TTTTGTCAAG  
 10 1201 AGGGCCTGTA GTACCAGGGG GATTTCTAAC AGCTGCTGTC TGGATGAGTT  
 1251 TTGCCCCAGC AGGAATCAAT CCTTTTGTCT GCATTTTCTC AAACAGGGAG  
 1301 CTGAGGCGCT GTTTCAGCAC AACCTTCTT TACTGCAGAA AATCCAGGTT  
 1351 ACCAAGGGAA CCTTACTGTG TTATATGAGG

Also preferred is a polynucleotide comprising nucleotides 266-1375 of SEQ ID NO:

15 13, which represent the portion of SEQ ID NO: 13 that encodes CON202 amino acids.  
 Nucleotides 1376-1378 represent a stop codon.

Another highly preferred polynucleotide of the invention comprises the  
 sequence set forth in SEQ ID NO: 15, which comprises a human CON222 encoding  
 DNA sequence:

20 1 ATGTTTAGAC CTCTGTGAA TCTCTCTCAC ATATATTTTA AGAAATTCCA  
 51 GTACTGTGGG TATGCACCAC ATGTTGCGAG CTGTAAACCA AACACTGATG  
 101 GAATTCATC TCTAGAGAAT CTCTGGCAA GCATTATCA GAGAGTATTT  
 151 GTCTGGGTTG TATCTGCAGT TACCTGCTTT GGAAACATTT TTGTCAATTTG  
 201 GATGCGACCT TATATCAGGT CTGAGAACAA GCTGTATGCC ATGTCAATCA  
 25 251 TTTCTCTCTG CTGTGCCGAC TGCTTAATGG GAATATATTT ATTCGTGATC  
 301 GGAGGCTTTG ACCTAAAGTT TCGTGAGAA TACAATAAGC ATGCGCAGCT  
 351 GTGGATGGAG AGTACTCATT GTCAGCTTGT AGGATCTTTG GCCATTCTGT  
 401 CCACAGAAGT ATCAGTTTTA CTGTTAACAT TTCTGACATT GGAAAAATAC  
 451 ATCTGCATTG TCTATCCTTT TAGATGTGTG AGACCTGGAA AATGCAGAAC  
 30 501 AATTACAGTT CTGATTCTCA TTTGGATTAC TGGTTTTATA GTGGCTTTCA  
 551 TTCCATTGAG CAATAAGGAA TTTTCAAAA ACTACTATGG CACCAATGGA  
 601 GTATGCTTCC CTCTCATTG AGAAGATACA GAAAGTATTG GAGCCCAGAT  
 651 TTATTCASTG GCAATTTTTT TGGTATTAA TTTGGCCGCA TTTATCATCA  
 701 TAGTTTTTTC CTATGGAAGC ATGTTTTATA GTGTTTATCA AAGTGCCATA  
 35 751 ACAGCAACTG AAATACGGAA TCAAGTTAAA AAAGAGATGA TCCTTGCCAA  
 801 ACGTTTTTTC TTTATAGTAT TTAAGTATGC ATTATGCTGG ATACCCATTT  
 851 TTGTAGTGAA ATTTCTTTCA CTGCTTCAGG TAGAAATACC AGGTACCATA  
 901 ACCTCTTGGG TAGTGATTTT TATTCTGCCC ATTAACAGTG CTTTGAACCC  
 951 AATTCTCTAT ACTCTGACCA CAAGACCATT TAAAGAAATG ATTCATCGGT  
 40 1001 TTTGGTATAA CTACAGACAA AGAAAATCTA TGGACAGCAA AGGTCAGAAA  
 1051 ACATATGCTC CATCATTCAT CTGGGTGGAA ATGTGGCCAC TGCAGGAGAT  
 1101 GCCACCTGAG TTAATGAAGC CGGACCTTTT CACATACCCC TGTGAAATGT

- 27 -

1151 CACTGATTTC TCAATCAACG AGACTCAATT CCTATTCA

The last three nucleotides of this sequence represent a stop codon.

Another highly preferred polynucleotide of the invention comprises the sequence set forth in SEQ ID NO: 17, which comprises a human CON215 encoding DNA sequence. Also preferred is a polynucleotide comprising the portion of SEQ ID NO: 17 set forth below, which represent the portion of SEQ ID NO: 17 that encodes CON215 amino acids (the last three nucleotides represent a stop codon).

ATGGGGTTCA ACTTGACGCT TGCAAAATTA CCAAATAACG AGCTGCACGG CCAAGAGAGT 60  
 CACAATTCAG GCAACAGGAG CGACGGGCCA GGAAAGAACA CCACCCTTCA CAATGAATTT 120  
 10 GACACAATTG TCTTGCCAGT GCTTATATCTC ATTATATTTG TGGCAAGCAT CTTGCTGAAT 180  
 GGTTTAGCAG TGTGGATCTT CTTCACATT AGGAATAAAA CCAGCTTCAT ATTCTATCTC 240  
 AAAACATAG TGGTTGCAGA CCTCATAATG ACGCTGACAT TTCCATTTG AATAGTCCAT 300  
 GATGCAGGAT TTGGACCTTG GTACTTCAAG TTTATTCTCT GCAGATACAC TTCAGTTTTG 360  
 TTTTATGCAA ACATGTATAC TTCCATCGTG TTCCTTGGGC TGATAAGCAT TGATCGCTAT 420  
 15 CTGAAGGTGG TCAAGCCATT TGGGGACTCT CGGATGTACA GCATAACCTT CACGAAGGTT 480  
 TTATCTGTTT GTGTTTGGGT GATCATGGCT GTTTTGTCTT TGCCAAACAT CATCCTGACA 540  
 AATGGTCAGC CAACAGAGGA CAATATCCAT GACTGCTCAA AACTTAAAAG TCCTTTGGGG 600  
 GTCAAATGGC ATACGGCAGT CACCTATGTG AACAGCTGCT TGTTTGTGGC CGTGTGTTG 660  
 ATTCTGATCG GATGTTACAT AGCCATATCC AGGTACATCC ACAAAATCCAG CAGGCAATTC 720  
 20 ATAAGTCAGT CAAGCCGAAA GCGAAAACAT AACCAGAGCA TCAGGGTTGT TGTGGCTGTG 780  
 TTTTITACCT GCTTCTTACC ATATCACTTG TGCAGAAITC CTTTACTTT TAGTCACTTA 840  
 GACAGGCTTT TAGATGAATC TGCACAAAAA ATCCTATATT ACTGCAAAGA AATTACACTT 900  
 TTCTTGCTG CGTGTAATGT TTGCCTGGAT CCAATAATTT ACTTTTTCAT GTGTAGGTCA 960  
 TTTTCAAGAA GGCTGTTCAA AAAATCAAAT ATCAGAACCA GGAGTGAAAG CATCAGATCA 1020  
 25 CTGCAAAGTG TGAGAAGATC GGAAGTTCTC ATATATTATG ATTATACTGA TGTGTAG 1077

Another preferred polynucleotide of the invention comprises the portion of the sequence set forth in SEQ ID NO: 19 which comprises a human CON217 encoding DNA sequence:

1 ATGTTAGCCA ACAGCTCCTC AACCAACAGT TCTGTCTCC CGTGTCTGA CTACCGACCT  
 30 61 ACCCACCGCC TGCACTTGGT GGTCTACAGC TTGGTGCTGG CTGCCGGGCT CCCCCTCAAC  
 121 GCGCTAGCCC TCTGGGTCTT CCTGCGCGCG CTGCGCGTGC ACTCGGTGGT GAGCGGTAC  
 181 ATGTGTAACC TGGCGGCCAG CGACCTGCTC TTCACCTCT CGCTGCCCGT TCGTCTCTCC  
 241 TACTACGCAC TGCACTCTG GCCCTTCCCC GACCTCCTGT GCCAGACGAC GGGCGCCATC  
 301 TTCCAGATGA ACATGTACGG CAGCTGCATC TTCCTGATGC TCATCAACGT GGACCGCTAC  
 35 361 GCGGCCATCG TGCACCGGCT GCGACTGCGC CACCTGCGGC GGCCCCGCGT GCGCGGCTG  
 421 CTCTGCTGG GCGTGTGGGC GCTCATCCTG GTGTTTGCCG TGCCCCGCGC CCGCGTGCAC  
 481 AGGCCCTCGC GTTGCCGCTA CCGGACCTC GAGGTGCGCC TATGCTTCGA GAGCTTCAGC  
 541 GACGAGCTGT GGAAAGGCAG GCTGCTGCCC CTCGTGCTGC TGGCCGAGGC GCTGGGCTTC  
 601 CTGCTGCCCC TGGCGGCGGT GGTCTACTCG TCGGGCCGAG TCTTCTGGAC GCTGGCGCGC  
 40 661 CCGACGCCA CGCAGAGCCA GCGGCGCGG AAGACCGTGC GCCTCCTGCT GGCTAACCTC  
 721 GTCATCTTCC TGCTGTGCTT CGTGCCCTAC AACAGCACGC TGGCGGTCTA CGGGCTGCTG  
 781 CGGAGCAAGC TGGTGGCGGC CAGCGTGCCT GCCCGGATC GCGTGCAGCG GGTGCTGATG

- 28 -

841 GTGATGGTGC TGCTGGCCGG CGCCAACTGC GTGCTGGACC CGCTGGTGTA C'ACTTTAGC  
 901 CCCGAGGGCT TCCGCAACAC CCTGCGCGGC CTGGGCACTC CGCACCAGGC CAGGACCTCG  
 961 GCCACCAACG GGACGCGGGC GCGCTCGCG CAATCCGAAA GTCCGCGGT CACCACCGAC  
 1021 GCCACCAGGC CGGATGCCGC CAGTCAGGGG CTGCTCCGAC CCTCCGACTC CCACTCTCTG  
 5 1081 TCTTCCTTCA CACAGTGTCC CCAGGATTCC GCCCTCTGA

The last three nucleotides of this sequence represent a stop codon.

The invention also includes polynucleotides differing from the sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17 and 19 and from their complementary strand by at least one nucleotide.

10 In a related embodiment, the invention provides vectors comprising a polynucleotide of the invention. Such vectors are useful, *e.g.*, for amplifying the polynucleotides in host cells to create useful quantities thereof. In preferred embodiments, the vector is an expression vector wherein the polynucleotide of the invention is operatively linked to a polynucleotide comprising an expression control  
 15 sequence. Such vectors are useful for recombinant production of polypeptides of the invention.

In another related embodiment, the invention provides host cells that are transformed or transfected (stably or transiently) with a polynucleotide of the invention or vectors of the invention. As stated above, such host cells are useful for  
 20 amplifying the polynucleotides and also for expressing the GPCR seven transmembrane receptor polypeptides or fragments thereof encoded by the polynucleotides. Such host cells are useful in assays as described herein.

In still another related embodiment, the invention provides a method for producing a seven transmembrane receptor polypeptide (or fragment thereof) of  
 25 the invention comprising the steps of growing a host cell of the invention in a nutrient medium and isolating the polypeptide or variant thereof from the cell or the medium. Since the GPCR polypeptides are seven transmembrane receptors, it will be appreciated that, for some applications, such as certain activity assays, the preferable isolation may involve isolation of cell membranes containing the polypeptide  
 30 embedded therein, whereas for other applications a more complete isolation may be preferable.

In still another embodiment, the invention provides antibodies that are specific for the GPCR seven transmembrane receptors of the invention. Antibody

- 29 -

specificity is described in greater detail below. However, it should be emphasized that antibodies that can be generated from polypeptides that have previously been described in the literature and that are capable of fortuitously cross-reacting with the GPCR polypeptides of the invention (e.g., due to the fortuitous existence of a similar epitope in both polypeptides) are considered "cross-reactive" antibodies. Such cross-reactive antibodies are not antibodies that are "specific" for the GPCR polypeptides. The determination of whether an antibody is specific for a GPCR polypeptide or is cross-reactive with another known receptor is made using Western blotting assays or several other assays well known in the literature. For identifying cells that express GPCR polypeptides and also for modulating GPCR -ligand binding activity, antibodies that specifically bind to an extracellular epitope of one of the GPCR seven transmembrane receptors of the present invention are preferred.

In one preferred variation, the invention provides monoclonal antibodies. Hybridomas that produce such antibodies also are intended as aspects of the invention. In yet another variation, the invention provides a humanized antibody. Humanized antibodies are useful for *in vivo* therapeutic indications.

In another variation, the invention provides a cell-free composition comprising polyclonal antibodies, wherein at least one of the antibodies is an antibody of the invention specific for a GPCR polypeptide of the present invention. Antisera isolated from an animal is an exemplary composition, as is a composition comprising an antibody fraction of an antisera that has been resuspended in water or in another diluent, excipient, or carrier.

In still another related embodiment, the invention provides anti-idiotypic antibodies specific for an antibody that is specific for a GPCR polypeptide of the present invention.

It is well known that antibodies contain relatively small antigen binding domains that can be isolated chemically or by recombinant techniques. Such domains are useful GPCR binding molecules themselves, and also may be reintroduced into human antibodies, or fused to toxins or other polypeptides. Thus, in still another embodiment, the invention provides a polypeptide comprising a fragment of a GPCR-specific antibody, wherein the fragment and the polypeptide bind to a

- 30 -

GPCR seven transmembrane receptor of the present invention. By way of non-limiting example, the invention provides polypeptides that are single chain antibodies and CDR-grafted antibodies.

Also within the scope of the invention are compositions comprising  
5 polypeptides, polynucleotides, or antibodies of the invention that have been formulated with, *e.g.*, a pharmaceutically acceptable carrier.

The invention also provides methods of using antibodies of the invention. For example, the invention provides a method for modulating ligand binding of a GPCR seven transmembrane receptor of the present invention comprising  
10 the step of contacting the seven transmembrane receptor with an antibody specific for the seven transmembrane receptor, under conditions wherein the antibody binds the receptor.

GPCR polypeptides are expressed in the brain, providing an indication that aberrant GPCR polypeptide signaling activity may correlate with one or more  
15 neurological disorders. The invention also provides a method for treating a neurological disorder comprising the step of administering to a mammal in need of such treatment an amount of an antibody-like polypeptide of the invention that is sufficient to modulate ligand binding of a GPCR seven transmembrane receptor of the present invention in neurons of the mammal. In addition to administration of  
20 antibody-like polypeptides, administration of natural ligands for GPCR polypeptides as well as modulators of GPCR polypeptide activity, such as small molecules that mimic, agonize or antagonize ligand-mediated GPCR polypeptide signaling, are contemplated. The expression pattern provides an indication that such molecules will have utility for treating neurological and/or psychiatric diseases, including but not  
25 limited to schizophrenia, depression, anxiety, bipolar disease, affective disorders, attention deficit hyperactivity disorder/attention deficit disorder (ADHD/ADO), epilepsy, neuritis, neurasthenia, neuropathy, neuroses, Alzheimer's disease, Parkinson's disease, migraine, senile dementia, and the like. Treatment of individuals having any of these disorders is contemplated as an aspect of the invention.

30 Thus, in yet another embodiment, the invention provides genetic screening procedures that entail analyzing a person's genome -- in particular their

- 31 -

alleles for GPCR's of the invention -- to determine whether the individual possesses a genetic characteristic found in other individuals that are considered to be afflicted with, or at risk for, developing a mental disorder or disease of the brain that is suspected of having a hereditary component. For example, in one embodiment, the invention provides a method for determining a potential for developing a disorder affecting the brain in a human subject comprising the steps of analyzing the coding sequence of one or more GPCR genes from the human subject; and determining development potential for the disorder in said human subject from the analyzing step.

More particularly, the invention provides a method of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor, comprising the steps of: (a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering the amino acid sequence, expression, or biological activity of at least one seven transmembrane receptor that is expressed in the brain, wherein the seven transmembrane receptor comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, and 20, or an allelic variant thereof, and wherein the nucleic acid corresponds to the gene encoding the seven transmembrane receptor; and (b) diagnosing the disorder or predisposition from the presence or absence of said mutation, wherein the presence of a mutation altering the amino acid sequence,

baptists

52

human f

- 30 -

GPCR seven transmembrane receptor of the present invention. By way of non-limiting example, the invention provides polypeptides that are single chain antibodies and CDR-grafted antibodies.

Also within the scope of the invention are compositions comprising  
5 polypeptides, polynucleotides, or antibodies of the invention that have been formulated with, *e.g.*, a pharmaceutically acceptable carrier.

The invention also provides methods of using antibodies of the invention. For example, the invention provides a method for modulating ligand binding of a GPCR seven transmembrane receptor of the present invention comprising  
10 the step of contacting the seven transmembrane receptor with an antibody specific for the seven transmembrane receptor, under conditions wherein the antibody binds the receptor.

GPCR polypeptides are expressed in the brain, providing an indication that aberrant GPCR polypeptide signaling activity may correlate with one or more  
15 neurological disorders. The invention also provides a method for treating a neurological disorder comprising the step of administering to a mammal in need of such treatment an amount of an antibody-like polypeptide of the invention that is sufficient to modulate ligand binding of a GPCR seven transmembrane receptor of the present invention in neurons of the mammal. In addition to administration of  
20 antibody-like polypeptides, administration of natural ligands for GPCR polypeptides as well as modulators of GPCR polypeptide activity, such as small molecules that mimic, agonize or antagonize ligand-mediated GPCR polypeptide signaling, are contemplated. The expression pattern provides an indication that such molecules will have utility for treating neurological and/or psychiatric diseases, including but not  
25 limited to schizophrenia, depression, anxiety, bipolar disease, affective disorders, attention deficit hyperactivity disorder/attention deficit disorder (ADHD/ADO), epilepsy, neuritis, neurasthenia, neuropathy, neuroses, Alzheimer's disease, Parkinson's disease, migraine, senile dementia, and the like. Treatment of individuals having any of these disorders is contemplated as an aspect of the invention.

30 Thus, in yet another embodiment, the invention provides genetic screening procedures that entail analyzing a person's genome -- in particular their

alleles for GPCR's of the invention -- to determine whether the individual possesses a genetic characteristic found in other individuals that are considered to be afflicted with, or at risk for, developing a mental disorder or disease of the brain that is suspected of having a hereditary component. For example, in one embodiment, the invention provides a method for determining a potential for developing a disorder affecting the brain in a human subject comprising the steps of analyzing the coding sequence of one or more GPCR genes from the human subject; and determining development potential for the disorder in said human subject from the analyzing step.

More particularly, the invention provides a method of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor, comprising the steps of: (a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering the amino acid sequence, expression, or biological activity of at least one seven transmembrane receptor that is expressed in the brain, wherein the seven transmembrane receptor comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, and 20, or an allelic variant thereof, and wherein the nucleic acid corresponds to the gene encoding the seven transmembrane receptor; and (b) diagnosing the disorder or predisposition from the presence or absence of said mutation, wherein the presence of a mutation altering the amino acid sequence, expression, or biological activity of allele in the nucleic acid correlates with an increased risk of developing the disorder. In preferred variations, the seven transmembrane receptor is CON202 comprising an amino acid sequence set forth in SEQ ID NO: 14, or an allelic variant thereof, and the disease is schizophrenia.

By "human subject" is meant any human being, human embryo, or human fetus. It will be apparent that methods of the present invention will be of particular interest to individuals that have themselves been diagnosed with a disorder affecting the brain or have relatives that have been diagnosed with a disorder affecting the brain.

By "screening for an increased risk" is meant determination of whether a genetic variation exists in the human subject that correlates with a greater likelihood of developing a disorder affecting the brain than exists for the human population as a

whole, or for a relevant racial or ethnic human sub-population to which the individual belongs. Both positive and negative determinations (i.e., determinations that a genetic predisposition marker is present or is absent) are intended to fall within the scope of screening methods of the invention. In preferred embodiments, the presence of a mutation altering the sequence or expression of at least one *CON202* seven transmembrane receptor allele in the nucleic acid is correlated with an increased risk of developing schizophrenia, whereas the absence of such a mutation is reported as a negative determination.

The "assaying" step of the invention may involve any techniques available for analyzing nucleic acid to determine its characteristics, including but not limited to well-known techniques such as single-strand conformation polymorphism analysis (SSCP) [Orita *et al.*, *Proc Natl. Acad. Sci. USA*, 86: 2766-2770 (1989)]; heteroduplex analysis [White *et al.*, *Genomics*, 12: 301-306 (1992)]; denaturing gradient gel electrophoresis analysis [Fischer *et al.*, *Proc. Natl. Acad. Sci. USA*, 80: 1579-1583 (1983); and Riesner *et al.*, *Electrophoresis*, 10: 377-389 (1989)]; DNA sequencing; RNase cleavage [Myers *et al.*, *Science*, 230: 1242-1246 (1985)]; chemical cleavage of mismatch techniques [Rowley *et al.*, *Genomics*, 30: 574-582 (1995); and Roberts *et al.*, *Nucl. Acids Res.*, 25: 3377-3378 (1997)]; restriction fragment length polymorphism analysis; single nucleotide primer extension analysis [Shumaker *et al.*, *Hum. Mutat.*, 7: 346-354 (1996); and Pastinen *et al.*, *Genome Res.*, 7: 606-614 (1997)]; 5' nuclease assays [Pease *et al.*, *Proc. Natl. Acad. Sci. USA*, 91:5022-5026 (1994)]; DNA Microchip analysis [Ramsay, G., *Nature Biotechnology*, 16: 40-48 (1999); and Chee *et al.*, U.S. Patent No. 5,837,832]; and ligase chain reaction [Whiteley *et al.*, U.S. Patent No. 5,521,065]. [See generally, Schafer and Hawkins, *Nature Biotechnology*, 16: 33-39 (1998).] All of the foregoing documents are hereby incorporated by reference in their entirety.

Thus, in one preferred embodiment involving screening *CON202* sequences, for example, the assaying step comprises at least one procedure selected from the group consisting of: (a) determining a nucleotide sequence of at least one codon of at least one *CON202* allele of the human subject; (b) performing a hybridization assay to determine whether nucleic acid from the human subject has a

nucleotide sequence identical to or different from one or more reference sequences;  
(c) performing a polynucleotide migration assay to determine whether nucleic acid  
from the human subject has a nucleotide sequence identical to or different from one or  
more reference sequences; and (d) performing a restriction endonuclease digestion to  
5 determine whether nucleic acid from the human subject has a nucleotide sequence  
identical to or different from one or more reference sequences.

In a highly preferred embodiment, the assaying involves sequencing of  
nucleic acid to determine nucleotide sequence thereof, using any available sequencing  
technique. [See, e.g., Sanger *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 74: 5463-5467  
10 (1977) (dideoxy chain termination method); Mirzabekov, *TIBTECH*, 12: 27-32 (1994)  
(sequencing by hybridization); Drmanac *et al.*, *Nature Biotechnology*, 16: 54-58  
(1998); U.S. Patent No. 5,202,231; and *Science*, 260: 1649-1652 (1993) (sequencing  
by hybridization); Kieleczawa *et al.*, *Science*, 258: 1787-1791 (1992) (sequencing by  
primer walking); (Douglas *et al.*, *Biotechniques*, 14: 824-828 (1993) (Direct  
15 sequencing of PCR products); and Akane *et al.*, *Biotechniques* 16: 238-241 (1994);  
Maxam and Gilbert, *Meth. Enzymol.*, 65: 499-560 (1977) (chemical termination  
sequencing), all incorporated herein by reference.] The analysis may entail sequencing  
of the entire seven transmembrane receptor gene genomic DNA sequence, or portions  
thereof; or sequencing of the entire seven transmembrane receptor coding sequence or  
20 portions thereof. In some circumstances, the analysis may involve a determination of  
whether an individual possesses a particular allelic variant, in which case sequencing  
of only a small portion of nucleic acid -- enough to determine the sequence of a  
particular codon characterizing the allelic variant -- is sufficient. This approach is  
appropriate, for example, when assaying to determine whether one family member  
25 inherited the same allelic variant that has been previously characterized for another  
family member, or, more generally, whether a person's genome contains an allelic  
variant that has been previously characterized and correlated with a mental disorder  
having a heritable component.

In another highly preferred embodiment, the assaying step comprises  
30 performing a hybridization assay to determine whether nucleic acid from the human  
subject has a nucleotide sequence identical to or different from one or more reference

- 34 -

sequences. In a preferred embodiment, the hybridization involves a determination of whether nucleic acid derived from the human subject will hybridize with one or more oligonucleotides, wherein the oligonucleotides have nucleotide sequences that correspond identically to a portion of the GPCR gene sequence taught herein, such as the CON202 coding sequence set forth in SEQ ID NO: 14, or that correspond identically except for one mismatch. The hybridization conditions are selected to differentiate between perfect sequence complementarity and imperfect matches differing by one or more bases. Such hybridization experiments thereby can provide single nucleotide polymorphism sequence information about the nucleic acid from the human subject, by virtue of knowing the sequences of the oligonucleotides used in the experiments.

Several of the techniques outlined above involve an analysis wherein one performs a polynucleotide migration assay, *e.g.*, on a polyacrylamide electrophoresis gel (or in a capillary electrophoresis system), under denaturing or non-denaturing conditions. Nucleic acid derived from the human subject is subjected to gel electrophoresis, usually adjacent to (or co-loaded with) one or more reference nucleic acids, such as reference GPCR-encoding sequences having a coding sequence identical to all or a portion of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, or 19 (or identical except for one known polymorphism). The nucleic acid from the human subject and the reference sequence(s) are subjected to similar chemical or enzymatic treatments and then electrophoresed under conditions whereby the polynucleotides will show a differential migration pattern, unless they contain identical sequences. [See generally Ausubel *et al.* (eds.), *Current Protocols in Molecular Biology*, New York: John Wiley & Sons, Inc. (1987-1999); and Sambrook *et al.*, (eds.), *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press (1989), both incorporated herein by reference in their entirety.]

In the context of assaying, the term "nucleic acid of a human subject" is intended to include nucleic acid obtained directly from the human subject (*e.g.*, DNA or RNA obtained from a biological sample such as a blood, tissue, or other cell or fluid sample); and also nucleic acid derived from nucleic acid obtained directly from the human subject. By way of non-limiting examples, well known procedures

- 35 -

exist for creating cDNA that is complementary to RNA derived from a biological sample from a human subject, and for amplifying (*e.g.*, via polymerase chain reaction (PCR)) DNA or RNA derived from a biological sample obtained from a human subject. Any such derived polynucleotide which retains relevant nucleotide sequence information of the human subject's own DNA/RNA is intended to fall within the definition of "nucleic acid of a human subject" for the purposes of the present invention.

In the context of assaying, the term "mutation" includes addition, deletion, and/or substitution of one or more nucleotides in the GPCR gene sequence (*e.g.*, as compared to the seven transmembrane receptor-encoding sequences set forth in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, or 19) and other polymorphisms that occur in introns (where introns exist) and that are identifiable via sequencing, restriction fragment length polymorphism, or other techniques. The various activity examples provided herein permit determination of whether a mutation modulates activity of the relevant receptor in the presence or absence of various test substances.

In a related embodiment, the invention provides methods of screening a person's genotype with respect to GPCR's of the invention, and correlating such genotypes with diagnoses for disease or with predisposition for disease (for genetic counseling). For example, the invention provides a method of screening for a CON202 hereditary schizophrenia genotype in a human patient, comprising the steps of: (a) providing a biological sample comprising nucleic acid from the patient, the nucleic acid including sequences corresponding to said patient's CON202 alleles; (b) analyzing the nucleic acid for the presence of a mutation or mutations; (c) determining a CON202 genotype from the analyzing step; and (d) correlating the presence of a mutation in a CON202 allele with a hereditary schizophrenia genotype. In a preferred embodiment, the biological sample is a cell sample containing human cells that contain genomic DNA of the human subject. The analyzing can be performed analogously to the assaying described in preceding paragraphs. For example, the analyzing comprises sequencing a portion of the nucleic acid (*e.g.*, DNA or RNA), the portion comprising at least one codon of the CON202 alleles.

- 36 -

Although more time consuming and expensive than methods involving nucleic acid analysis, the invention also may be practiced by assaying protein of a human subject to determine the presence or absence of an amino acid sequence variation in GPCR protein from the human subject. Such protein analyses may be performed, e.g., by fragmenting GPCR protein via chemical or enzymatic methods and sequencing the resultant peptides; or by Western analyses using an antibody having specificity for a particular allelic variant of the GPCR.

The invention also provides materials that are useful for performing methods of the invention. For example, the present invention provides oligonucleotides useful as probes in the many analyzing techniques described above. In general, such oligonucleotide probes comprise 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides that have a sequence that is identical, or exactly complementary, to a portion of a human GPCR gene sequence taught herein (or allelic variant thereof), or that is identical or exactly complementary except for one nucleotide substitution. In a preferred embodiment, the oligonucleotides have a sequence that corresponds in the foregoing manner to a human GPCR coding sequence taught herein, and in particular, the coding sequences set forth in SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, or 19. In one variation, an oligonucleotide probe of the invention is purified and isolated. In another variation, the oligonucleotide probe is labeled, e.g., with a radioisotope, chromophore, or fluorophore. In yet another variation, the probe is covalently attached to a solid support. [See generally Ausubel *et al.* And Sambrook *et al.*, *supra*.]

In a related embodiment, the invention provides kits comprising reagents that are useful for practicing methods of the invention. For example, the invention provides a kit for screening a human subject to diagnose schizophrenia or a genetic predisposition therefor, comprising, in association: (a) an oligonucleotide useful as a probe for identifying polymorphisms in a human CON202 seven transmembrane receptor gene, the oligonucleotide comprising 6-50 nucleotides that have a sequence that is identical or exactly complementary to a portion of a human CON202 gene sequence or CON202 coding sequence, except for one sequence

- 37 -

difference selected from the group consisting of a nucleotide addition, a nucleotide deletion, or nucleotide substitution; and (b) a media packaged with the oligonucleotide containing information identifying polymorphisms identifiable with the probe that correlate with schizophrenia or a genetic predisposition therefor.

5 Exemplary information-containing media include printed paper package inserts or packaging labels; and magnetic and optical storage media that are readable by computers or machines used by practitioners who perform genetic screening and counseling services. The practitioner uses the information provided in the media to correlate the results of the analysis with the oligonucleotide with a diagnosis. In a  
10 preferred variation, the oligonucleotide is labeled.

In still another embodiment, the invention provides methods of identifying those allelic variants of GPCR's of the invention that correlate with mental disorders. For example, the invention provides a method of identifying a seven transmembrane allelic variant that correlates with a mental disorder, comprising steps  
15 of: (a) providing a biological sample comprising nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny; (b) analyzing the nucleic acid for the presence of a mutation or mutations in at least one seven transmembrane receptor that is expressed in the brain, wherein the at least one seven transmembrane receptor comprises an amino acid sequence selected  
20 from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, and 20, or an allelic variant thereof, and wherein the nucleic acid includes sequence corresponding to the gene or genes encoding the at least one seven transmembrane receptor; (c) determining a genotype for the patient for the at least one seven transmembrane receptor from said analyzing step; and (d) identifying an allelic variant that correlates  
25 with the mental disorder from the determining step. To expedite this process, it may be desirable to perform linkage studies in the patients (and possibly their families) to correlate chromosomal markers with disease states. The chromosomal localization data provided herein facilitates identifying an involved GPCR with a chromosomal marker.

30 The foregoing method can be performed to correlate GPCR's of the invention to a number of disorders having hereditary components that are causative or

- 38 -

that predispose persons to the disorder. For example, in one preferred variation, the disorder is schizophrenia, and the at least one seven transmembrane receptor comprises CON202 having an amino acid sequence set forth in SEQ ID NO: 14, or an allelic variant thereof.

5                   Also contemplated as part of the invention are polynucleotides that comprise the allelic variant sequences identified by such methods, and polypeptides encoded by the allelic variant sequences, and oligonucleotide and oligopeptide fragments thereof that embody the mutations that have been identified. Such materials are useful in *in vitro* cell-free and cell-based assays for identifying lead compounds and therapeutics for treatment of the disorders. For example, the variants are used in activity assays, binding assays, and assays to screen for activity modulators described herein. In one preferred embodiment, the invention provides a purified and isolated polynucleotide comprising a nucleotide sequence encoding a CON202 receptor allelic variant identified according to the methods described above; and an oligonucleotide 10 that comprises the sequences that differentiate the allelic variant from the CON202 sequences set forth in SEQ ID NOs: 13 and 14. The invention also provides a vector comprising the polynucleotide (preferably an expression vector); and a host cell transformed or transfected with the polynucleotide or vector. The invention also provides an isolated cell line that is expressing the allelic variant GPCR polypeptide; 20 purified cell membranes from such cells; purified polypeptide; and synthetic peptides that embody the allelic variation amino acid sequence. In one particular embodiment, the invention provides a purified polynucleotide comprising a nucleotide sequence encoding a CON202 seven transmembrane receptor protein of a human that is affected with schizophrenia; wherein said polynucleotide hybridizes to the complement of SEQ ID NO: 13 under the following hybridization conditions: (a) hybridization for 16 25 hours at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate and (b) washing 2 times for 30 minutes at 60°C in a wash solution comprising 0.1x SSC and 1% SDS; and wherein the polynucleotide encodes a CON202 amino acid sequence that differs from SEQ ID NO: 14 at at least one 30 residue.

- 39 -

An exemplary assay for using the allelic variants is a method for identifying a modulator of CON202 biological activity, comprising the steps of: (a) contacting a cell expressing the allelic variant in the presence and in the absence of a putative modulator compound; (b) measuring CON202 biological activity in the cell; and (c) identifying a putative modulator compound in view of decreased or increased CON202 biological activity in the presence versus absence of the putative modulator.

In still another example, the invention provides for a method of diagnosing schizophrenia or a susceptibility to schizophrenia comprising the steps of: determining the presence or amount of expression of CON202 polypeptide as set out as SEQ ID NO: 14 or the polypeptide encoded by the nucleic acid molecule having SEQ ID NO: 13 in a sample; and comparing the level of CON202 polypeptide in a biological, tissue or cellular sample from normal subjects or the subject at an earlier time, wherein the susceptibility to schizophrenia is based on the presence or amount of CON202 polypeptide expression.

The invention also provides for a method of treating schizophrenia comprising the step of administering to a human diagnosed with schizophrenia an amount of a modulator of CON202 receptor activity sufficient to modulate CON202 receptor activity or CON202 ligand binding in said human.

The invention also provides assays to identify compounds that bind GPCR seven transmembrane receptors. One such assay comprises the steps of: (a) contacting a composition comprising one of the GPCR seven transmembrane receptor polypeptides of the invention with a compound suspected of binding a GPCR polypeptide of the invention; and (b) measuring binding between the compound and the GPCR polypeptide. In one variation, the composition comprises a cell expressing a GPCR polypeptide of the invention on its surface. In another variation, an isolated GPCR polypeptide of the invention or cell membranes comprising a GPCR polypeptide of the invention are employed. The binding may be measured directly, e.g., using a labeled compound, or may be measured indirectly by several techniques, including measuring intracellular signaling of a GPCR polypeptide of the invention induced by the compound (or measuring changes in the level of GPCR polypeptide signaling).

The invention also provides a method for identifying a modulator of binding between a GPCR seven transmembrane receptor of the invention and a GPCR polypeptide binding partner, comprising the steps of: (a) contacting a GPCR polypeptide binding partner and a composition comprising one of the GPCR seven transmembrane receptors of the invention in the presence and in the absence of a putative modulator compound; (b) detecting binding between the binding partner and the GPCR polypeptide of the invention; and (c) identifying a putative modulator compound in view of decreased or increased binding between the binding partner and the GPCR polypeptide in the presence of the putative modulator, as compared to binding in the absence of the putative modulator.

GPCR polypeptide binding partners that stimulate GPCR seven transmembrane receptors of the present invention are useful as agonists in disease states characterized by insufficient GPCR polypeptide signaling (*e.g.*, as a result of insufficient expression of active GPCR polypeptide ligand). GPCR polypeptide binding partners that block ligand-mediated GPCR polypeptide signaling are useful as GPCR polypeptide antagonists to treat disease states characterized by excessive GPCR polypeptide signaling.

Additional features and variations of the invention will be apparent to those skilled in the art from the entirety of this application, including the detailed description, and all such features are intended as aspects of the invention. Likewise, features of the invention described herein can be re-combined into additional embodiments that also are intended as aspects of the invention, irrespective of whether the combination of features is specifically mentioned above as an aspect or embodiment of the invention. Also, only such limitations which are described herein as critical to the invention should be viewed as such; variations of the invention lacking limitations which have not been described herein as critical are intended as aspects of the invention.

In addition to the foregoing, the invention includes, as an additional aspect, all embodiments of the invention narrower in scope in any way than the variations specifically mentioned above. Although the applicant(s) invented the full scope of the claims appended hereto, the claims appended hereto are not intended to

- 41 -

encompass within their scope the prior art work of others. Therefore, in the event that statutory prior art within the scope of a claim is brought to the attention of the applicants by a Patent Office or other entity or individual, the applicant(s) reserve the right to exercise amendment rights under applicable patent laws to redefine the subject matter of such a claim to specifically exclude such statutory prior art or obvious variations of statutory prior art from the scope of such a claim. Variations of the invention defined by such amended claims also are intended as aspects of the invention.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention provides purified and isolated polynucleotides (*e.g.*, DNA sequences and RNA transcripts, both sense and complementary antisense strands, both single and double stranded, including splice variants thereof) encoding human G protein-coupled receptors referred to herein as GPCR polypeptides. DNA polynucleotides of the invention include genomic DNA, cDNA, and DNA that has been chemically synthesized in whole or in part. "Synthesized" as used herein and understood in the art, refers to polynucleotides produced by purely chemical, as opposed to enzymatic, methods. "Wholly" synthesized DNA sequences are therefore produced entirely by chemical means, and "partially" synthesized DNAs embrace those wherein only portions of the resulting DNA were produced by chemical means.

Genomic DNA of the invention comprises the protein coding region for a polypeptide of the invention and is also intended to include allelic variants thereof. It is widely understood that, for many genes, genomic DNA is transcribed into RNA transcripts that undergo one or more splicing events wherein intron (*i.e.*, non-coding regions) of the transcripts are removed, or "spliced out." RNA transcripts that can be spliced by alternative mechanisms, and therefore be subject to removal of different RNA sequences but still encode a GPCR polypeptide of the present invention, are referred to in the art as splice variants which are embraced by the invention. Splice variants comprehended by the invention therefore are encoded by the same original genomic DNA sequences but arise from distinct mRNA transcripts. Allelic variants are modified forms of a wild type gene sequence, the modification

resulting from recombination during chromosomal segregation or exposure to conditions which give rise to genetic mutation. Allelic variants, like wild type genes, are naturally occurring sequences (as opposed to non-naturally occurring variants which arise from *in vitro* manipulation).

5                   The invention also comprehends cDNA that is obtained through reverse transcription of an RNA polynucleotide encoding a GPCR of the present invention (conventionally followed by second strand synthesis of a complementary strand to provide a double-stranded DNA).

                  A preferred DNA sequence encoding a human GPCR polypeptide is set out in SEQ ID NO: 1, wherein nucleotides 157 to 1122 represent the CON193 coding sequence, with termination codon (surrounded by upstream and downstream untranslated sequences). Another preferred DNA sequence encoding a human GPCR polypeptide is set out in SEQ ID NO: 3, wherein nucleotides 1 to 1014 represent the CON166 coding sequence and stop codon. Still another preferred DNA sequence encoding a human GPCR polypeptide is set out in SEQ ID NO: 5, wherein nucleotides 691 to 1845 represent the CON103 coding sequence with stop codon (surrounded by upstream and downstream untranslated sequences). Another preferred DNA sequence encoding a human GPCR polypeptide is set out in SEQ ID NO: 7, wherein nucleotides 146 to 1147 represent the CON203 coding sequence with stop codon (surrounded by upstream and downstream untranslated sequences). A preferred DNA sequence encoding a human GPCR polypeptide is set out in SEQ ID NO: 9, wherein nucleotides 1 to 957 represent the CON198 coding sequence with stop codon. Another preferred DNA sequence encoding a human GPCR polypeptide is set out in SEQ ID NO: 11, wherein nucleotides 1 to 924 represent the CON197 coding sequence with stop codon (followed by downstream untranslated sequences). A preferred DNA sequence encoding a human GPCR polypeptide is set out in SEQ ID NO: 13, wherein nucleotides 266 to 1378 represent the CON202 coding sequence and termination codon (surrounded by upstream and downstream untranslated sequences). A preferred DNA sequence encoding a human GPCR polypeptide is set out in SEQ ID NO: 15, wherein nucleotides 1 to 1191 represent the CON222 coding sequence and termination codon. A preferred DNA sequence encoding a human GPCR polypeptide

- 43 -

is set out in SEQ ID NO: 17, wherein nucleotides 13 to 1089 represent the CON215 coding sequence and termination codon (surrounded by upstream and downstream untranslated sequences). A preferred DNA sequence encoding a human GPCR polypeptide is set out in SEQ ID NO: 19, wherein nucleotides 42 to 1157 represent  
5 the CON217 coding sequence (surrounded by upstream and downstream untranslated sequences). The foregoing sequences without their termination codons also comprise preferred sequences.

The worker of skill in the art will readily appreciate that the preferred DNA of the invention comprises a double stranded molecule, for example the  
10 molecule having any one of the sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, or 19 (or coding portions thereof) along with the complementary molecule (the "non-coding strand" or "complement") having a sequence deducible from the sequence of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, or 19 according to Watson-Crick base pairing rules for DNA. Also preferred are other polynucleotides encoding  
15 the GPCR polypeptides of the invention set forth in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18 and 20 which differ in sequence from the polynucleotide of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, or 19, respectively, by virtue of the well-known degeneracy of the universal genetic code.

The invention further embraces species, preferably mammalian,  
20 homologs of the human GPCR DNAs. Species homologs, sometimes referred to as "orthologs," in general, share at least 35%, at least 40%, at least 45%, at least 50%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99% homology with human DNA of the invention. Percent sequence "homology" with respect to polynucleotides of the  
25 invention is defined herein as the percentage of nucleotide bases in the candidate sequence that are identical to nucleotides in the GPCR sequence set forth in any one of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, or 19 after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity.

The polynucleotide sequence information provided by the invention  
30 makes possible large scale expression of the encoded polypeptide by techniques well known and routinely practiced in the art. Polynucleotides of the invention also permit

identification and isolation of polynucleotides encoding related GPCR polypeptides, such as human allelic variants and species homologs, by well known techniques including Southern and/or Northern hybridization, and polymerase chain reaction (PCR). Examples of related polynucleotides include human and non-human genomic sequences, including allelic variants, as well as polynucleotides encoding polypeptides homologous to GPCR polypeptides and structurally related the polypeptides sharing one or more biological, immunological, and/or physical properties of the GPCR polypeptides. Non-human species genes encoding proteins homologous to GPCR polypeptides can also be identified by Southern and/or PCR analysis and are useful in animal models for GPCR-related disorders. Knowledge of the sequence of a human GPCR DNA also makes possible, through use of Southern hybridization or polymerase chain reaction (PCR), the identification of genomic DNA sequences encoding GPCR expression control regulatory sequences such as promoters, operators, enhancers, repressors, and the like. Polynucleotides of the invention are also useful in hybridization assays to detect the capacity of cells to express GPCR polypeptides. Polynucleotides of the invention may also be the basis for diagnostic methods useful for identifying a genetic alteration(s) in a GPCR locus that underlies a disease state or states, which information is useful both for diagnosis and for selection of therapeutic strategies.

The disclosure herein of full length polynucleotides encoding GPCR polypeptides of the present invention makes readily available to the worker of ordinary skill in the art every possible fragment of the full length polynucleotides. The invention therefore provides fragments of GPCR-encoding polynucleotides comprising at least 14-15, and preferably at least 18, 20, 25, 50, or 75 consecutive nucleotides of a polynucleotide encoding GPCR polypeptides. Preferably, fragment polynucleotides of the invention comprise sequences unique to the GPCR-encoding polynucleotide sequence, and therefore hybridize under highly stringent or moderately stringent conditions only (*i.e.*, "specifically") to polynucleotides encoding GPCR polypeptides (or fragments thereof). Polynucleotide fragments of genomic sequences of the invention comprise not only sequences unique to the coding region, but also include fragments of the full length sequence derived from introns, regulatory regions,

and/or other non-translated sequences. Sequences unique to polynucleotides of the invention are recognizable through sequence comparison to other known polynucleotides, and can be identified through use of alignment programs routinely utilized in the art, *e.g.*, those made available in public sequence databases. Such sequences also are recognizable from Southern and Northern hybridization analyses to determine the number of fragments of genomic DNA and RNA to which a polynucleotide will hybridize. Polynucleotides of the invention can be labeled in a manner that permits their detection, including radioactive, fluorescent, and enzymatic labeling.

Fragment polynucleotides are particularly useful as probes for detection of full length or other fragment GPCR polynucleotides. One or more fragment polynucleotides can be included in kits that are used to detect the presence of a polynucleotide encoding a GPCR polypeptide, or used to detect variations in a polynucleotide sequences encoding GPCR polypeptides.

The invention also embraces DNAs encoding GPCR polypeptides which DNAs hybridize under moderately stringent or high stringency conditions to the non-coding strand, or complement, of the polynucleotide in any one of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17 or 19.

Exemplary highly stringent hybridization conditions are as follows: hybridization at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% Dextran sulfate, and washing twice for 30 minutes at 60°C in a wash solution comprising 0.1x SSC and 1% SDS. It is understood in the art that conditions of equivalent stringency can be achieved through variation of temperature and buffer, or salt concentration as described Ausubel, *et al.* (Eds.), *Protocols in Molecular Biology*, John Wiley & Sons (1994), pp. 6.0.3 to 6.4.10. Modifications in hybridization conditions can be empirically determined or precisely calculated based on the length and the percentage of guanosine/cytosine (GC) base pairing of the probe. The hybridization conditions can be calculated as described in Sambrook *et al.*, (Eds.), *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press: Cold Spring Harbor, New York (1989), pp. 9.47 to 9.51.

- 46 -

Autonomously replicating recombinant expression constructs such as plasmid and viral DNA vectors incorporating polynucleotides of the invention are also provided. Expression constructs wherein GPCR-encoding polynucleotides are operatively linked to an endogenous or exogenous expression control DNA sequence and a transcription terminator are also provided. Expression control DNA sequences include promoters, enhancers, and operators, and are generally selected based on the expression systems in which the expression construct is to be utilized. Preferred promoter and enhancer sequences are generally selected for the ability to increase gene expression, while operator sequences are generally selected for the ability to regulate gene expression. Expression constructs of the invention may also include sequences encoding one or more selectable markers that permit identification of host cells bearing the construct. Expression constructs may also include sequences that facilitate, and preferably promote, homologous recombination in a host cell. Preferred constructs of the invention also include sequences necessary for replication in a host cell.

Expression constructs are preferably utilized for production of an encoded protein, but also may be utilized simply to amplify GPCR-encoding polynucleotide sequences.

According to another aspect of the invention, host cells are provided, including prokaryotic and eukaryotic cells, comprising a polynucleotide of the invention (or vector of the invention) in a manner which permits expression of the encoded GPCR polypeptide. Polynucleotides of the invention may be introduced into the host cell as part of a circular plasmid, or as linear DNA comprising an isolated protein coding region or a viral vector. Methods for introducing DNA into the host cell well known and routinely practiced in the art include transformation, transfection, electroporation, nuclear injection, or fusion with carriers such as liposomes, micelles, ghost cells, and protoplasts. Expression systems of the invention include bacterial, yeast, fungal, plant, insect, invertebrate, and mammalian cells systems.

Host cells of the invention are a valuable source of immunogen for development of antibodies specifically immunoreactive with GPCR polypeptides. Host cells of the invention are also useful in methods for large scale production of

GPCR polypeptides wherein the cells are grown in a suitable culture medium and the desired polypeptide products are isolated from the cells or from the medium in which the cells are grown by purification methods known in the art, *e.g.*, conventional chromatographic methods including immunoaffinity chromatography, receptor  
5 affinity chromatography, hydrophobic interaction chromatography, lectin affinity chromatography, size exclusion filtration, cation or anion exchange chromatography, high pressure liquid chromatography (HPLC), reverse phase HPLC, and the like. Still other methods of purification include those wherein the desired protein is expressed and purified as a fusion protein having a specific tag, label, or chelating moiety that is  
10 recognized by a specific binding partner or agent. The purified protein can be cleaved to yield the desired protein, or be left as an intact fusion protein. Cleavage of the fusion component may produce a form of the desired protein having additional amino acid residues as a result of the cleavage process.

Knowledge of GPCR DNA sequences allows for modification of cells  
15 to permit, or increase, expression of endogenous GPCR. Cells can be modified (*e.g.*, by homologous recombination) to provide increased expression by replacing, in whole or in part, the naturally occurring GPCR promoter with all or part of a heterologous promoter so that the cells express GPCR polypeptides at higher levels. The heterologous promoter is inserted in such a manner that it is operatively linked to  
20 endogenous GPCR polypeptide encoding sequences. [See, for example, PCT International Publication No. WO 94/12650, PCT International Publication No. WO 92/20808, and PCT International Publication No. WO 91/09955.] It is also contemplated that, in addition to heterologous promoter DNA, amplifiable marker DNA (*e.g.*, *ada*, *dhfr*, and the multifunctional CAD gene which encodes carbamyl  
25 phosphate synthase, aspartate transcarbamylase, and dihydroorotase) and/or intron DNA may be inserted along with the heterologous promoter DNA. If linked to the GPCR coding sequence, amplification of the marker DNA by standard selection methods results in co-amplification of the GPCR coding sequences in the cells.

The DNA sequence information provided by the present invention also  
30 makes possible the development through, *e.g.* homologous recombination or "knock-out" strategies [Capecchi, *Science* 244: 1288-1292 (1989)], of animals that

fail to express functional GPCR polypeptides or that express a variant of GPCR polypeptides. Such animals (especially small laboratory animals such as rats, rabbits, and mice) are useful as models for studying the *in vivo* activities of GPCR polypeptides and modulators of GPCR polypeptides.

5                   Also made available by the invention are anti-sense polynucleotides which recognize and hybridize to polynucleotides encoding GPCR polypeptides. Full length and fragment anti-sense polynucleotides are provided. Fragment anti-sense molecules of the invention include those which specifically recognize and hybridize to GPCR RNA (as determined by sequence comparison of DNA encoding GPCR  
10 polypeptides to DNA encoding other known molecules). Identification of sequences unique to GPCR-encoding polynucleotides, can be deduced through use of any publicly available sequence database, and/or through use of commercially available sequence comparison programs. The uniqueness of selected sequences in an entire genome can be further verified by hybridization analyses. After identification of the  
15 desired sequences, isolation through restriction digestion or amplification using any of the various polymerase chain reaction techniques well known in the art can be performed. Antisense polynucleotides are particularly relevant to regulating expression of GPCR polypeptides by those cells expressing GPCR mRNA.

                  Antisense nucleic acids (preferably 10 to 20 base pair oligonucleotides)  
20 capable of specifically binding to GPCR expression control sequences or GPCR RNA are introduced into cells (*e.g.*, by a viral vector or colloidal dispersion system such as a liposome). The antisense nucleic acid binds to the GPCR target nucleotide sequence in the cell and prevents transcription or translation of the target sequence. Phosphorothioate and methylphosphonate antisense oligonucleotides are specifically  
25 contemplated for therapeutic use by the invention. The antisense oligonucleotides may be further modified by poly-L-lysine, transferrin polylysine, or cholesterol moieties at their 5' end. Suppression of GPCR polypeptide expression at either the transcriptional or translational level is useful to general cellular and/or animal models for diseases characterized by aberrant expression. Suppression of GPCR polypeptide  
30 expression at either the transcriptional or translational level is useful to generate

- 49 -

cellular animal models for diseases characterized by aberrant GPCR polypeptide expression.

The GPCR polynucleotide and polypeptide sequences taught in the present invention facilitate the design of novel transcription factors for modulating GPCR polypeptide expression in native cells and animals, and cells transformed or transfected with GPCR polynucleotides. For example, the Cys<sub>2</sub>-His<sub>2</sub> zinc finger proteins, which bind DNA via their zinc finger domains, have been shown to be amenable to structural changes that lead to the recognition of different target sequences. These artificial zinc finger proteins recognize specific target sites with high affinity and low dissociation constants, and are able to act as gene switches to modulate gene expression. Knowledge of the particular GPCR target sequence of the present invention facilitates the engineering of zinc finger proteins specific for the target sequence using known methods such as a combination of structure-based modeling and screening of phage display libraries [Segal *et al.*, *Proc Natl Acad Sci USA* 96: 2758-2763 (1999); Liu *et al.*, *Proc Natl Acad Sci USA* 94: 5525-30 (1997); Greisman and Pabo *Science* 275: 657-61 (1997); Choo *et al.*, *J Mol Biol* 273: 525-32 (1997)]. Each zinc finger domain usually recognizes three or more base pairs. Since a recognition sequence of 18 base pairs is generally sufficient in length to render it unique in any known genome, a zinc finger protein consisting of 6 tandem repeats of zinc fingers would be expected to ensure specificity for a particular sequence [Segal *et al.*, *Proc Natl Acad Sci USA* 96: 2758-2763 (1999)]. The artificial zinc finger repeats, designed based on GPCR polynucleotide sequences, are fused to activation or repression domains to promote or suppress GPCR polypeptides expression [Liu *et al.*, *Proc Natl Acad Sci USA* 94: 5525-30 (1997)]. Alternatively, the zinc finger domains can be fused to the TATA box-binding factor (TBP) with varying lengths of linker region between the zinc finger peptide and the TBP to create either transcriptional activators or repressors [Kim *et al.*, *Proc Natl Acad Sci USA* 94: 3616-3620 (1997)]. Such proteins, and polynucleotides that encode them, have utility for modulating GPCR polypeptide expression *in vivo* in both native cells, animals and humans; and/or cells transfected with GPCR polynucleotide-encoding sequences. The novel transcription factor can be delivered to the target cells by transfecting constructs that

- 50 -

express the transcription factor (gene therapy), or by introducing the protein.

Engineered zinc finger proteins can also be designed to bind RNA sequences for use in therapeutics as alternatives to antisense or catalytic RNA methods [McColl *et al.*, *Proc Natl Acad Sci USA* 96:9521-6 (1999); Wu *et al.*, *Proc Natl Acad Sci USA* 92:344-348 (1995)]. The present invention contemplates methods of designing such transcription factors based on the gene sequence of the invention, as well as customized zinc finger proteins, that are useful to modulate GPCR polypeptide expression in cells (native or transformed) whose genetic complement includes these sequences.

The invention also provides purified and isolated mammalian GPCR polypeptides encoded by a polynucleotide of the invention. Presently preferred is a human GPCR polypeptide comprising the amino acid sequence set out in any one of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18 or 20.

The invention also embraces polypeptides that have at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55% or at least 50% identity and/or homology to a preferred polypeptide of the invention. Percent amino acid sequence "identity" with respect to the preferred polypeptide of the invention is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the residues in a GPCR polypeptide sequence after aligning both sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Percent sequence "homology" with respect to the preferred polypeptide of the invention is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the residues in a GPCR sequence after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and also considering any conservative substitutions as part of the sequence identity.

In one aspect, percent homology is calculated as the percentage of amino acid residues in the smaller of two sequences which align with identical amino acid residue in the sequence being compared, when four gaps in a length of 100 amino

acids may be introduced to maximize alignment [Dayhoff, in *Atlas of Protein Sequence and Structure*, Vol. 5, p. 124, National Biochemical Research Foundation, Washington, D.C. (1972), incorporated herein by reference].

Polypeptides of the invention may be isolated from natural cell  
5 sources or may be chemically synthesized, but are preferably produced by recombinant procedures involving host cells of the invention. Use of mammalian host cells is expected to provide for such post-translational modifications (*e.g.*, glycosylation, truncation, lipidation, and phosphorylation) as may be needed to confer optimal biological activity on recombinant expression products of the invention.  
10 Glycosylated and non-glycosylated forms of GPCR polypeptides are embraced.

The invention also embraces variant (or analog) GPCR polypeptides. In one example, insertion variants are provided wherein one or more amino acid residues supplement a GPCR amino acid sequence. Insertions may be located at either or both termini of the protein, or may be positioned within internal regions of  
15 the GPCR amino acid sequence. Insertional variants with additional residues at either or both termini can include for example, fusion proteins and proteins including amino acid tags or labels.

Insertion variants include GPCR polypeptides wherein one or more amino acid residues are added to a GPCR amino acid sequence, or to a biologically  
20 active fragment thereof.

Variant products of the invention also include mature GPCR polypeptide products, *i.e.*, GPCR polypeptide products wherein leader or signal sequences are removed, with additional amino terminal residues. The additional amino terminal residues may be derived from another protein, or may include one or  
25 more residues that are not identifiable as being derived from a specific proteins. GPCR polypeptide products with an additional methionine residue at position -1 (Met<sup>-1</sup>-GPCR) are contemplated, as are variants with additional methionine and lysine residues at positions -2 and -1 (Met<sup>-2</sup>-Lys<sup>-1</sup>-GPCR). Variants of GPCR polypeptide with additional Met, Met-Lys, Lys residues (or one or more basic residues in general)  
30 are particularly useful for enhanced recombinant protein production in bacterial host cell.

- 52 -

The invention also embraces GPCR polypeptide variants having additional amino acid residues which result from use of specific expression systems. For example, use of commercially available vectors that express a desired polypeptide as part of glutathione-S-transferase (GST) fusion product provides the desired polypeptide having an additional glycine residue at position -1 after cleavage of the GST component from the desired polypeptide. Variants which result from expression in other vector systems are also contemplated.

Insertional variants also include fusion proteins wherein the amino and/or carboxy termini of a GPCR polypeptide is fused to another polypeptide.

In another aspect, the invention provides deletion variants wherein one or more amino acid residues in a GPCR polypeptide are removed. Deletions can be effected at one or both termini of the GPCR polypeptide, or with removal of one or more residues within the GPCR amino acid sequence. Deletion variants, therefore, include all fragments of a GPCR polypeptide.

The invention also embraces polypeptide fragments of the sequence set out in SEQ ID NO: 2 wherein the fragments maintain biological (e.g., ligand binding and/or intracellular signaling) or immunological properties of a GPCR polypeptide. Fragments comprising at least 5, 10, 15, 20, 25, 30, 35, or 40 consecutive amino acids of SEQ ID NO: 2 are comprehended by the invention. Preferred polypeptide fragments display antigenic properties unique to or specific for human GPCR and its allelic and species homologs. Fragments of the invention having the desired biological and immunological properties can be prepared by any of the methods well known and routinely practiced in the art.

In still another aspect, the invention provides substitution variants of GPCR polypeptides. Substitution variants include those polypeptides wherein one or more amino acid residues of a GPCR polypeptide are removed and replaced with alternative residues. In one aspect, the substitutions are conservative in nature, however, the invention embraces substitutions that are also non-conservative. Conservative substitutions for this purpose may be defined as set out in Tables A, B, or C below.

- 53 -

Variant polypeptides include those wherein conservative substitutions have been introduced by modification of polynucleotides encoding polypeptides of the invention. Amino acids can be classified according to physical properties and contribution to secondary and tertiary protein structure. A conservative substitution is recognized in the art as a substitution of one amino acid for another amino acid that has similar properties. Exemplary conservative substitutions are set out in Table A (from WO 97/09433, page 10, published March 13, 1997 (PCT/GB96/02197, filed 9/6/96), immediately below.

10

**Table A****Conservative Substitutions I**

	<b>SIDE CHAIN</b>	
	<b><u>CHARACTERISTIC</u></b>	<b><u>AMINO ACID</u></b>
15	Aliphatic	
	Non-polar	G A P I L V
	Polar - uncharged	C S T M N Q
	Polar - charged	D E K R
	Aromatic	H F W Y
20	Other	N Q D E

Alternatively, conservative amino acids can be grouped as described in Lehninger, [Biochemistry, Second Edition; Worth Publishers, Inc. NY:NY (1975), pp.71-77] as set out in Table B, immediately below.

25

**Table B**  
**Conservative Substitutions II**

5	<b>SIDE CHAIN</b>	
	<b><u>CHARACTERISTIC</u></b>	<b><u>AMINO ACID</u></b>
	Non-polar (hydrophobic)	
	A. Aliphatic:	A L I V P
10	B. Aromatic:	F W
	C. Sulfur-containing:	M
	D. Borderline:	G
	Uncharged-polar	
	A. Hydroxyl:	S T Y
15	B. Amides:	N Q
	C. Sulfhydryl:	C
	D. Borderline:	G
	Positively Charged (Basic):	
	Negatively Charged (Acidic):	
20		K R H
		D E

As still an another alternative, exemplary conservative substitutions are set out in Table C, immediately below.

**Table C**  
**Conservative Substitutions III**

	<u>Original</u>	<u>Exemplary Substitution</u>
5	<u>Residue</u>	
	Ala (A)	Val, Leu, Ile
	Arg (R)	Lys, Gln, Asn
	Asn (N)	Gln, His, Lys, Arg
	Asp (D)	Glu
10	Cys (C)	Ser
	Gln (Q)	Asn
	Glu (E)	Asp
	His (H)	Asn, Gln, Lys, Arg
	Ile (I)	Leu, Val, Met, Ala, Phe,
15	Leu (L)	Ile, Val, Met, Ala, Phe
	Lys (K)	Arg, Gln, Asn
	Met (M)	Leu, Phe, Ile
	Phe (F)	Leu, Val, Ile, Ala
	Pro (P)	Gly
20	Ser (S)	Thr
	Thr (T)	Ser
	Trp (W)	Tyr
	Tyr (Y)	Trp, Phe, Thr, Ser
25	Val (V)	Ile, Leu, Met, Phe, Ala

GPCR polypeptide variants that display ligand binding properties of native GPCR polypeptides and are expressed at higher levels, and variants that provide for constitutive active receptor are particularly useful in assays of the

- 56 -

invention. Such variants also are useful in cellular and animal models for diseases characterized by aberrant GPCR polypeptide expression/activity.

It should be understood that the definition of polypeptides of the invention is intended to include polypeptides bearing modifications other than insertion, deletion, or substitution of amino acid residues. By way of example, the modifications may be covalent in nature, and include for example, chemical bonding with polymers, lipids, other organic, and inorganic moieties. Such derivatives may be prepared to increase circulating half-life of a polypeptide, or may be designed to improve targeting capacity for the polypeptide to desired cells, tissues, or organs.

Similarly, the invention further embraces GPCR polypeptides that have been covalently modified to include one or more water soluble polymer attachments such as polyethylene glycol, polyoxyethylene glycol, or polypropylene glycol.

In a related embodiment, the present invention provides compositions comprising purified polypeptides of the invention. Preferred compositions comprise, in addition to the polypeptide of the invention, a pharmaceutically acceptable (*i.e.*, sterile and non-toxic) liquid, semisolid, or solid diluents that serve as pharmaceutical vehicles, excipients, or media. Any diluent known in the art may be used. Exemplary diluents include, but are not limited to, water, saline solutions, polyoxyethylene sorbitan monolaurate, magnesium stearate, methyl- and propylhydroxybenzoate, talc, alginates, starches, lactose, sucrose, dextrose, sorbitol, mannitol, glycerol, calcium phosphate, mineral oil, and cocoa butter.

Also comprehended by the present invention are antibodies (*e.g.*, monoclonal and polyclonal antibodies, single chain antibodies, chimeric antibodies, bifunctional/bispecific antibodies, humanized antibodies, human antibodies, and complementary determining region (CDR)-grafted antibodies, including compounds which include CDR sequences which specifically recognize a polypeptide of the invention) specific for GPCR polypeptides of the invention or fragments thereof. Preferred antibodies of the invention are human antibodies which can be produced and identified according to methods described in WO93/11236, published June 20, 1993, which is incorporated herein by reference in its entirety. Antibody fragments, including Fab, Fab', F(ab')<sub>2</sub>, and F<sub>v</sub>, are also provided by the invention. The term

“specific for,” when used to describe antibodies of the invention, indicates that the variable regions of the antibodies of the invention recognize and bind GPCR polypeptides exclusively (*i.e.*, able to distinguish GPCR polypeptides from other known GPCR polypeptides by virtue of measurable differences in binding affinity, despite the possible existence of localized sequence identity, homology, or similarity between GPCR polypeptides and such polypeptides). It will be understood that specific antibodies may also interact with other proteins (for example, *S. aureus* protein A or other antibodies in ELISA techniques) through interactions with sequences outside the variable region of the antibodies, and in particular, in the constant region of the molecule. Screening assays to determine binding specificity of an antibody of the invention are well known and routinely practiced in the art. For a comprehensive discussion of such assays, see Harlow *et al.* (Eds), *Antibodies A Laboratory Manual*; Cold Spring Harbor Laboratory; Cold Spring Harbor, NY (1988), Chapter 6. Antibodies that recognize and bind fragments of the GPCR polypeptides of the invention are also contemplated, provided that the antibodies are, first and foremost, specific for GPCR polypeptides. Antibodies of the invention can be produced using any method well known and routinely practiced in the art.

Non-human antibodies may be humanized by any methods known in the art. In one method, the non-human CDRs are inserted into a human antibody or consensus antibody framework sequence. Further changes can then be introduced into the antibody framework to modulate affinity or immunogenicity.

Antibodies of the invention are useful for, for example, therapeutic purposes (by modulating activity of GPCR polypeptides), diagnostic purposes to detect or quantitate GPCR polypeptides, as well as purification of GPCR polypeptides. Kits comprising an antibody of the invention for any of the purposes described herein are also comprehended. In general, a kit of the invention also includes a control antigen for which the antibody is immunospecific.

Specific binding molecules, including natural ligands and synthetic compounds, can be identified or developed using isolated or recombinant GPCR polypeptide products, GPCR polypeptide variants, or preferably, cells expressing such products. Binding partners are useful for purifying GPCR polypeptide products and

- 58 -

detection or quantification of GPCR polypeptide products in fluid and tissue samples using known immunological procedures. Binding molecules are also manifestly useful in modulating (*i.e.*, blocking, inhibiting or stimulating) biological activities of GPCR polypeptides, especially those activities involved in signal transduction.

5           The DNA and amino acid sequence information provided by the present invention also makes possible identification of binding partner compounds with which a GPCR polypeptide or polynucleotide will interact. Methods to identify binding partner compounds include solution assays, *in vitro* assays wherein GPCR polypeptides are immobilized, and cell based assays. Identification of binding partner  
10       compounds of GPCR polypeptides provides candidates for therapeutic or prophylactic intervention in pathologies associated with GPCR polypeptide normal and aberrant biological activity.

          The invention includes several assay systems for identifying GPCR polypeptide binding partners. In solution assays, methods of the invention comprise  
15       the steps of (a) contacting a GPCR polypeptide with one or more candidate binding partner compounds and (b) identifying the compounds that bind to the GPCR polypeptide. Identification of the compounds that bind the GPCR polypeptide can be achieved by isolating the GPCR polypeptide/binding partner complex, and separating  
20       the GPCR polypeptide from the binding partner compound. An additional step of characterizing the physical, biological, and/or biochemical properties of the binding partner compound is also comprehended in another embodiment of the invention. In one aspect, the GPCR polypeptide/binding partner complex is isolated using a antibody immunospecific for either the GPCR polypeptide or the candidate binding partner compound.

25           In still other embodiments, either the GPCR polypeptide or the candidate binding partner compound comprises a label or tag that facilitates its isolation, and methods of the invention to identify binding partner compounds include a step of isolating the GPCR polypeptide/binding partner complex through interaction with the label or tag. An exemplary tag of this type is a poly-histidine sequence,  
30       generally around six histidine residues, that permits isolation of a compound so labeled using nickel chelation. Other labels and tags, such as the FLAG<sup>®</sup> tag

- 59 -

(Eastman Kodak, Rochester, NY), well known and routinely used in the art, are embraced by the invention.

In one variation of an *in vitro* assay, the invention provides a method comprising the steps of (a) contacting an immobilized GPCR polypeptide with a candidate binding partner compound and (b) detecting binding of the candidate compound to GPCR polypeptide. In an alternative embodiment, the candidate binding partner compound is immobilized and binding of GPCR polypeptide is detected. Immobilization is accomplished using any of the methods well known in the art, including covalent bonding to a support, a bead, or a chromatographic resin, as well as non-covalent, high affinity interaction such as antibody binding, or use of streptavidin/biotin binding wherein the immobilized compound includes a biotin moiety. Detection of binding can be accomplished (i) using a radioactive label on the compound that is not immobilized, (ii) using a fluorescent label on the non-immobilized compound, (iii) using an antibody immunospecific for the non-immobilized compound, (iv) using a label on the non-immobilized compound that excites a fluorescent support to which the immobilized compound is attached, as well as other techniques well known and routinely practiced in the art.

The invention also provides cell-based assays to identify binding partner compounds of a GPCR polypeptide. In one embodiment, the invention provides a method comprising the steps of contacting a GPCR polypeptide expressed on the surface of a cell with a candidate binding partner compound and detecting binding of the candidate binding partner compound to the GPCR polypeptide. In a preferred embodiment, the detection comprises detecting a calcium flux or other physiological cellular events caused by the binding of the molecule.

Agents that modulate (*i.e.*, increase, decrease, or block) GPCR polypeptide activity or expression may be identified by incubating a putative modulator with a cell expressing a GPCR polypeptide or polynucleotide and determining the effect of the putative modulator on GPCR polypeptide activity or expression. The selectivity of a compound that modulates the activity of GPCR polypeptides can be evaluated by comparing its effects on GPCR polypeptides to its effect on other G coupled-protein receptor compounds. Selective modulators may

- 60 -

include, for example, antibodies and other proteins, peptides, or organic molecules which specifically bind to a G coupled-protein receptor polypeptide or a G coupled-protein receptor-encoding nucleic acid. Modulators of GPCR polypeptide activity will be therapeutically useful in treatment of diseases and physiological conditions in which normal or aberrant GPCR polypeptide activity is involved.

Methods of the invention to identify modulators include variations on any of the methods described above to identify binding partner compounds, the variations including techniques wherein a binding partner compound has been identified and the binding assay is carried out in the presence and absence of a candidate modulator. A modulator is identified in those instances where binding between the GPCR polypeptide and the binding partner compound changes in the presence of the candidate modulator compared to binding in the absence of the candidate modulator compound. A modulator that increases binding between the GPCR polypeptide and the binding partner compound is described as an enhancer or activator, and a modulator that decreases binding between the GPCR polypeptide and the binding partner compound is described as an inhibitor.

The invention also comprehends high throughput screening (HTS) assays to identify compounds that interact with or inhibit biological activity (*i.e.*, inhibit enzymatic activity, binding activity, *etc.*) of a GPCR polypeptide. HTS assays permit screening of large numbers of compounds in an efficient manner. Cell-based HTS systems are contemplated to investigate GPCR receptor-ligand interaction. HTS assays are designed to identify "hits" or "lead compounds" having the desired property, from which modifications can be designed to improve the desired property. Chemical modification of the "hit" or "lead compound" is often based on an identifiable structure/activity relationship between the "hit" and the GPCR polypeptide.

Mutations in the GPCR gene that result in loss of normal function of the GPCR gene product underlie GPCR polypeptide-related human disease states. The invention comprehends gene therapy to restore activity to treat those disease states. Delivery of a functional GPCR gene to appropriate cells is effected *ex vivo*, *in situ*, or *in vivo* by use of vectors, and more particularly viral vectors (*e.g.*, adenovirus,

- 61 -

adeno-associated virus, or a retrovirus), or *ex vivo* by use of physical DNA transfer methods (e.g., liposomes or chemical treatments). See, for example, Anderson, *Nature*, supplement to vol. 392, no. 6679, pp.25-20 (1998). For additional reviews of gene therapy technology see Friedmann, *Science*, 244: 1275-1281 (1989); Verma, *Scientific American*: 68-84 (1990); and Miller, *Nature*, 357: 455-460 (1992).

Alternatively, it is contemplated that in other human disease states, preventing the expression of or inhibiting the activity of GPCR polypeptides of the invention will be useful in treating the disease states. It is contemplated that antisense therapy or gene therapy could be applied to negatively regulate the expression of GPCR polypeptides of the invention.

Additional features of the invention will be apparent from the following Examples.

## EXAMPLE 1

### Cloning of G Protein-Coupled Receptors

The Incyte and Genbank expressed sequence tag (EST) databases were searched with the NCBI program Blastall using either the transmembrane VI region of known dopamine receptors (leading to the identification of CON193, CON166, CON103 and CON 203) or all known GPCR's *except* olfactory and opsin receptors (leading to the identification of CON198, CON197, CON202, CON222, CON215) as query sequences, to find patterns suggestive of novel G protein-coupled receptors. Positive hits from the find-pattern program were further analyzed with the GCG program BLAST to determine which ones were the most likely candidates to encode a GPCR, using the standard (default) alignment produced by BLAST as a guide.

#### A. Cloning of CON193 G Protein-Coupled Receptor

##### A.1. Database Search Results

Searching identified Clone 3091220H1 in the Incyte database as an interesting candidate sequence. The 3091220H1 Clone was obtained and sequenced directly using an ABI377 fluorescence-based sequencer (Perkin-Elmer/Applied Biosystems Division, PE/ABD, Foster City, CA) and the ABI PRISM™ Ready

- 62 -

Dye-Deoxy Terminator kit with Taq FSTM polymerase. Each ABI cycle sequencing reaction contained about 0.5 µg of plasmid DNA. Cycle-sequencing was performed using an initial denaturation at 98°C for 1 minute, followed by 50 cycles using the following parameters: 98°C for 30 seconds, annealing at 50°C for 30 seconds, and extension at 60°C for 4 minutes. Temperature cycles and times were controlled by a Perkin-Elmer 9600 thermocycler. Extension products were purified using Centriflex™ gel filtration cartridges (Advanced Genetic Technologies Corp., Gaithersburg, MD). Each reaction product was loaded by pipette onto the column, which was then centrifuged in a swinging bucket centrifuge (Sorvall model RT6000B tabletop centrifuge) at 1500 x g for 4 minutes at room temperature. Column-purified samples were dried under vacuum for about 40 minutes and then dissolved in 5 µl of a DNA loading solution (83% deionized formamide, 8.3 mM EDTA, and 1.6 mg/ml Blue Dextran). The samples were then heated to 90°C for three minutes and loaded into the gel sample wells for sequence analysis using the ABI377 sequencer.

Sequence analysis was done by importing ABI377 files into the Sequencer program (Gene Codes, Ann Arbor, MI). Generally, sequence reads of 700 bp were obtained. Potential sequencing errors were minimized by obtaining sequence information from both DNA strands and by re-sequencing difficult areas using primers annealing at different locations until all sequencing ambiguities were removed.

From the sequence it was deduced that Clone 3091220H1 contained only an amino-terminal fragment of a putative GPCR corresponding to the third through the seventh transmembrane regions (3TM-7TM) of a GPCR. Referring to SEQ ID NO: 1, the nucleotide sequence of Clone 3091220H1 corresponds to nucleotides 404 to 1308 of what was eventually determined to be the complete sequence of a novel seven-transmembrane receptor designated CON193. A database search with this partial sequence showed a 56% match to members of the olfactory receptor gene family, *e.g.*, the gene encoding mouse odorant receptor S19.

#### **A.2 Screening of a Genomic Phage Library to Obtain a Full-Length GPCR Clone:**

The PCR technique was used to prepare a genomic fragment for use as a probe specific for the genomic CON193 Clone. Based on the complete sequence of Clone 3091220H1, two oligonucleotide primers were designed: Primer LW1282: 5'-

TAATACCTGCACTGCCCCAC-3' (SEQ ID NO: 21; see nucleotides 876-894 of SEQ ID NO:1) and Primer LW1283: 5'-TCTTTCCTTCTCTTCTCACTCC-3' (SEQ ID NO: 22 see nucleotides 1137-1158 of SEQ ID NO:1). These primers were designed to amplify a 283 base-pair fragment of genomic DNA containing a portion of the CON193 coding region found in Clone 3091220H1 (assuming the absence of introns in this region).

Initially, a suitable human genomic library constructed in EMBL3 SP6/T7 (Clontech Laboratories) was amplified to provide the materials required for screening. Two microliters of the human genomic library (approximately  $10^8$  plaque-forming units per milliliter; Clontech Laboratories, catalog number HL1067J) were added to 6 ml of an overnight culture of K802 cells (Clontech Laboratories), and 250  $\mu$ l aliquots were distributed into each of 24 tubes. The tubes were incubated at 37°C for 15 minutes, and then 7 ml of 0.8% agarose (*i.e.*, top agarose) at 50°C were added to each tube. After mixing, the contents of the tubes were poured onto 150 mm LB plates and incubated overnight at 37°C to allow clone amplification, evident as plaque formation (typically, confluent lysis was observed rather than discrete plaques). To each plate, 5 ml of SM phage buffer (0.1 M NaCl, 8.1  $\mu$ M  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 50 mM Tris-HCl (pH 7.5), and 0.0001 % gelatin) was added and the top agarose was removed by scraping with a microscope slide. Top agarose slurries containing phage were then placed in individual 50 ml centrifuge tubes. A drop of chloroform was added and each tube was placed in a 37°C shaker for 15 minutes, followed by centrifuging at 2,750 x g for 15 minutes. The supernatants were isolated and separately stored at 4°C as 24 stock solutions of amplified library clones.

As noted above, polymerase chain reaction (PCR) was selected as a technique for screening the phage library. Each PCR reaction was done in a 20  $\mu$ l reaction volume containing 8.84  $\mu$ l  $\text{H}_2\text{O}$ , 2  $\mu$ l 10X PCR buffer II (Perkin-Elmer), 2  $\mu$ l 25 mM  $\text{MgCl}_2$ , 0.8  $\mu$ l dNTP mixture (dATP, dCTP, dGTP, dCTP, each at 10 mM), 0.12  $\mu$ l primer LW1282 (approximately 1  $\mu$ g/ $\mu$ l), 0.12  $\mu$ l primer LW1283 (approximately 1  $\mu$ g/ $\mu$ l), 0.12  $\mu$ l AmpliTaq Gold polymerase (5 Units/ $\mu$ l, with "Units" as defined by the supplier, Perkin-Elmer) and 2  $\mu$ l of phage from one of the 24 stock tubes. The PCR reaction involved 1 cycle at 95°C for 10 minutes and 80°C for 20

- 64 -

minutes, followed by 22 cycles at 95°C for 30 seconds, 72-51°C for 2 minutes (72°C for this stage of the second cycle, with a decrease of one degree for this stage in each succeeding cycle), 72°C for one minute, followed by 30 cycles at 95°C for 15 seconds, 50°C for 30 seconds, and 72°C for one minute.

5                   Following PCR cycling, the contents from each reaction tube were loaded onto a 2% agarose gel and electrophoresed adjacent to known size standards to screen for PCR products of the expected size, indicative of a clone containing the 283 bp portion of Clone 3091220H1 amplified by the two selected primers. A positive signal (*i.e.*, a fragment of the expected size) was found in one of the 24 PCR  
10 reactions, thereby identifying a single stock genomic library tube containing positive clones.

                  From the original genomic library tube that had given a PCR product of the correct size, a 5 µl phage aliquot was used to establish a set of five serial dilutions (1/100, v/v) that were plated and incubated in the same manner as described  
15 for the amplification of the phage library. Following incubation, BA85 nitrocellulose filters (Schleicher & Schuell) were placed on top of each of the plates for 1 hour to adsorb phage from the plaques that had formed in the top agarose during incubation. Each filter was then gently removed, placed phage side up in an individual petri dish, and covered with 4 ml of SM buffer for 15 minutes to elute the phage. One milliliter  
20 of SM containing eluted phage was removed from each plate and used to set up a PCR reaction as described above. The plate containing the most dilute phage solution to yield a PCR product of the expected size was then subdivided using the following procedure. A BA85 filter was placed on the top agar of the plate and the medium with applied filter was physically divided into 24 sections. After one hour to allow  
25 phage adsorption to the 24 filters, each filter was removed and separately incubated in 1 ml of SM buffer at room temperature for 15 minutes. Two microliters of each eluted phage solution were then used as a PCR substrate. Those plate sections yielding positive PCR results were then subdivided into 12 subsections by removing the top agar and incubating it in 200 µl of SM buffer for one hour at room  
30 temperature. Again, 2 µl of the eluted phage solutions were plated and lifted using BA85 filters, and PCR reactions were repeated. The procedure for progressive

- 65 -

dilution of phage was continued until a single plaque was isolated. Subsequently, 10  $\mu$ l of eluted phage from that single plaque were added to 100  $\mu$ l SM and 200  $\mu$ l of K802 cells for plating in a single petri dish as described above. A total of 7 plates were inoculated in this manner. Following incubation at 37°C for 16 hours, the top agarose from each of the 7 plates was removed to recover the phage, which were used to prepare purified genomic phage DNA using the Qiagen Lambda Midi Kit.

The purified CON193 genomic phage DNA was sequenced using the ABI PRISM™ 310 Genetic Analyzer (Perkin-Elmer/Applied Biosystems) which uses advanced capillary electrophoresis technology and the ABI PRISM™ BigDye™

Terminator Cycle Sequencing Ready Reaction Kit. The cycle-sequencing reaction contained 18  $\mu$ l of H<sub>2</sub>O, 16  $\mu$ l of BigDye™ Terminator mix, 3  $\mu$ l of genomic phage DNA (0.26  $\mu$ g/ $\mu$ l), and 3  $\mu$ l primer (25 ng/ $\mu$ l). The reaction was performed in a Perkin-Elmer 9600 thermocycler at 95°C for 5 minutes, followed by 75 cycles of 95°C for 30 seconds, 55°C for 20 seconds, and 60°C for 4 minutes. The final subclone was also sequenced using the ABI PRISM™ 310 Genetic Analyzer. The cycle-sequencing reaction contained 6  $\mu$ l of H<sub>2</sub>O, 8  $\mu$ l of BigDye™ Terminator mix, 5  $\mu$ l of miniprep clone DNA (0.1  $\mu$ g/ $\mu$ l), and 1  $\mu$ l primer (25 ng/ $\mu$ l). The reaction was performed in a Perkin-Elmer 9600 thermocycler at 25 cycles of 96°C for 10 seconds, 50°C for 10 seconds, and 60°C for 4 minutes. The product of the PCR reaction was purified using Centriflex™ gel filtration cartridges, dried under vacuum, and dissolved in 16  $\mu$ l of Template Suppression Reagent (PE-Applied Biosystems). The samples were then incubated at 95°C for 5 minutes and placed in the 310 Genetic Analyzer. These efforts resulted in the determination of the CON193 polynucleotide sequence set forth in SEQ ID NO:1 and the deduced amino acid sequence of the encoded CON193 polypeptide which is set forth in SEQ ID NO:2.

### A.3 Subcloning of the Coding Region of CON193 via PCR

Additional experiments were conducted to subclone the coding region of CON193 and place the isolated coding region into a useful vector. Two additional PCR primers were designed based on the coding region of CON193. The first PCR primer, designated Primer LW1373, has the sequence 5'-GCATAAGCTTATGCTA-ACACTGAATAAAACAG-3' (SEQ ID NO: 23), nucleotides 11-32 of which

- 66 -

correspond to nucleotides 157-178 of SEQ ID NO: 1. The second PCR primer is Primer LW1374, which has the sequence 5'-GCATCTCGAGTCACA-TGCTGTAGGATTTGG-3' (SEQ ID NO: 24, nucleotides 11-30 of which correspond to the complement of nucleotides 1102-1121 of SEQ ID NO: 1. To protect against  
5 exonucleolytic attack during subsequent exposure to enzymes, *e.g.*, Taq polymerase, primers were routinely synthesized with a protective run of nucleotides at the 5' end that were not necessarily complementary to the desired target.

PCR was performed in a 50  $\mu$ l reaction containing 35  $\mu$ l H<sub>2</sub>O, 5  $\mu$ l 10X TT buffer (140 mM ammonium sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5  $\mu$ l  
10 15 mM MgSO<sub>4</sub>, 2  $\mu$ l dNTP mixture (dGTP, dATP, dTTP, and dCTP, each at 10 mM), 2  $\mu$ l genomic phage DNA (0.26  $\mu$ g/ $\mu$ l), 0.3  $\mu$ l Primer LW1373 (1  $\mu$ g/ $\mu$ l), 0.3  $\mu$ l Primer LW1374 (1  $\mu$ g/ $\mu$ l), 0.4  $\mu$ l High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction was started with 1 cycle of 94°C for 2 minutes; followed by 15 cycles at 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1.3  
15 minutes.

The contents from the PCR reaction were loaded onto a 2% agarose gel, fractionated and electroeluted. The DNA band of expected size was excised from the gel, placed in a GenElute Agarose spin column (Supelco) and spun for 10 minutes at maximum speed in a microcentrifuge. The eluted DNA was precipitated with  
20 ethanol and resuspended in 6  $\mu$ l H<sub>2</sub>O for ligation.

The PCR-amplified DNA fragment containing the CON193 coding region was cloned into pCR2.1 using a protocol standard in the art. In particular, the ligation reaction consisted of 6  $\mu$ l of CON193 DNA, 1  $\mu$ l 10X ligation buffer, 2  $\mu$ l pCR2.1 (25 ng/ $\mu$ l, Invitrogen), and 1  $\mu$ l T4 DNA ligase (Invitrogen). The reaction  
25 mixture was incubated overnight at 14°C and the reaction was then stopped by heating at 65°C for 10 minutes. Two microliters of the ligation reaction were transformed into One Shot cells (Invitrogen) and plated onto ampicillin plates. A single colony containing an insert was used to inoculate a 5 ml culture of LB medium. The culture was grown for 18 hours and the plasmid DNA was purified using the Concert Rapid Plasmid Miniprep System (GibcoBRL) and sequenced. Following confirmation of the  
30 sequence, pCR-CON193 was identified, and a 50 ml culture of LB medium was

- 67 -

inoculated and recombinant plasmid DNA was purified using a Qiagen Plasmid Midi Kit to yield purified pCR-CON193.

## **B. Cloning of CON166 G Protein-Coupled Receptor**

### **5 B.1 Database Search Results**

The database searching identified clone 2553280H1 in the Incyte database as an interesting candidate sequence. The 2553280H1 clone was obtained and sequenced directly using an ABI377 fluorescence-based sequencer and the ABI PRISM™ Ready Dye-Deoxy Terminator kit with Taq FSTM polymerase as described above for CON 193 in Example 1A.1. From the sequence it was deduced that clone 2553280H1 contained 349 nucleotides of a GPCR coding region comprising a carboxy-terminal fragment of a putative GPCR corresponding to the sixth and seventh transmembrane regions (6TM and 7TM). In addition, clone 2553280H1 contained 1.2 kb of the 3' untranslated sequence of that GPCR. Referring to SEQ ID NO: 3, the nucleotide sequence of Clone 2553280H1 corresponds to nucleotides 663 to 1,014 of what was eventually determined to be the complete sequence of a novel seven-transmembrane receptor that was designated CON166. A database search with this partial sequence showed a 44% match to an activated T cell-specific G protein-coupled receptor.

### **20 B2. Screening of a Genomic Phage Library to Obtain a Full-Length GPCR Clone**

The PCR technique was used to prepare a genomic fragment for use as a probe specific for the genomic CON166 clone. Based on the complete sequence of clone 2553280H1, two oligonucleotide primers were designed: Primer LW1278: 5'-ACCGCTGCCTTTTCTAGTC-3' (SEQ ID NO: 28; see nucleotides 715 to 732 of SEQ ID NO: 3 and Primer LW1279: 5'-CCTTCTTTCTGGGTACATAAGTC-3' (SEQ ID NO: 29; see the reverse complement of nucleotides 951-973 of SEQ ID NO: 3). These primers were designed to amplify a 259 base-pair fragment of genomic DNA containing a portion of the CON166 coding region found in clone 2553280H1 (assuming the absence of introns in this region).

Initially, a suitable human genomic library constructed in EMBL SP6/T7 was amplified to provide the materials required for screening as described

- 68 -

above for CON193 in Example 1A.2. Polymerase chain reaction (PCR) was selected as a technique for screening the phage library. Each PCR reaction was done in a 20  $\mu$ l reaction volume containing 8.84  $\mu$ l H<sub>2</sub>O, 2  $\mu$ l 10X PCR buffer II (Perkin-Elmer), 2  $\mu$ l 25 mM MgCl<sub>2</sub>, 0.8  $\mu$ l dNTP mixture (dATP, dCTP, dGTP, dCTP, each at 10 mM), 0.12  $\mu$ l primer LW1278 (approximately 1  $\mu$ g/ $\mu$ l), 0.12  $\mu$ l primer LW1279 (approximately 1  $\mu$ g/ $\mu$ l), 0.12  $\mu$ l AmpliTaq Gold polymerase (5 Units/ $\mu$ l, with "Units" as defined by the supplier, Perkin-Elmer) and 2  $\mu$ l of phage from one of the 24 stock tubes. The PCR reaction involved 1 cycle at 95°C for 10 minutes and 80°C for 20 minutes, followed by 12 cycles at 95°C for 30 seconds, 72-61°C for 2 minutes (72°C for this stage of the second cycle, with a decrease of one degree for this stage in each succeeding cycle), 72°C for 30 seconds, followed by 30 cycles at 95°C for 15 seconds, 60°C for 30 seconds, and 72°C for 30 seconds.

Following PCR cycling, the contents from each reaction tube were loaded onto a 2% agarose gel and electrophoresed adjacent to known size standards to screen for PCR products of the expected size of 259 bp, indicative of a clone containing the portion of clone 2553280H1 amplified by the two selected primers. A positive signal (*i.e.*, a fragment of the expected size) was found in one of the 24 PCR reactions, thereby identifying a single stock genomic library tube containing positive clones.

From the original genomic library tube that had given a PCR product of the correct size, a 5  $\mu$ l phage aliquot was used to amplify the CON166 genomic phage DNA as described for CON 193 above in Example 1A.2. For the amplification of the phage library, the plates containing the diluted phage solution were subdivided into 12 sections unlike that of CON193; otherwise the procedures were identical.

The purified CON166 genomic phage DNA was sequenced using the ABI PRISM™ 310 Genetic Analyzer which uses advanced capillary electrophoresis technology and the ABI PRISM™ BigDye™ Terminator Cycle Sequencing Ready Reaction Kit as described above for CON193 in Example 1A.2. These efforts resulted in the determination of the CON166 polynucleotide sequence set forth in SEQ ID NO: 3 and the deduced amino acid sequence of the encoded CON166 polypeptide which is set forth in SEQ ID NO: 4.

### B.3 Subcloning of the Coding Region of CON166 via PCR

Additional experiments were conducted to subclone the coding region of CON166 from the genomic clone and place the isolated coding region into a useful vector. Two additional PCR primers were designed based on the coding region of CON166. The first PCR primer, designated Primer LW1405, has the sequence 5'-AAGCATAACATGGATGAAACAGGAAATCTG-3' (SEQ ID NO: 29, nucleotides 10-30 of which correspond to nucleotides 1-21 of SEQ ID NO: 3). To protect against exonucleolytic attack during subsequent exposure to enzymes, *e.g.*, Taq polymerase, primers were routinely synthesized with a protective run of nucleotides at the 5' end that were not necessarily complementary to the desired target. The second PCR primer is Primer LW1406, which has the sequence 5'-AAGCATAACTATACTTTACATATTTCTTC-3' (SEQ ID NO: 30, nucleotides 9-29 of which correspond to the reverse complement of nucleotides 994-1014 of SEQ ID NO: 3).

PCR was performed in a 50  $\mu$ l reaction containing 34  $\mu$ l H<sub>2</sub>O, 5  $\mu$ l 10X TT buffer (140 mM ammonium sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5  $\mu$ l 15 mM MgSO<sub>4</sub>, 2  $\mu$ l dNTP mixture (dGTP, dATP, dTTP, and dCTP, each at 10 mM), 3  $\mu$ l genomic phage DNA (0.25  $\mu$ g/ $\mu$ l), 0.3  $\mu$ l Primer LW1405 (1  $\mu$ g/ $\mu$ l), 0.3  $\mu$ l Primer LW1406 (1  $\mu$ g/ $\mu$ l), 0.4  $\mu$ l High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction was started with 1 cycle of 94°C for 2 minutes; followed by 25 cycles at 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1.3 minutes.

The contents from the PCR reaction were loaded onto a 2% agarose gel and fractionated. The DNA band of expected size (1,031 bp) was excised from the gel, placed in a GenElute Agarose spin column (Supelco) and spun for 10 minutes at maximum speed in a microfuge. The eluted DNA was precipitated with ethanol and resuspended in 6  $\mu$ l H<sub>2</sub>O for ligation.

The PCR-amplified DNA fragment containing the CON166 coding region was cloned into pCR2.1 to generate pCR-CON166 using a protocol standard in the art. In particular, the ligation reaction was carried out as described for CON193 in Example 1A.3. The resulting plasmid DNA was purified using the Concert Rapid

- 70 -

Plasmid Miniprep System (GibcoBRL) and sequenced. Following confirmation of the sequence, a 50 ml culture of LB medium was inoculated with the transformed One Shot cells, cultured, and processed using a Qiagen Plasmid Midi Kit to yield purified pCR-CON166.

5

## **C. Cloning of CON103 G Protein-Coupled Receptor**

### **C.1 Database Search Results**

The database searching identified clone 1581220H1 in the Incyte database as an interesting candidate sequence. The 1581220H1 clone was obtained and sequenced directly using an ABI377 fluorescence-based sequencer and the ABI PRISM™ Ready Dye-Deoxy Terminator kit with Taq FSTM polymerase as described for CON 193 in Example 1A.1. From the sequence it was deduced that clone 1581220H1 contained 454 nucleotides of a GPCR coding region comprising a carboxy-terminal fragment of a putative GPCR corresponding to the sixth and seventh transmembrane regions (6<sup>TM</sup> and 7<sup>TM</sup>). In addition, clone 1581220H1 contained 1.2 kb of the 3' untranslated sequence of that GPCR. Referring to SEQ ID NO: 5, the nucleotide sequence of clone 1581220H1 corresponds to nucleotides 698 to 1190 of what was eventually determined to be the complete sequence of a novel seven-transmembrane receptor designated CON103. A database search with this partial sequence showed a 44% match to an activated T cell-specific G protein-coupled receptor.

20

### **C.2 Screening of a Genomic Phage Library to Obtain a Full-Length GPCR Clone**

The PCR technique was used to prepare a genomic fragment for use as a probe specific for the genomic CON103 clone. Based on the complete sequence of clone 1581220H1, two oligonucleotide primers were designed: Primer LW1280: 5'-TCTGCACACAGCTCTTCCATGG-3' (SEQ ID NO: 32; see nucleotides 1568-1589 of SEQ ID NO: 5) and Primer LW1281: 5'-TCCCTTGTCAGTTGGTTGAGG-3' (SEQ ID NO: 33; see nucleotides 1926 to 1947 of SEQ ID NO: 5. These primers were designed to amplify a 380 base-pair fragment of genomic DNA containing a portion of the CON103 coding region found in clone 1581220H1 (assuming the absence of introns in this region).

25

30

- 71 -

Initially, a suitable human genomic library constructed in EMBL SP6/T7 was amplified to provide the materials required for screening as described above for CON193 in Example 1A.2. Polymerase chain reaction (PCR) was selected as a technique for screening the phage library. Each PCR reaction was done in a 20  $\mu$ l reaction volume containing 8.84  $\mu$ l H<sub>2</sub>O, 2  $\mu$ l 10X PCR buffer II (Perkin-Elmer), 2  $\mu$ l 25 mM MgCl<sub>2</sub>, 0.8  $\mu$ l dNTP mixture (dATP, dTTP, dGTP, dCTP, each at 10 mM), 0.12  $\mu$ l primer LW1280 (approximately 1  $\mu$ g/ $\mu$ l), 0.12  $\mu$ l primer LW1281 (approximately 1  $\mu$ g/ $\mu$ l), 0.12  $\mu$ l AmpliTaq Gold polymerase (5 Units/ $\mu$ l, with "Units" as defined by the supplier, Perkin-Elmer) and 2  $\mu$ l of phage from one of the 24 stock tubes. PCR amplification reactions using each one of the other 23 stock collections of genomic clones were performed under the same conditions. The PCR reaction involved 1 cycle at 95°C for 10 minutes and 80°C for 20 minutes, followed by 12 cycles at 95°C for 30 seconds, 72-61°C for 2 minutes (72°C for this stage of the second cycle, with a decrease of one degree for this stage in each succeeding cycle), 72°C for one minute, followed by 30 cycles at 95°C for 15 seconds, 60°C for 30 seconds, and 72°C for 30 seconds.

Following PCR cycling, the contents from each reaction tube were loaded onto a 2% agarose gel and electrophoresed adjacent to known size standards to screen for PCR products of the expected size of 380 bp, indicative of a clone containing the portion of clone 1581220H1 amplified by the two selected primers. A positive signal (*i.e.*, a fragment of the expected size) was found in one of the 24 PCR reactions, thereby identifying a single stock genomic library tube containing positive clones.

From the original genomic library tube that had given a PCR product of the correct size, a 5  $\mu$ l phage aliquot was used to amplify the CON 103 genomic phage DNA as described above for CON193 in Example 1A.2. A total of 8 plates were inoculated with eluted phage in this manner described above. Following incubation at 37°C for 16 hours, the top agarose from each of the 8 plates was removed to recover the phage, which were used to prepare purified genomic phage DNA using the Qiagen Lambda Midi Kit.

The CON103 clone was sequenced using the ABI PRISM™ 310 Genetic Analyzer. The cycle-sequencing reaction contained 6 µl of H<sub>2</sub>O, 8 µl of BigDye™ Terminator mix, 5 µl of miniprep clone DNA (0.1 µg/µl), and 1 µl primer (25 ng/µl). The reaction was performed in a Perkin-Elmer 9600 thermocycler at 25 cycles of 96°C for 10 seconds, 50°C for 10 seconds, and 60°C for 4 minutes. The product of the PCR reaction was purified using Centriflex™ gel filtration cartridges, dried under vacuum, and dissolved in 16 µl of Template Suppression Reagent (PE-Applied Biosystems). The samples were then incubated at 95°C for 5 minutes and placed in the 310 Genetic Analyzer. These efforts resulted in the determination of the CON103 polynucleotide sequence set forth in SEQ ID NO: 5 and the deduced amino acid sequence of the encoded CON103 polypeptide which is set forth in SEQ ID NO: 6.

### C.3 Subcloning of the Coding Region of CON103 via PCR

Additional experiments were conducted to subclone the coding region of CON103 from the genomic clone and place the isolated coding region into a useful vector. Two additional PCR primers were designed based on the sequence of the coding region of CON103: Primer LW1385 (5'-GCATAAGCT-TCCATGGAAGTTCATAACCTG-3'; SEQ ID NO: 34, nucleotides 13-30 of which correspond to nucleotides 1-18 of SEQ ID NO: 5) and Primer LW1386 (5'-GCATCTCGAGTTACCCCCACAGCGCTGCAG-3'; SEQ ID NO: 35, nucleotides 11-30 of which correspond to the reverse complement of nucleotides 1171-1190 of SEQ ID NO: 5). To protect against exonucleolytic attack during subsequent exposure to enzymes, e.g., Taq polymerase, primers were routinely synthesized with a protective run of nucleotides at the 5' end that were not necessarily complementary to the desired target.

PCR was performed in a 50 µl reaction containing 22.6 µl H<sub>2</sub>O, 5 µl 10X TT buffer (140 mM ammonium sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5 µl 15 mM MgSO<sub>4</sub>, 10 µl rapid dye (Origene), 2 µl dNTP mixture (dGTP, dATP, dTTP, and dCTP, each at 10 mM), 0.5 µl genomic phage DNA (0.97 µg/µl), 0.3 µl Primer LW1385 (1 µg/µl), 0.3 µl Primer LW1386 (1 µg/µl), and 0.4 µl High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction was started with 1 cycle

- 73 -

of 94°C for 2 minutes, followed by 12 cycles at 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1.3 minutes.

The contents from the PCR reaction were loaded onto a 2% agarose gel and fractionated. The DNA band of expected size (1,212 bp) was excised from the gel, placed in a GenElute Agarose spin column (Supelco) and spun for 10 minutes at maximum speed in a microcentrifuge. The eluted DNA was precipitated with ethanol and resuspended in 6 µl H<sub>2</sub>O for ligation.

The PCR-amplified DNA fragment containing the CON103 coding region was cloned into pCR2.1 using a protocol standard in the art. In particular, the ligation reaction was carried out as described above for CON193 in Example 1A.3. The resulting plasmid DNA was purified using the Concert Rapid Plasmid Miniprep System (GibcoBRL) and sequenced. Following confirmation of the sequence, pCR-CON103 was identified, and a 50 ml culture of LB medium was inoculated, cultured, and processed using a Qiagen Plasmid Midi Kit to yield purified pCR-CON103.

#### **D. Cloning of CON203 G Protein-Coupled Receptor**

##### **D.1 Database Search Results**

The database searching identified clone 3210396H1 in the Incyte database as an interesting candidate sequence. The 3210396H1 clone was obtained and sequenced directly using an ABI377 fluorescence-based sequencer and the ABI PRISM™ Ready Dye-Deoxy Terminator kit with Taq FSTM polymerase as described above for CON193 in Example 1A.1. From the sequence it was deduced that clone 3210396H1 contained all 1,002 nucleotides of a GPCR coding region (see SEQ ID NO: 7). A database search with this sequence showed a 33% match to a platelet activating receptor (Gene H963, GenBank Acc. No. AF002986).

##### **D.2 Subcloning of the Coding Region of CON203 via PCR**

Additional experiments were conducted to subclone the coding region of CON203 and place the isolated coding region into a useful vector. Two additional PCR primers were designed based on the sequence of the coding region of CON203: Primer LW1329: 5'-GCATCTCGAGTCAGCCTAAGGTTATGTTG-3' (SEQ ID NO: 36; see nucleotides 984 to 1,002 of SEQ ID NO: 7 for the reverse complement of

- 74 -

nucleotides 9-29 of SEQ ID NO: 36) and Primer LW1377: 5'-

GCATAAGCTTATGAACACCACAGTGATGC-3' (SEQ ID NO: 37; see

nucleotides 1-19 of SEQ ID NO: 7 which correspond to nucleotides 11-29 of SEQ ID NO: 37). To protect against exonucleolytic attack during subsequent exposure to

5 enzymes, *e.g.*, Taq polymerase, primers were routinely synthesized with a protective run of nucleotides at the 5' end that were not necessarily complementary to the desired target. These primers were designed to amplify a 1,020 base-pair fragment of clone 3210396H1 containing the complete coding region of CON203.

10 PCR was performed in a 50  $\mu$ l reaction containing 34  $\mu$ l H<sub>2</sub>O, 5  $\mu$ l 10X TT buffer (140 mM ammonium sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5  $\mu$ l 15 mM MgSO<sub>4</sub>, 2  $\mu$ l dNTP mixture (dGTP, dATP, dTTP, and dCTP, each at 10 mM), 3  $\mu$ l clone 3210396H1 (miniprep DNA), 0.3  $\mu$ l Primer LW1329 (1  $\mu$ g/ $\mu$ l), 0.3  $\mu$ l Primer LW1377 (1  $\mu$ g/ $\mu$ l), and 0.4  $\mu$ l High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction was started with 1 cycle of 94°C for 2 minutes, 15 followed by 12 cycles at 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1.3 minutes.

The contents from the PCR reaction were loaded onto a 1.2% agarose gel and fractionated. The DNA band of expected size (1,020 bp) was excised from the gel, placed in a GenElute Agarose spin column (Supelco) and spun for 10 minutes 20 at maximum speed in a microcentrifuge. The eluted DNA was precipitated with ethanol and resuspended in 6  $\mu$ l H<sub>2</sub>O for ligation.

The PCR-amplified DNA fragment containing the CON203 coding region was cloned into pCR2.1 using a standard protocol and the Original TA Cloning Kit (Invitrogen). Ligation reactions were carried out as described above for CON193 25 in Example 1A.3. The resulting plasmid DNA was purified using the Concert Rapid Plasmid Miniprep System (GibcoBRL) and sequenced. Following confirmation of the sequence, pCR-C203 was identified, and a 50 ml culture of LB medium was inoculated, cultured, and processed using a Qiagen Plasmid Midi Kit to yield purified pCR-C203.

30 The CON203 clone was sequenced using the ABI PRISM™ 310 Genetic Analyzer (P-E Applied Biosystems), which uses advanced capillary

- 75 -

electrophoresis technology and the ABI Prism™ BigDye™ Terminator Cycle Sequencing Ready Reaction Kit. The cycle-sequencing reaction contained 6 µl of H<sub>2</sub>O, 8 µl of BigDye™ Terminator mix, 5 µl of miniprep clone DNA (0.1 µg/µl), and 1 µl primer (25 ng/µl). The reaction was performed in a Perkin-Elmer 9600 thermocycler using the following conditions: 25 cycles of 96°C for 10 seconds, 50°C for 10 seconds, and 60°C for 4 minutes. The product of the PCR reaction was purified using Centriflex™ gel filtration cartridges, dried under vacuum, and dissolved in 16 µl of Template Suppression Reagent (PE-Applied Biosystems). The samples were then incubated at 95°C for 5 minutes and placed in the 310 Genetic Analyzer.

Initially, these efforts showed that the CON203 coding region cloned into pCR2.1 had a single bp difference from the corresponding sequence of clone 3210396H1. The single bp change in the pCR2.1 clone was eliminated by conforming that sequence to the sequence of clone 3210396H1 using the QuikChange Site-Directed Mutagenesis Kit (Stratagene). The method involves modification of a sequence during PCR amplification, for which PCR primers LW1387 (5'-GAGAAATATTTTCTAAAAAACCTGTTTTTGCAAAAACGG-3'; SEQ ID NO: 38) and LW1388 (5'-CCGTTTTTGCAAAAACAGGTTTTTTAGAAAA-ATATTTCTC-3'; SEQ ID NO: 39) were used. The PCR reaction contained 40 µl H<sub>2</sub>O, 5 µl 10X proprietary Reaction Buffer (Stratagene), 1 µl pCR-C203 (0.125 µg/µl) mini-prep DNA, 1 µl dNTP mixture (dGTP, dATP, dTTP, and dCTP, each at 10 mM), 1 µl Pfu DNA polymerase (2.5 Units/µl), 1 µl LW1387 (125 ng/µl) and 1 µl LW1388 (125 ng/l). The cycle conditions were 95°C for 30 seconds, followed by 12 cycles at 95°C for 30 seconds, 55°C for 1 minute, and 68°C for 12 minutes. The tube was then placed on ice for 2 minutes and 1 µl of *DpnI* was added. The tube was then incubated at 37°C for one hour. One microliter of the *DpnI*-treated DNA was transformed into Epicurian coli XL1-Blue supercompetent *E. coli* cells. Following isolation of pCR-C203, the entire insert was re-sequenced, thereby successfully verifying repair of the single-site polymorphism. As expected, the sequence of the CON203 coding region determined using this pCR2.1 clone is in complete agreement

- 76 -

with the CON203 coding region sequence of SEQ ID NO: 7 which specifies the amino acid sequence set forth in SEQ ID NO: 8.

## **E. Cloning of CON198 G Protein-Coupled Receptor**

### **5 E.1 Database Search Results**

The database searching identified Clone 3359808HI in the Incyte database as an interesting candidate sequence. The 3359808HI clone was obtained and sequenced using standard techniques. From the sequence it was deduced that Clone 3359808HI contained the entire coding region for a previously unidentified GPCR, which was designated "CON198." The DNA and deduced amino acid sequences for CON198 are set forth in SEQ ID NOS: 9 and 10, respectively. A database search with this CON198 DNA sequence showed a 61% match to the rat putative GPCR designated RA1c [Raming *et. al.*, *Recept Channels*, 6: 141-151 (1998)] and 46% identity to an olfactory receptor.

### **15 E.2 Subcloning of the Coding Region of CON198 via PCR**

Additional experiments were conducted to subclone the coding region of the CON198 clone into a useful vector. Two PCR primers were designed based on the coding region of CON198 for the purpose of PCR amplification of the CON198 coding sequence. The first, Primer LW1326, from 5' to 3' (SEQ ID NO: 42):  
20 GCATGAATTCATGATGGTGGATCCCAATGG, includes the 5' end of the CON198 coding sequence (underlined) as well as a *EcoRI* restriction site, useful for subsequent expression work. The second, Primer LW1327, from 5' to 3' (SEQ ID NO: 43): GCATCTCGAGCCTAGGGCTCTGAAGCG, includes sequence  
25 complementary to the 3' end of the CON198 coding sequence (underlined), preceded by a *XhoI* restriction site sequence useful for subsequent cloning and expression work.

The PCR was performed in a 50 µl reaction containing 34 µl H<sub>2</sub>O, 5 µl of 10X TT buffer (140 mM Ammonium Sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5 µl of 15 mM MgSO<sub>4</sub>, 2 µl of 10 mM dNTPs (dATP, dCTP, dTTP, dGTP), 2 µl of Clone 3359808HI mini-prep DNA (approx. 0.125 µg/µl), 0.3 µl of Primer  
30 LW1326 (1 µg/µl), 0.3 µl of Primer LW1327 (1 µg/µl), and 0.5 µl of High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction was started with 1 cycle

- 77 -

of 94°C for 2 minutes; followed by 12 cycles at 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1 minute.

5 The contents from the PCR reaction were loaded onto a 1.2% agarose gel and electrophoresed. The DNA band of expected size was excised from the gel, placed in a GenElute Agarose spin column (Supelco) and spun for 10 minutes at maximum speed in a microcentrifuge. The eluted DNA was ethanol-precipitated and resuspended in 6 µl H<sub>2</sub>O for ligation.

10 The purified PCR fragment containing the CON198 coding sequence was ligated into a commercial vector using Invitrogen's Original TA Cloning Kit. The ligation reaction was carried out as described above for CON193 in Example 1A.3. The resulting plasmid DNA was isolated using a Concert Rapid Plasmid Miniprep System (GibcoBRL) and sequenced to confirm that the plasmid contained the CON198 insert. Sequencing of the subcloned CON198 construct revealed that the PCR amplification had introduced a mutation (relative to the sequence of the original  
15 clone) at the nucleotide corresponding to position 204 of SEQ ID NO: 9. A site-directed mutagenesis experiment was performed using the QuikChange Site-Directed Mutagenesis Kit (Stratagene) to repair the mutation.

Two primers were designed to revert the mutated A nucleotide at position 204 back to a G nucleotide via polymerase chain reaction. Primer LW1415  
20 (SEQ ID NO: 44) contained the sequence:

5'-CCATGTATATATTTCTTTGCATGCTTTCAGGCATTGACATCC-3'; and primer LW1416 (SEQ ID NO: 45) contained the sequence:

5'-GGATGTCAATGCCTGAAAGCATGCAAAGAAATATATACATGG-3'. The PCR reaction contained 40 µl of H<sub>2</sub>O, 5 µl of 10x Reaction buffer, 1 µl of mini-prep  
25 DNA (approx. 0.125 µg/µl) from the CON198-pCR2.1 clone (as template), 1 µl of primer LW1415 (125 ng/µl), 1 µl of primer LW1416 (125 ng/µl), 1 µl of 10 mM dNTPs, 1 µl Pfu DNA polymerase. The PCR cycle conditions were as follows: initial denaturation at 95°C for 30 seconds, then 14 cycles at 95°C for 30 seconds, 55°C annealing for 1 minute, and 68°C extension for 12 minutes. Thereafter, the reaction  
30 tube was placed on ice for 2 minutes.

- 78 -

After PCR, 1  $\mu$ l of *DpnI* was added and the tube incubated at 37°C for one hour to digest the methylated parental DNA template. One microliter of the *DpnI*-treated DNA was transformed into Epicurian coli XL1-Blue supercompetent cells and the entire insert was re-sequenced. The resequencing confirmed that position 204 of SEQ ID NO: 9 had been successfully reverted to a guanine nucleotide.

Upon confirmation of the insert, the *E. coli* transformant was used to inoculate a 50 ml culture of LB medium. The culture was grown for 16 hours at 37°C, and centrifuged into a cell pellet. Plasmid DNA was purified from the pellet using a Qiagen Plasmid Midi Kit and again sequenced to confirm successful cloning of the CON198 insert, using an ABI377 fluorescence-based sequencer and the ABI PRISM™ Ready Dye-Deoxy Terminator kit with Taq FS™ polymerase as described above for CON 193 in Example 1A.1.

## **F. Cloning of CON197 G Protein-Coupled Receptor**

### **F.1 Database Search Results**

The database searching identified Clone 866390H1 in the Incyte database as an interesting candidate sequence. The 866390H1 clone was obtained and sequenced using standard techniques. From the sequence it was deduced that Clone 866390H1 contained the entire coding region for a previously unidentified GPCR, which was designated "CON197." The DNA and deduced amino acid sequences for CON197 are set forth in SEQ ID NOs: 11 and 12, respectively. A database search with this CON197 DNA sequence showed a 42% match to an olfactory receptor.

### **F.2 Subcloning of the Coding Region of CON197 via PCR**

Additional experiments were conducted to subclone the coding region of the CON197 clone into a useful vector. Two PCR primers were designed based on the coding region of CON197 for the purpose of PCR amplification of the CON197 coding sequence. The first, Primer LW1324, from 5' to 3' (SEQ ID NO: 48): GATCGGATCCATGGAAAGCGAGAACAG, includes the 5' end of the CON197 coding sequence (underlined) as well as a *Bam*HI restriction site, useful for subsequent expression work. The second, Primer LW1325, from 5' to 3' (SEQ ID NO: 49): GATCCTCGAGTCAGGCTATGTGCTTATTAAACACC, includes

- 79 -

sequence complementary to the 3' end of the CON197 coding sequence (underlined), preceded by a *Xho*I restriction site sequence useful for subsequent cloning and expression work.

The PCR was performed in a 50 µl reaction containing 24 µl H<sub>2</sub>O,  
5 10 µl Rapid Dye Loading buffer (Origene) 5 µl 10X TT buffer (140 mM Ammonium Sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5 µl of 15 mM MgSO<sub>4</sub>, 2 µl of 10 mM dNTPs (dATP, dCTP, dTTP, dGTP), 3 µl of Clone 866390H1 mini-prep DNA (approx. 0.125 µg/µl), 0.3 µl of Primer LW1324 (1 µg/µl), 0.3 µl of Primer LW1325 (1 µg/µl), and 0.5 µl of High Fidelity Taq polymerase (Boehringer Mannheim). The  
10 PCR reaction was started with 1 cycle of 94°C for 2 minutes; followed by 12 cycles at 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1 minute.

The contents from the PCR reaction was loaded onto a 1.2% agarose gel and electrophoresed. The DNA band of expected size was excised from the gel, placed in GenElute Agarose spin column (Supelco) and spun for 10 minutes at  
15 maximum speed in a Savant microcentrifuge. The eluted DNA was ethanol-precipitated and resuspended in 6 µl H<sub>2</sub>O for ligation.

The purified PCR fragment containing the CON197 coding sequence was ligated into a commercial vector using Invitrogen's Original TA Cloning Kit. The resulting plasmid DNA from the culture was isolated using a Concert Rapid Plasmid  
20 Miniprep System (GibcoBRL) and sequenced to confirm that the plasmid contained the CON197 insert.

Upon confirmation of the insert, the same transformant was used to inoculate a 50 ml culture of LB medium. The culture was grown for 16 hours at 37°C, and centrifuged into a cell pellet. Plasmid DNA was purified from the pellet  
25 using a Qiagen Plasmid Midi Kit and again sequenced to confirm successful cloning of the CON197 insert, using an ABI377 fluorescence-based sequencer (Perkin Elmer/Applied Biosystems Division, PE/ABD, Foster City, CA) and the ABI PRISM™ Ready Dye-Deoxy Terminator kit with Taq FS™ polymerase as described above for CON193 in Example 1A.1.

30

## **G. Cloning of CON202 G Protein-Coupled Receptor**

### **G.1 Database Search Results**

The database searching identified Clone Number 1305513H1 in the Incyte database as an interesting candidate sequence. The 1305513H1 clone was obtained and sequenced using an ABI377 fluorescence-based sequencer (Perkin Elmer/Applied Biosystems Division, PE/ABD, Foster City, CA) and the ABI PRISM™ Ready Dye-Deoxy Terminator kit with Taq FS™ polymerase as described above for CON193 in Example 1A.1.

Sequencing of Incyte Clone 1305513H1 revealed a sequence corresponding to nucleotides 1054 to 1378 of SEQ ID NO: 13. Using a FORTRAN computer program called "tmrest.all" [Parodi *et al.*, *Comput. Appl. Biosci.*, 5: 527-535 (1994)], Clone 1305513H1 was deduced to contain two transmembrane-spanning domains (TMVI and TMVII) and an extracellular loop for a previously unidentified GPCR, which was designated as "CON202". The sequence obtained was used as a tool to identify a full length GPCR clone as described in the next section.

### **G.2 PCR Screening of Genomic Clones**

A human genomic phage library was selected as a source from which to attempt to clone the CON202 gene. The genomic library was amplified as described above for CON193 in Example 1A.2.

This genomic library was screened by PCR using the primers: GV599 (5'GGCAGAAGAAGGCTATTGGTCTTAGACGAG3'; SEQ ID NO: 52), and GV600 (5'CTGAAACAGCGCCTCAGCTCCCC3'; SEQ ID NO: 53). These primers were designed from the sequence of Clone 1305513H1 to amplify a 253 base pair fragment (corresponding to nucleotides 1064 to 1317 of SEQ ID NO: 13) from any corresponding genomic clone in the library. The 20 µl PCR reactions each contained 12.8 µl of H<sub>2</sub>O, 2 µl of 10x PCR buffer II (Perkin-Elmer), 2 µl of 25 mM MgCl<sub>2</sub>, 0.8 µl of 10 mM dNTP's (dATP, dGTP, dCTP, dTTP), 0.12 µl of primer GV599 (1 µg/ml), 0.12 µl of primer GV600 (1 µg/ml), 0.2 µl AmpliTaq Gold polymerase (5 Units/µl, with "Units" as defined by the supplier, Perkin Elmer) and 2 µl of phage from one of the 24 tubes. The PCR reaction consisted of 1 cycle at 95°C for 10 minutes; then 17 cycles at 95°C for 20 seconds, 72°C for 2 minutes decreasing 1°C

- 81 -

each cycle, 72°C for 30 seconds followed by 30 cycles at 95°C for 20 seconds, 55°C for 30 seconds, and 72°C for 30 seconds.

The PCR products were visualized on a 2% agarose gel. For those tubes which produced the correct sized band of 253 bp, five microliters from each original phage culture tube were used to amplify the CON202 genomic phage DNA as described above for CON 193 in Example 1A.2.

The genomic DNA from the single phage isolate, was sequenced with the ABI PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary electrophoresis technology and the ABI PRISM™ Big Dye™ Terminator Cycle Sequencing Ready Reaction Kit. The cycle-sequencing reaction contained 20 ml of H<sub>2</sub>O, 16 ml of BigDye™ Terminator Mix, 1 ml of genomic phage DNA (1.1 mg/ml), and 3 ml primer (25 ng/ml). The reaction was performed in a Perkin-Elmer 9600 thermocycler at 95°C for 5 minutes, followed by 99 cycles of 95°C for 30 seconds, 55°C for 20 seconds and 60°C for 4 minutes. The product was purified using a Centriflex™ gel filtration cartridge, dried under a vacuum, then dissolved in 16 ml of Template Suppression Reagent. The samples were heated at 95°C for 5 minutes then placed in the 310 Genetic Analyzer.

### G.3 Subcloning of the Coding Region of CON202 via PCR

Additional experiments were conducted to subclone the coding region of the CON202 clone into a more useful vector. Two PCR primers were designed based on the coding region of CON202 for the purpose of PCR amplification of the CON202 coding sequence. The first, Primer LW1482 (5'AGCTATGGCGAACTATAGCCATGCAGC3'; SEQ ID NO: 54) included the 5' end of the CON202 coding sequence (underlined). The second, Primer LW148 (5'AGTCCTCATATAACACAGTAAGGTTCC3'; SEQ ID NO: 55) included the sequence complementary to the 3' end of the CON202 coding sequence (underlined).

The PCR was performed in a 50 µl reaction containing 36.5 µl of H<sub>2</sub>O, 5 µl of 10x TT buffer (140 mM Ammonium Sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5 µl of 15 mM MgSO<sub>4</sub>, 2 µl of 10 mM dNTP's (dATP, dCTP, dTTP, dGTP), 0.5 µl of CON202 genomic phage DNA (approx. 1.1 µg/µl), 0.3 µl of Primer LW1482 (1 µg/µl), 0.3 µl of Primer LW1483 (1 µg/µl), and 0.4 µl of High Fidelity Taq

- 82 -

polymerase (Boehringer Mannheim). The PCR reaction was started with 1 cycle of 94°C for 2 minutes; followed by 12 cycles at 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1.3 minutes.

5 The contents from the PCR reaction were loaded onto a 2.1% agarose gel and electrophoresed. The DNA band of expected size (1.1 kb) was excised from the gel, placed on a GenElute Agarose spin column (Supelco), and spun for 10 minutes at maximum speed in a microfuge. The eluted DNA was ethanol-precipitated and resuspended in 6 µl of H<sub>2</sub>O for ligation.

10 The purified PCR fragment, containing the CON202 coding sequence, was ligated into a commercial vector using Invitrogen's Original TA Cloning Kit. The ligation reaction was carried out as described above for CON193 in Example 1A.3. The resulting plasmid DNA from the culture was isolated using a Concert Rapid Plasmid Miniprep System (GibcoBRL) and sequenced to confirm that the plasmid contained the CON202 insert. The resulting construct was denoted as pCR-CON202.

15 The final subclone was sequenced using the ABI PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary electrophoresis technology and the ABI PRISM™ Terminator Cycle Sequencing Ready Reaction Kit. The cycle-sequencing reaction contained 6 ml of H<sub>2</sub>O, 8 ml of BigDye™ Terminator mix, 5 ml miniprep DNA (0.1 mg/ml), and 1 ml primer (25  
20 ng/ml). The reaction was performed in a Perkin-Elmer 9600 thermocycler at 25 cycles of 96°C for 10 seconds, 50°C for 10 seconds, and 60°C for 4 minutes. The product was purified using Centriflex™ gel filtration cartridges, dried under vacuum, then dissolved in 16 ml of Template Suppression Reagent. The samples were heated to 95°C for 5 minutes then placed in the 310 Genetic Analyzer.

25 Upon confirmation of the insert, the same transformant was used to inoculate a 50 ml culture of LB medium. The culture was grown for 16 hours at 37°C, and centrifuged into a cell pellet. Plasmid DNA was purified from the pellet using a Qiagen Plasmid Midi Kit and again sequenced to confirm successful cloning of the CON202 insert, as described above.

30

## H. Cloning of CON222 G Protein-Coupled Receptor

### H.1 Database Search Results

The database searching in the Incyte database identified Sequence Number 2488822CB1 as an interesting candidate sequence. This Incyte sequence is a consensus sequence derived by compiling multiple, shorter contiguous (apparently overlapping) partial sequences from cDNA clones. A single clone known to contain the complete consensus sequence was not available from Incyte. The following experiments were performed to clone a piece of human DNA which corresponds to the region of the theoretical Incyte Sequence Number 2488822CB that was deduced to encode a heretofore undescribed GPCR. The human DNA and protein that was eventually isolated is referred to herein as CON222.

### H.2 Isolation of CON222 Genomic DNA using PCR

To isolate a clone of CON222, PCR primers were designed based on the 5' and 3' ends of the open reading frame that was identified in the Incyte Sequence Number 2488822CB1. The first primer, designated as LW1440, has the sequence 5'AAGCGGATGTTTAGACCTCTTGTG3' (SEQ ID NO: 60) which corresponds to nucleotides 1 to 18 of SEQ ID NO: 15 (underlined). The second primer, designated LW1441, has the sequence 5'AACAGTCATGAATAGGAATTGAG3' (SEQ ID NO: 61) which is the reverse complement of nucleotides 1173 to 1191 of SEQ ID NO: 15 (underlined).

PCR was performed in a 50 ml reaction containing 22.1 ml H<sub>2</sub>O, 10 ml Rapid Dye Loading Buffer (Origene), 5 ml 10x TT buffer (140 mM Ammonium Sulfate, 0.1% gelatin, 0.6 M Tris-tricine pH 8.4), 5 ml 15 mM MgSO<sub>4</sub>, 2 ml 10 mM dNTP's (dATP, dCTP, dGTP, dTTP), 5 ml human genomic DNA (0.03 mg/ml) (Clontech, Cat# 6550-1), 0.3 ml of Primer LW1440 (1 mg/ml) (SEQ ID NO: 59), 0.3 ml of LW1441 (1 mg/ml) (SEQ ID NO: 60), 0.4 ml High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction was started with 1 cycle of 94°C for 2 minutes followed by 10 cycles at 94°C for 30 seconds, 55°C for 2 minutes, 72°C for 2 minutes then 25 cycles at 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 2 minutes. The PCR reaction was loaded onto a 1.2% agarose gel. The resulting band was not 1.2 kb in length as expected, indicating that this method was unsuccessful in

- 84 -

identifying an appropriate clone from the selected Clontech genomic DNA library containing the coding region of CON222.

A human genomic DNA phage library was selected as an alternate source from which to attempt to clone the CON222 gene. Internal primers were designed to attempt to isolate from a genomic library a single phage which expresses the complete coding region. The procedure was carried out as described above for CON193 in Example 1A.2.

PCR was performed to identify a phage that contained a genomic DNA insert which corresponds to the deduced complete coding region of Incyte Sequence Number 2488822CB1 using the primers: Primer LW1442: 5'GCCATTCTGTCCACAGAAG3' (SEQ ID NO: 58; see nucleotides 391 to 410 of SEQ ID NO: 15) and Primer LW1443: 5'TCAGTTGCTGTTATGGCAC3' (SEQ ID NO: 59; see reverse complement of nucleotides 744 to 761 of SEQ ID NO: 15). These primers were designed based on the deduced coding region of Incyte Sequence Number 2488822CB1, to amplify a 370 bp fragment (corresponding to nucleotides 391 to 761 of SEQ ID NO: 1) from any corresponding genomic clone in the library. The 50 µl PCR reactions each contained 32 µl of H<sub>2</sub>O, 5 µl of 10x PCR gold buffer (PE Applied Biosystems), 5 µl of 25 mM MgCl<sub>2</sub>, 2 µl of 10 mM dNTP's (dATP, dCTP, dGTP, dTTP), 0.3 µl of primer LW1442 (1µg/ml), 0.3 µl of primer LW1443 (1µg/ml), 0.4 µl AmpliTaq Gold polymerase (5 U/µl, with "Units" defined by the supplier; PE Applied Biosystems) and 5 µl of phage isolated human genomic DNA (0.03 µg/µl). The PCR reaction consisted of 1 cycle at 95°C for 10 minutes, then 17 cycles at 95°C for 20 seconds and 72°C for 2 minutes decreasing 1 degree each cycle, and 72°C for 1 minute, followed by 30 cycles at 95°C for 20 seconds, 55°C for 30 seconds, and 72°C for 1 minute. An aliquot of the PCR reaction was loaded onto a 1.2% agarose gel and electrophoresed. Although the internal primers were designed to produce a 370 bp PCR fragment, the resulting band was approximately 1.4 kb in length.

The DNA band was excised from the gel, placed on GenElute Agarose spin columns (Supelco) and spun for 10 minutes at maximum speed in a

- 85 -

microcentrifuge. The eluted DNA was ethanol-precipitated and resuspended in 10  $\mu$ l of H<sub>2</sub>O and 5  $\mu$ l was used to sequence the PCR band.

The PCR fragment was sequenced with an ABI PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary electrophoresis technology and the ABI PRISM™ BigDye™ Terminator Cycle Sequencing Ready Reaction Kit. Each cycle-sequencing reaction contained 6 ml of H<sub>2</sub>O, 8 ml of BigDye Terminator mix, 5 ml PCR fragment DNA (0.2 mg/ml), and 1 ml Primer LW1442 (25 ng/ml) and Primer LW1443 (25 ng/ml). The reaction was performed in a Perkin-Elmer 9600 thermocycler with 25 cycles of 96°C for 10 seconds, 50°C for 10 seconds, and 60°C for 4 minutes. The product was purified using Centriflex™ gel Reagent (PE Applied Biosystems). The samples were heated at 95°C for 5 minutes then placed in the 310 Genetic Analyzer.

The sequence analysis determined that there is an intron in the middle of the 5th transmembrane-spanning domain between nucleotides 673 and 674 in SEQ ID NO: 15. This intron was responsible for the unexpectedly large PCR fragment.

### H.3 Isolation of Full Length cDNA

Since attempts to isolate an uninterrupted coding region from genomic DNA were unsuccessful, a fetal brain cDNA was used to generate the complete coding region of Incyte Sequence Number 2488833CB1. The PCR primers described above, LW1440 (SEQ ID NO: 60) and LW1441 (SEQ ID NO: 61), which correspond to the 5' and 3' end of CON222 respectively, were used to generate the full length coding region.

The 50  $\mu$ l PCR reaction contained 37.4  $\mu$ l of H<sub>2</sub>O, 5  $\mu$ l of 10x cDNA PCR buffer (Clontech), 1  $\mu$ l of 10 mM dNTP's (dATP, dCTP, dTTP, dGTP), 5  $\mu$ l of Marathon-Ready Fetal Brain cDNA (Clontech), 0.3  $\mu$ l of Primer LW1440 (1  $\mu$ g/ $\mu$ l), 0.3  $\mu$ l of Primer LW1441 (1  $\mu$ g/ $\mu$ l), and 1  $\mu$ l of 50x Advantage cDNA polymerase (Clontech). The PCR reaction was started with 1 cycle of 94°C for 1 minute, followed by 30 cycles at 94°C for 30 seconds, 50°C for 30 seconds, and 68°C for 3 minutes.

The contents from the PCR reaction were loaded onto a 1.2% agarose gel and electrophoresed. The DNA band of expected size (1.2 kb) was excised from

- 86 -

the gel, placed on a GenElute Agarose spin column (Supelco), and spun for 10 minutes at maximum speed in a microfuge. The eluted DNA was ethanol-precipitated and resuspended in 6  $\mu$ l H<sub>2</sub>O for ligation.

#### H.4 Subcloning of Coding Region of CON222 via PCR

5                   After a cDNA containing the full length CON222 open reading frame was obtained, the coding region of CON222 was then subcloned into a more useful vector as follows.

                  The purified PCR fragment described above, containing the CON222 coding sequence, was ligated into a commercial vector using Invitrogen's Original TA  
10                  Cloning Kit. The ligation reaction was carried out as described above for CON193 in Example 1A.3. The resulting plasmid DNA from the culture was isolated using a Concert Rapid Plasmid Miniprep System (GibcoBRL) and sequenced to confirm that the plasmid contained the CON222 insert.

                  The subcloned insert in pCR2.1 was sequenced using the ABI  
15                  PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary technology and the ABI PRISM™ BigDye™ Terminator Cycle Sequencing Ready Reaction Kit. Each cycle-sequence reaction contained 6 ml of H<sub>2</sub>O, 8 ml of BigDye™ Terminator mix, 5 ml mini-prep DNA (0.1 mg/ml), and 1 ml of primer (25 ng/ml) and was performed in a Perkin-Elmer 9600 thermocycler with 25 cycles of  
20                  96°C for 10 seconds, 50°C for 10 seconds, and 60°C for 4 minutes. The product was purified using a Centriflex™ gel filtration cartridge, vacuum dried and dissolved in 16 ml of Template Suppression Reagent (PE Applied Biosystems). The samples were heated at 95°C for 5 minutes then placed in the 310 Genetic Analyzer.

                  Upon confirmation of the insert, the same transformant was used to  
25                  inoculate a 50 ml culture of LB medium. The culture was grown for 16 hours at 37°C, and centrifuged into a cell pellet. Plasmid DNA was purified from the pellet using a Qiagen Plasmid Midi Kit and again sequenced to confirm successful cloning of the CON222 insert, as described above.

30

- 87 -

**I. Cloning of CON215 G Protein-Coupled Receptor****I.1 Database Search Results**

The database searching identified Clone 1452259H1 in the Incyte database as an interesting candidate sequence. The sequence from 1452259H1 clone was used to search the Incyte full-length database and matched the entry 1650519CB1. An inspection of the clones that made up 1650519CB1 indicated that Incyte Clone 2796157H1 probably contained the full-length coding region. Sequence analysis of Incyte Clone 2796157H1 indicated that it contains the entire coding region for a previously unidentified GPCR, which was designated "CON215", along with 12 nucleotides of 5' untranslated region, 63 nucleotides of 3' untranslated region and a poly A<sup>+</sup> tail. The DNA and deduced amino acid sequences for CON215 are set forth in SEQ ID NOS: 17 and 18, respectively. A database search with this CON215 sequence showed a 47% match to the human probable G protein-coupled receptor KIA0001.

Since the untranslated regions were relatively short, it was not necessary to remove the coding region of CON215 from the pINCY vector (Incyte) and the construct is referred to as pINCY-CON215. The Incyte Clone 2796157H1 was sequenced using the ABI PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary electrophoresis technology and the ABI PRISM™ BigDye™ Terminator Cycle Sequencing Ready Reaction Kit as described above for CON222 in Example 1H.4.

**J. Cloning of CON217 G Protein-Coupled Receptor****J.1 Database Search Results**

The Incyte database search identified EST 3700658H1 as an interesting candidate sequence. The EST sequence No. 3700658H1 was used to search the Incyte full length database. This search identified Incyte clone No. 3356166H1 as a clone that potentially contained a full length GPCR corresponding to the selected EST.

The 3356166H1 clone was obtained from Incyte and sequenced using an ABI377 fluorescence-based sequencer ( and the ABI PRISM™ Ready Dye-Deoxy

- 88 -

Terminator kit with Taq FS™ polymerase as described above for CON193 in Example 1A.1.

Sequencing of Incyte Clone No. 3356166H1 revealed a 2480 basepair sequence as shown in SEQ NO: 19. Using a FORTRAN computer program called "tmrest.all" [Parodi *et al.*, *Comput. Appl. Biosci.*, 5: 527-535 (1994)], Clone No. 3356166H1 was deduced to contain seven transmembrane-spanning domains (TMI-TMVII) and was designated as "CON217" (SEQ ID NO: 20). The following experiments were performed to subclone and isolate the full length coding sequence of CON217 from Incyte Clone No. 3356166H1.

## J.2 Subcloning of the Coding Region of GPCR217

To subclone the full length coding sequence of CON217, PCR primers were designed based on the 5' and 3' ends of the open reading frame that was identified in the Incyte Clone No. 3356166H1. The first primer, designated as LW1448, has the sequence 5'AAGCGGTACCATGTTAGCCAACAGCTCCTC3' (SEQ ID NO: 66) which corresponds to nucleotides 42 to 62 of SEQ ID NO: 19 (underlined). The second primer, designated LW1449, has the sequence 5'AAGCTCTAGATCAGAGGGCGGAATCCTGG3' (SEQ ID NO: 67) which is the reverse complement of nucleotides 1142 to 1160 of SEQ ID NO: 20 (underlined). The primers also include recognition sequences (bold) for the restriction enzymes KpnI and XbaI, respectively.

PCR was performed in a 50 ml reaction containing 32.5 ml of H<sub>2</sub>O, 5 ml of 10x Pfx Amplification buffer (GibcoBRL), 5 ml of 10x PCR Enhancer solution (GibcoBRL), 1.5 ml of 50 mM MgSO<sub>4</sub>, 2 ml of 10 mM dNTP's (dATP, dCTP, dGTP, dTTP), 3 ml 3356166H1 mini-prep DNA (0.125 mg/ml obtained with the Concert Rapid Plasmid Miniprep System; GibcoBRL), 0.3 ml of Primer LW1448 (1 mg/ml) (SEQ ID NO: 3), 0.3 ml of Primer LW1449 (1 mg/ml) (SEQ ID NO: 4), 0.5 ml Platinum Pfx DNA polymerase (2.5 U/ml; GibcoBRL). The PCR reaction was started with 1 cycle of 94°C for 2 minutes followed by 25 cycles at 94°C for 30 seconds, 55°C for 30 seconds, 68°C for 1.3 minutes.

The contents from the PCR reaction were loaded onto a 1.2% agarose gel and electrophoresed. The DNA band of expected size (~1.1 kb) was excised from

- 89 -

the gel, placed on a GenElute Agarose spin column (Supelco), and spun for 10 minutes at maximum speed in a microfuge. The eluted DNA was ethanol-precipitated and resuspended in 6  $\mu$ l of H<sub>2</sub>O for ligation.

5 The purified PCR fragment, containing the CON217 coding sequence, was ligated into a commercial vector designated pCR2.1 using Invitrogen's Original TA Cloning Kit. The ligation reaction was carried out as described above for CON193 in Example 1A.3. The resulting plasmid DNA from the culture was isolated using a Concert Rapid Plasmid Miniprep System (GibcoBRL) and sequenced to confirm that the plasmid contained the CON217 insert and to confirm that no errors were introduced during PCR amplification. The resulting construct was denoted as pCR-CON217.

10 The final subclone was sequenced using the ABI PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary electrophoresis technology and the ABI PRISM™ Terminator Cycle Sequencing Ready Reaction Kit as described above for CON222 in Example 1H.4.

## EXAMPLE 2

### Analysis of G Protein-Coupled Receptor Sequence

#### A. CON193

20 The DNA and deduced amino acid sequence for CON193 are set forth in SEQ ID NOS: 1 and 2, respectively. Beginning with the initiation codon (methionine), the CON193 genomic Clone contains an open reading frame of 963 nucleotides encoding 321 amino acids, followed by a stop codon. Using a FORTRAN computer program called "tmrest.all" [Parodi *et al.*, *Comput. Appl. Biosci.*, 5: 527-535 (1994)], CON193 was shown to contain seven transmembrane-spanning domains corresponding to residues 30-49 (1TM), 61-81 (2TM), 103-122 (3TM), 146-165 (4TM), 199-222 (5TM), 243-262 (6TM), and 270-295 (7TM) of SEQ ID NO: 2. These transmembrane domains define first ("N-terminal," residues 1-29), second ("first EC loop," residues 82-102), third ("second EC loop," residues 166-198), and fourth ("third EC loop," residues 263-269) extracellular domains, as well as first ("first IC loop," residues 50-60), second ("second IC loop," residues 123-145), third

25

30

- 90 -

("third IC loop," residues 223-242), and fourth ("C-terminal," residues 296-321) intracellular domains.

Inspection of the CON193 amino acid sequence (SEQ ID NO: 2) reveals that this GPCR contains a DRY sequence following the third transmembrane domain (3TM) and a PIVY sequence found in the sixth transmembrane domain (TM6). In addition, the CON193 polynucleotide sequence was compared to sequences of known genes. CON193 is 45% identical and 72% similar to the mouse olfactory receptor gene S19 [see Malnic et al., Cell 96:713-723 (1999)]. This level of sequence similarity suggests that CON193 is a novel GPCR.

The CON193 cDNA clone (SEQ ID NO:1) was deposited with the National Center for Agricultural Utilization Research at the United States Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 in accordance with the Budapest Treaty on January 18, 2000. The clone was given accession no. B-30250.

#### B. CON166

The DNA and deduced amino acid sequence for CON166 are set forth in SEQ ID NOS: 3 and 4, respectively. Beginning with the initiation codon (methionine), the CON166 genomic clone contains an open reading frame of 1,011 nucleotides encoding 337 amino acids, followed by a stop codon. Using a FORTRAN computer program called "tmrest.all" [Parodi *et al.*, *Comput. Appl. Biosci.*, 5: 527-535 (1994)], CON166 was shown to contain seven transmembrane-spanning domains corresponding to the following residues presented in SEQ ID NO: 4: 1TM (30-49), 2TM (59-79), 3TM (99-119), 4TM (141-161), 5TM (191-215), 6TM (231-251), and 7TM (277-296). These transmembrane domains define first ("N-terminal," residues 1-29), second ("first EC loop," residues 80-98), third ("second EC loop," residues 162-190), and fourth ("third EC loop," residues 252-276), extracellular domains as well as first ("first IC loop," residues 50-58), second ("second IC loop," residues 120-140), third ("third IC loop," residues 216-230), and fourth ("C-terminal," residues 297-337) intracellular domains.

- 91 -

Inspection of the CON166 amino acid sequence (SEQ ID NO:2) reveals that this GPCR contains an FRC sequence following the third transmembrane domain (3TM), which is typically occupied by a consensus DRY sequence in other GPCRs; a PLLY sequence is also found in the seventh transmembrane domain (7TM). In addition, the CON166 polynucleotide sequence was compared to sequences of known genes. CON166 is 44% identical and 62% similar to a T-cell-specific G protein-coupled receptor of *Gallus gallus* found in the TREMBL database (Accession No. L06109). This level of sequence similarity suggests that CON166 is a novel GPCR.

The CON166 cDNA clone (SEQ ID NO:3) was deposited with the National Center for Agricultural Utilization Research at the United States Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 in accordance with the Budapest Treaty on January 18, 2000. The clone was given accession no. B-30248.

#### C. CON103

The DNA and deduced amino acid sequence for CON103 are set forth in SEQ ID NOS: 5 and 6, respectively. Beginning with the initiation codon (methionine), the CON103 genomic clone contains an open reading frame of 1,152 nucleotides encoding 384 amino acids, followed by a stop codon and a short open reading frame (SEQ ID NO: 5). Using a FORTRAN computer program called "tmrest.all" [Parodi *et al.*, *Comput. Appl. Biosci.*, 5: 527-535 (1994)], CON103 was shown to contain seven transmembrane-spanning domains corresponding to the following residues in SEQ ID NO: 6: 54-77 (1TM), 89-108 (2TM), 134-149 (3TM), 167-188 (4TM), 216-240 (5TM), 258-283 (6TM), and 301-320 (7TM). These transmembrane domains define first ("N-terminal," residues 1-53), second ("first EC loop," residues 109-133), third ("second EC loop," residues 189-215), and fourth ("third EC loop," residues 284-300) extracellular domains, as well as first ("first IC loop," residues 78-88), second ("second IC loop," residues 150-166), third ("third IC loop," residues 241-257), and fourth ("C-terminal," residues 321-384) intracellular domains.

- 92 -

Inspection of the CON103 amino acid sequence (SEQ ID NO: 6) reveals that this GPCR contains an NRY sequence following the third transmembrane domain (3TM), which is typically occupied by a consensus DRY sequence in other GPCRs. In addition, the CON103 polynucleotide sequence was compared to sequences of known genes. CON103 is 36% identical to GPR31 (GenBank Accession No. U65402) and 31% identical to the P2Y1 purinergic receptor (GenBank Accession No. S81950). This level of sequence similarity indicates that CON103 is a novel GPCR.

The CON103 cDNA clone (SEQ ID NO:5) was deposited with the National Center for Agricultural Utilization Research at the United States Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 in accordance with the Budapest Treaty on January 18, 2000. The clone was given accession no. B-30247.

#### D. CON203

The DNA and deduced amino acid sequence for CON203 are set forth in SEQ ID NOS: 7 and 8, respectively. Beginning with the initiation codon (methionine), the CON203 genomic clone contains an open reading frame of 999 nucleotides encoding 333 amino acids, followed by a stop codon. Using a FORTRAN computer program called "tmrest.all" [Parodi *et al.*, *Comput. Appl. Biosci.*, 5: 527-535 (1994)], CON203 was shown to contain seven transmembrane-spanning domains corresponding to the following residues of SEQ ID NO: 7: nucleotides 29-53 (1TM), 63-82 (2TM), 97-118 (3TM), 136-160 (4TM), 189-211 (5TM), 232-252 (6TM), and 281-300 (7TM). These transmembrane domains define first ("N-terminal," residues 1-28), second ("first EC loop," residues 83-96), third ("second EC loop," residues 161-188), and fourth ("third EC loop," residues 253-280) extracellular domains, as well as first ("first IC loop," residues 54-62), second ("second IC loop," residues 119-135), third ("third IC loop," residues 212-231), and fourth ("C-terminal," residues 301-333) intracellular domains.

Inspection of the CON203 amino acid sequence (SEQ ID NO: 8) reveals that this GPCR contains a DRF sequence following the third transmembrane

- 93 -

domain (3TM), which is typically occupied by a consensus DRY sequence in other GPCRs; CON203 also exhibited a PLIY sequence in the seventh transmembrane domain (7TM). In addition, the CON203 polynucleotide sequence was compared to sequences of known genes. CON203 is 33% identical to a platelet activating receptor (GenBank Accession No. AF002986. This level of sequence similarity suggests that CON203 is a novel GPCR.

The CON203 cDNA clone (SEQ ID NO: 7) was deposited with the National Center for Agricultural Utilization Research at the United States Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 in accordance with the Budapest Treaty on January 18, 2000. The clone was given accession no. B-30254.

#### E. CON198

The DNA and deduced amino acid sequence for CON198 are set forth in SEQ ID NO: 9 and 10 respectively. Beginning with the initiator methionine, the CON198 genomic clone contains an open reading frame of 954 nucleotides encoding 318 amino acids, followed by a stop codon. It will be appreciated that residue 2 of SEQ ID NO: 10 also is a methionine. Amino-terminal sequencing of purified native or recombinant CON198 protein will provide an indication as to which methionine acts as an initiator codon *in vivo*. Using a FORTRAN computer program called "tmrest.all" [Parodi *et al.*, *Comput. Appl. Biosci.*, 5: 527-535 (1994)], CON198 was deduced to contain seven transmembrane-spanning domains corresponding to residues 28-52 (TM1), 61-80 (TM2), 104-123 (TM3), 147-167 (TM4), 200-226 (TM5), 239-263 (TM6), and 274-295 (TM7) of SEQ ID NO: 10. These transmembrane domains define first ("N-terminal," residues 1-27 or 2-27), second ("first EC loop," residues 81-103), third ("second EC loop," residues 168-199), and fourth ("third EC loop," residues 264-273) extracellular domains as well as first ("first IC loop," residues 53-60), second ("second IC loop," residues 124-146), third ("third IC loop," residues 227-238), and fourth ("C-terminal," residues 296-318) intracellular domains.

CON198 contains a DRY sequence following the third transmembrane domain (TM3), a feature that is conserved in most GPCR. The most similar sequence

- 94 -

in a public database, at the time of initial screening, was that of rat GPCR RA1c, which shared only 61% identity at the amino acid level.

The CON198 cDNA clone (SEQ ID NO: 9) was deposited with the National Center for Agricultural Utilization Research at the United States Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 in accordance with the Budapest Treaty on January 18, 2000. The clone was given accession no. B-30252.

#### F. CON197

The DNA and deduced amino acid sequence for CON197 are set forth in SEQ ID NO: 11 and 12, respectively. Beginning with the initiator methionine, the CON197 genomic clone contains an open reading frame of 921 nucleotides encoding 307 amino acids, followed by a stop codon. Using a FORTRAN computer program called "tmrest.all" [Parodi *et al.*, *Comput. Appl. Biosci.*, 5: 527-535 (1994)], CON197 was deduced to contain seven transmembrane-spanning domains corresponding to residues 23-47 (TM1), 58-78 (TM2), 99-120 (TM3), 142-164 (TM4), 195-219 (TM5), 237-258 (TM6), and 270-289 (TM7) of SEQ ID NO: 12. These transmembrane domains define first ("N-terminal" residues 1-22), second ("first EC loop" residues 79-98), third ("second EC loop" residues 165-194), and fourth ("third EC loop" residues 259-269) extracellular domains as well as first ("first IC loop" residues 48-57), second ("second IC loop" residues 121-141), third ("third IC loop" residues 220-236), and fourth ("C-terminal" residues 290-309) intracellular domains.

CON197 contains a DRY sequence following the third transmembrane domain (TM3), a feature that is conserved in most GPCR. The most similar sequence in a public database, at the time of initial screening, was that of an olfactory receptor, which shared only 42% identity at the amino acid level.

The CON197 cDNA clone (SEQ ID NO: 11) was deposited with the National Center for Agricultural Utilization Research at the United States Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 in accordance with the Budapest Treaty on January 18, 2000. The clone was given accession no. B-30251.

**G. CON202**

The DNA and deduced amino acid sequence for this phage insert, termed "CON202", are set forth in SEQ ID NO: 13 and 14, respectively. The CON202 open reading frame, as depicted in SEQ ID NO: 14, begins with the initiator methionine and spans 1110 nucleotides which encode 370 amino acids, followed by a stop codon. Since this gene was isolated from genomic DNA and there are no apparent interruptions in the sequence, it is likely that CON202 contains no introns within the coding region. The full length clone of CON202 contained seven transmembrane-spanning domains corresponding to residues, 24 to 46 (TM1), 57 to 77 (TM2), 96 to 117 (TM3), 135 to 159, (TM4) TMV comprises 184 to 202 (TM5), 286 to 308 (TM6), 316 to 339 (TM7) of SEQ ID NO: 14. TM2 terminates with PFVC instead of the characteristic PXXY. TM3 is followed by the sequence TRY instead of the characteristic DRY. These transmembrane domains define first ("N-terminal," residues 1-23), second ("first EC loop," residues 78-95), third ("second EC loop," residues 160-183), and fourth ("third EC loop," residues 309-315) extracellular domains as well as first ("first IC loop," residues 47-56), second ("second IC loop," residues 118-134), third ("third IC loop," residues 203-285), and fourth ("C-terminal," residues 340-370) intracellular domains.

The CON202 cDNA clone (SEQ ID NO: 13) was deposited with the National Center for Agricultural Utilization Research at the United States Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 in accordance with the Budapest Treaty on January 18, 2000. The clone was given accession no. B-30253.

**H. CON222**

The sequence of CON222 coding region deduced the DNA and amino acid sequence set forth in SEQ ID NO: 15 and 16, respectively. The open reading frame that is depicted in SEQ ID NO: 16 begins with an initiator codon and spans 1188 nucleotides which encode 396 amino acids, followed by a stop codon.

The full length clone of CON222 contains seven transmembrane-spanning domains corresponding to residues 42-65 (TM1) 79-103, (TM2), 125-156,

- 96 -

(TM3), 167-188 (TM4), 217-241 (TM5), 268-290 (TM6), 301-320 (TM7) of SEQ ID NO: 16. TM2 is followed by a FRC sequence and TM7 contains a PILY sequence within. These transmembrane domains define first ("N-terminal," residues 1-41), second ("first EC loop," residues 104-124), third ("second EC loop," residues 189-216), and fourth ("third EC loop," residues 291-300) extracellular domains as well as first ("first IC loop," residues 66-78), second ("second IC loop," residues 157-166), third ("third IC loop," residues 242-267), and fourth ("C-terminal," residues 321-396) intracellular domains. A search of the public database indicated that CON222 is about 35% identical to a unique GPCR found in the nervous system of *Lymnaea stagnalis*.

The CON222 cDNA clone (SEQ ID NO: 15) was deposited with the National Center for Agricultural Utilization Research at the United States Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 in accordance with the Budapest Treaty on January 18, 2000. The clone was given accession no. B-30257.

#### I. CON215

The DNA and deduced amino acid sequence for CON215 are set forth in SEQ ID NO: 17 and 18, respectively. Beginning with the initiator methionine, the CON215 genomic clone contains an open reading frame of 1074 nucleotides encoding 358 amino acids, followed by a stop codon. Using a FORTRAN computer program called "tmrest.all" [Parodi *et al.*, *Comput. Appl. Biosci.*, 5: 527-535 (1994)], CON215 was deduced to contain seven transmembrane-spanning domains corresponding to residues 42-66 (TM1), 81-99 (TM2), 116-137 (TM3), 156-180 (TM4), 210-234 (TM5), 256-275 (TM6), and 308-328 (TM7) of SEQ ID NO: 18. These transmembrane domains define first ("N-terminal," residues 1-41), second ("first EC loop," residues 100-115), third ("second EC loop," residues 181-209), and fourth ("third EC loop," residues 276-307) extracellular domains as well as first ("first IC loop," residues 67-80), second ("second IC loop," residues 138-155), third ("third IC loop," residues 235-255), and fourth ("C-terminal," residues 329-358) intracellular domains.

- 97 -

CON215 contains a DRY sequence following the third transmembrane domain (TM3), a feature that is conserved in most GPCR. CON215 also contains a PLIY sequence within the seventh transmembrane domain (TM7).

The CON215 cDNA clone (SEQ ID NO: 17) was deposited with the National Center for Agricultural Utilization Research at the United States Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 in accordance with the Budapest Treaty on January 18, 2000. The clone was given accession no. B-30255.

#### 10 J. CON217

The DNA and deduced amino acid sequences of CON217 are set forth as SEQ ID NO: 19 and 20, respectively. The open reading frame that is depicted in SEQ ID NO: 2 begins with an initiator methionine codon and spans 1116 nucleotides which encode 372 amino acids, followed by a stop codon. In addition, the nucleotide sequence consists of 41 bp in the 5' untranslated region and 1323 bp in the 3' untranslated region.

The full length clone of CON217 contains seven transmembrane-spanning domains as indicated by the FORTRAN computer program "tmrest.all" [Parodi *et al.*, *Comput. Appl. Biosci.*, 5: 527-535 (1994)] which corresponds to 29-50 (TM1), 57-75 (TM2), 96-117 (TM3), 137-160 (TM4), 188-210 (TM5), 235-258 (TM6), 277-297 (TM7). TM3 is followed by a DRY sequence and TM7 contains a PLVY sequence within. These transmembrane domains define first ("N-terminal," residues 1-28), second ("first EC loop," residues 76-95), third ("second EC loop," residues 161-187), and fourth ("third EC loop," residues 259-276) extracellular domains as well as first ("first IC loop," residues 51-56), second ("second IC loop," residues 118-136), third ("third IC loop," residues 211-234), and fourth ("C-terminal," residues 298-372) intracellular domains. A search of the public database indicated that CON217 is about 41% identical to GPR23 (Genebank Accession No.: U66578) and to a purinergic receptor P2Y9 (Genebank Accession No.: U90322).

The CON215 cDNA clone (SEQ ID NO: 19) was deposited with the National Center for Agricultural Utilization Research at the United States Department

- 98 -

of Agriculture 1815 North University Street, Peoria, Illinois 61604 in accordance with the Budapest Treaty on January 18, 2000. The clone was given accession no. B-30256.

#### 5 K. Summary of Deposits

The polynucleotides (SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15 and 17) encoding the GPCR polypeptides of the invention were deposited with the Agricultural Research Service Culture Collection (NRRL) at the National Center Agricultural Utilization Research at the U.S. Department of the Agriculture 1815 North University Street, Peoria, Illinois 61604. These deposits were made in accordance with the Budapest Treaty on the International Recognition of the Deposit of Microorganism for the Purposes of Patent Procedures. The table below lists the details of these deposits.

	<u>GPCR</u>	<u>SEQ ID NO:</u>	<u>NRRL No.</u>	<u>Deposit Date</u>
15	CON193	1	B-30250	1/18/00
	CON166	3	B-30248	1/18/00
	CON103	5	B-30247	1/18/00
	CON203	7	B-30254	1/18/00
	CON198	9	B-30252	1/18/00
20	CON197	11	B-30251	1/18/00
	CON202	13	B-30253	1/18/00
	CON222	15	B-30257	1/18/00
	CON215	17	B-30255	1/18/00
	CON217	19	B-30256	1/18/00

25

#### EXAMPLE 3

##### Hybridization Analysis Demonstrates that the GPCRs are Expressed in the Brain

The expression of GPCR polynucleotides in mammals, such as the rat, was investigated by *in situ* hybridization histochemistry. Coronal and sagittal rat

30

- 99 -

brain cryosections (20  $\mu$ m thick) were prepared using a Reichert-Jung cryostat. Individual sections were thaw-mounted onto silanized, nuclease-free slides (CEL Associates, Inc., Houston, TX), and stored at -80°C. Sections were processed starting with post-fixation in cold 4% paraformaldehyde, rinsed in cold phosphate-buffered saline (PBS), acetylated using acetic anhydride in triethanolamine buffer, and dehydrated through a series of alcohol washes in 70%, 95%, and 100% alcohol at room temperature. Subsequently, sections were delipidated in chloroform, followed by rehydration through successive exposure to 100% and 95% alcohol at room temperature. Microscope slides containing processed cryosections were allowed to air dry prior to hybridization.

#### A. CON193

A CON193-specific probe was generated using PCR. The probe consisted of a 270 bp fragment containing sequence at the 3' end of CON-193. The primers for PCR amplification were LW 1248 [5'-GCATGAATTCCAATATACTTCCCCATACCTAC-3'; SEQ ID NO: 26) and LW 1249 [5'-GCATGGATCCGGAAGAAGGAGAAGAAAG-3'; SEQ ID NO: 27), which introduced terminal *Eco*RI and *Bam*HI restriction sites into the PCR product. Following PCR amplification, the fragment was digested with *Eco*RI and *Bam*HI and cloned into pBluescriptII cleaved with the same enzymes. For production of a probe specific for the sense strand of CON193, the CON193 Clone in pBluescriptII was linearized with *Bam*HI, which provided a substrate for labeled run-off transcripts (*i.e.*, cRNA riboprobes) using the vector-borne T7 promoter and commercially available T7 RNA polymerase. A probe specific for the antisense strand of CON193 was also readily prepared using the CON193 Clone in pBluescriptII by cleaving the recombinant plasmid with *Eco*RI to generate a linearized substrate for the production of labeled run-off cRNA transcripts using the T3 promoter and cognate polymerase. The riboprobes were labeled with [<sup>35</sup>S]-UTP to yield a specific activity of 0.81 x 10<sup>6</sup> cpm/pmol for antisense riboprobes and 0.55 x 10<sup>6</sup> cpm/pmol for sense-strand riboprobes. Both riboprobes were subsequently denatured by incubating at 70°C for 3 minutes and added (2 pmol/ml) to hybridization buffer which contained 50%

- 100 -

formamide, 10% dextran, 0.3 M NaCl, 10 mM Tris (pH 8.0), 1 mM EDTA, 1X Denhardt's Solution, and 10 mM dithiothreitol. Microscope slides containing sequential brain cryosections were independently exposed to 45 µl of hybridization solution per slide and silanized cover slips were placed over the sections being  
5 exposed to hybridization solution. Sections were incubated overnight (15-18 hours) at 52°C to allow hybridization to occur. Equivalent series of cryosections were exposed to sense or antisense CON193-specific cRNA riboprobes.

Following the hybridization period, coverslips were washed off the slides in 1X SSC. Slides were subjected to RNase A treatment by incubation in a  
10 buffer containing 20 µg/ml RNase A, 10 mM Tris (pH 8.0), 0.5 M NaCl and 1 mM EDTA for 45 minutes at 37°C. The cryosections were then subjected to three high-stringency washes in 0.1 X SSC at 52°C for 20 minutes each. Following the series of washes, cryosections were dehydrated by consecutive exposure to 70%, 95%, and 100% ammonium acetate in alcohol, followed by air drying and exposure to Kodak  
15 BioMax MR-1 film. After 13 days of exposure, the film was developed. Based on these results, brain sections that gave rise to positive hybridization signals were coated with Kodak NTB-2 nuclear track emulsion and the slides were stored in the dark for 32 days. The slides were then developed and counterstained with hematoxylin. Emulsion-coated sections were analyzed microscopically to determine the specificity  
20 of labeling. The signal was determined to be specific if autoradiographic grains (generated by antisense probe hybridization) were clearly associated with crystal violet-stained cell bodies. Autoradiographic grains found between cell bodies indicates non-specific binding.

Specific labeling with the antisense probe occurred at low levels in the  
25 cortex and in the substantia nigra-pars compacta (SN-c). The specificity of labeling was confirmed by microscopic analysis of emulsion-coated cryosections, as described above. In contrast, hybridization using the riboprobe specific for the sense strand of CON193 did not result in specific tissue labeling. The observed regional distribution of CON193 mRNA suggests that ligands for this GPCR may be involved in signal  
30 transductions important for cellular processes underlying neurological functioning. In addition, expression of CON193 in the brain provides an indication that modulators of

- 101 -

CON193 activity have utility for treating neurological disorders, including but not limited to, schizophrenia, depression, anxiety, bipolar disease, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, and the like. Use of CON193 modulators, including CON193 ligands and anti-CON193 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

#### B. CON166

A CON166-specific probe was generated using PCR as described above for CON193 in Example 3A (but using CON166-specific primers). The probe consisted of a 259 bp fragment containing sequence at the 3' end of CON-166 (nucleotides 715-974 of SEQ ID NO:1) and containing terminal *Eco*RI and *Bam*HI restriction sites. The riboprobes were labeled with [<sup>35</sup>S]-UTP to yield a specific activity of  $0.40 \times 10^6$  cpm/pmol for antisense riboprobes and  $0.65 \times 10^6$  cpm/pmol for sense-strand riboprobes. Hybridization with the riboprobes and subsequent washing of the slides was carried out as described above for CON193 in Example 3A.

Specific labeling with the antisense probe occurred in cortical regions, including the piriform complex, neostriatum, thalamus and hippocampus. The specificity of labeling was confirmed by microscopic analysis of emulsion-coated cryosections. These sections revealed that the autoradiographic grains resulting from antisense riboprobe *in situ* hybridizations were distributed over cell bodies rather than trapped between cell bodies. In contrast, hybridization using the riboprobe specific for the sense strand of CON166 produced a faint signal in the hippocampus only, but even this signal was found to be non-specific upon microscopic examination. The observed regional distribution of CON166 mRNA suggests that ligands for this GPCR may be involved in signal transductions important for cellular processes underlying neurological functioning. In addition, expression of CON166 in the brain provides an indication that modulators of CON166 activity have utility for treating neurological disorders, including but not limited to, schizophrenia, affective disorders, ADHD/ADD (*i.e.*, Attention Deficit-Hyperactivity Disorder/Attention Deficit Disorder), and neural disorders such as Alzheimer's disease, Parkinson's disease, migraine, and senile dementia. Some other diseases for which modulators of

- 102 -

CON166 may have utility include depression, anxiety, bipolar disease, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, and the like. Use of CON166 modulators, including CON166 ligands and anti-CON166 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

5

### C. CON 103

A cocktail of two CON103-specific antisense oligonucleotide probes (CON103a and CON103b) were used because of the relatively high GC content of the CON103 coding region. The CON103a sequence  
10 (5'TTTATTAATATTGGAAGGGACAACTGGAGAGCACAGAACAT3'; SEQ ID NO: 72) corresponds to the reverse complement of nucleotides 2196-2237 of SEQ ID NO: 5 and CON103b sequence (5'AAAGCCACCATGGA  
AGCCATGCCAAAGATGATGCTGGGCAAGAA 3'; SEQ ID NO: 73) corresponds to the reverse complement of nucleotides 195-1538 of SEQ ID NO: 5. Terminal  
15 deoxynucleotidyltransferase and [ $\alpha$ -<sup>33</sup>P]dATP were used to 3' end-label CON103a (1.36 x 10<sup>7</sup> cpm/pmol) and CON103b (9.1 x 10<sup>6</sup> cpm/pmol). The probes were denatured by incubation at 70°C for three minutes and added to hybridization buffer containing 50% formamide, 10% dextran, 0.3 M NaCl, 10 mM Tris (pH 8.0), 1 mM EDTA, 1X Denhardt's Solution, and 200 mM dithiothreitol. The final concentration  
20 of each radiolabeled probe was 2 pmol/ml of hybridization solution. Microscope slides containing sequential brain cryosections were independently exposed to 45  $\mu$ l of hybridization solution (containing the antisense oligonucleotide probes CON103a and CON103b) per slide and silanized cover slips were placed over the sections being exposed to hybridization solution. Sections were incubated overnight (15-18 hours) at  
25 37°C to allow hybridization to occur.

Following the hybridization period, coverslips were washed off the slides in 1X SSC. The cryosections were then subjected to three high-stringency washes in 1 X SSC at 65°C for 20 minutes each. Following two room-temperature washes, cryosections were dehydrated by consecutive exposure to 70%, 95%, and  
30 100% ethanol (0.3 M ammonium acetate added to 70% and 95% ethanol solutions), followed by air drying and exposure to Kodak BioMax MR-1 film. After 28 days of

- 103 -

exposure, the film was developed. Based on these results, brain sections that showed positive hybridization signals were coated with Kodak NTB-2 nuclear track emulsion and the slides were stored in the dark for four months. The slides were then developed and counterstained with hematoxylin. Emulsion-coated sections were analyzed microscopically to determine the specificity of labeling. The signal was determined to be specific if autoradiographic grains (generated by antisense probe hybridization) were present over cell bodies and not trapped between cell bodies.

Specific labeling with the antisense probe occurred in all cortical regions, including the piriform cortex and hippocampus. The specificity of labeling was confirmed by microscopic analysis of emulsion-coated cryosections. These sections revealed that the autoradiographic grains resulting from antisense riboprobe *in situ* hybridizations were distributed over cell bodies rather than trapped between cell bodies. The observed distribution of CON103 mRNA in the cortical and paralimbic regions of the mammalian brain suggests that ligands for this GPCR may be involved in signal transductions important for cellular processes underlying neurological functioning. In addition, expression of CON103 in the brain provides an indication that modulators of CON103 activity have utility for treating neurological and neuropsychiatric disorders, including but not limited to, schizophrenia, depression, anxiety, attention deficit disorder (with or without hyperactivity), bipolar disease, epilepsy, migraine, neuritis, neurasthenia, neuropathy, neuroses, obesity, Parkinson's disease, other dementias, and the like. Use of CON103 modulators, including CON103 ligands and anti-CON103 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

#### D. CON203

CON203-specific cRNA probes were prepared using conventional techniques. Initially, a 293 bp fragment of the CON203 coding region, with a *Bam*HI site and an *Eco*RI site disposed on opposite ends, was prepared by PCR using primers LW1314 (5'-GCATGAATTCCCACCTTCATCATCTACCTC-3'; SEQ ID NO: 40) and LW1315 (5'-GCATGGATCCGAAGACCAAAAAGACCCAG-3'; SEQ ID NO: 41). LW1314 includes an *Eco*RI site and additional protective residues at its 5'

- 104 -

terminus, with the rest of the sequence corresponding to CON203 coding nucleotides 164-183, which correspond to positions 309-328 of SEQ ID NO: 7. LW1315 includes 5' protective nucleotides and a *Bam*HI site, with the rest of the sequence corresponding to the complement of CON203 coding nucleotides 438-456, which correspond to positions 583-601 of SEQ ID NO: 7. The PCR-amplified fragment was then digested with *Bam*HI and *Eco*RI and ligated into the corresponding sites of pBluescript II to yield pCon203 BS. The recombinant clone was then linearized either with *Bam*HI or *Eco*RI. Linearization with *Bam*HI provided a substrate for *in vitro* expression of a sense-strand cRNA probe using the vector-borne T7 promoter. Digestion with *Eco*RI was used to provide a substrate for *in vitro* transcription using the vector-borne T3 promoter to generate an anti-sense cRNA probe. *In vitro* transcriptions were performed in the presence of [<sup>35</sup>S] UTP, thereby yielding sense- and anti-sense strand riboprobes having specific radioactivities of  $5.38 \times 10^7$  cpm/pmol and  $5.34 \times 10^7$  cpm/pmol, respectively. Hybridization with the riboprobes and subsequent washing of the slides was carried out as described above for CON193 in Example 3A. Subsequently, the slides were exposed to Kodak BioMax MR-1 film. After 9 days of exposure, the film was developed. Based on these results, brain sections that gave rise to positive hybridization signals were coated with Kodak NTB-2 nuclear track emulsion and the slides were stored in the dark for 25 days. The slides were then developed as described above for CON193 in Example 3A.

Specific labeling with the antisense probe occurred in several limbic and paralimbic regions, as well as areas thought to be involved in voluntary motor control. In particular, the probe hybridized to CON203 mRNAs in the following regions of the brain: cortical regions, including the piriform cortex, neostriatum, lateral olfactory tract, hypothalamic nuclei, bed nucleus of the stria terminalis, amygdala, hippocampus, reticular thalamus and other thalamic regions, subthalamic nucleus, and the red nucleus. The specificity of labeling was confirmed by microscopic analysis of emulsion-coated cryosections. These sections revealed that the autoradiographic grains resulting from antisense riboprobe *in situ* hybridizations were distributed over cell bodies rather than trapped between cell bodies. Confirming expression of CON203 mRNA, the sense-strand riboprobe did not show specific

- 105 -

hybridization. The observed distribution of CON203 mRNA in the cortical (particularly, motor circuits) and paralimbic regions of the mammalian brain suggests that CON203 and the ligands for this GPCR may be involved in signal transductions important for cellular processes underlying neurological functioning. In addition, expression of CON203 in the brain provides an indication that modulators of CON203 activity have utility for treating neurological disorders, including but not limited to, schizophrenia, depression, anxiety, bipolar disease, epilepsy, migraine, attention deficit disorder (with or without hyperactivity), neuritis, neurasthenia, neuropathy, neuroses, Parkinson's disease, dementia, obesity, and the like. Use of CON203 modulators, including CON203 ligands and anti-CON203 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

#### E. CON198

A 266 bp fragment of CON198 containing *EcoRI* and *BamHI* restriction sites was amplified from the full-length clone by PCR, using the primers LW1308: 5'-GCATGAATTC<sup>ACTCACTTCTCATCTCCTTC</sup>-3' (SEQ ID NO: 46) and LW1309: 5'-GCATGGATCC<sup>AATCTCCTTTGTCTTCACTC</sup>-3' (SEQ ID NO: 47). Primer LW1308 contains an *EcoRI* site (underlined) followed by sequence identical to nucleotides 638-657 of SEQ ID NO: 9. Primer LW1309 contains a *BamHI* site (underlined) followed by sequence complementary to nucleotides 903-884 of SEQ ID NO: 9. The amplification product was digested with *EcoRI* and *BamHI*, and then subcloned into an *EcoRI*- and *BamHI*-digested pBluescript II vector (*Stratagene*). The 266 amplified and subcloned basepairs correspond to nucleotides 638 to 903 of SEQ ID NO: 9.

The subcloned CON198-Bluescript construct was used to generate strand-specific probes for the *in situ* hybridization experiments. The construct was linearized with *BamHI*, for labeling with T7 polymerase (sense), or *EcoRI*, for T3 polymerase (antisense), and used as a template for *in vitro* transcription of sense and antisense cRNA riboprobes. The riboprobes were labeled with <sup>35</sup>S-UTP to yield a specific activity of 0.45 x 10<sup>6</sup> cpm/pmol for antisense and 0.732 x 10<sup>6</sup> cpm/pmol for

- 106 -

sense probe. Hybridization with the riboprobes and subsequent washing of the slides was carried out as described above for CON193 in Example 3A.

Specific labeling with the antisense probe showed distribution of CON198 mRNA in the rat brain in several limbic and paralimbic regions as well as areas thought to be involved in voluntary motor control. Labelled regions included cortical regions, piriform cortex, hypothalamic nuclei (paraventricular nucleus, supraoptic nucleus, suprachiasmatic nucleus), hippocampus, reticular thalamus, substantia nigra-pars compacta (SN-C), ventral tegmental area, and the red nucleus. The specificity of labeling was confirmed by microscopic analysis of emulsion coated sections. These sections revealed that the autoradiographic grains generated by the antisense probe were distributed over cell bodies rather than trapped between cell bodies. Sense probe did not generate specific labeling.

The observed regional distribution of CON198 mRNA provides a therapeutic indication for natural ligands for CON198 as well as modulators of CON198 activity, such as anti-CON198 antibody substances or small molecules that agonize or antagonize ligand-mediated CON198 signalling. In particular, the expression pattern provides an indication that such molecules will have utility for treating neurological and/or psychiatric diseases, including but not limited to schizophrenia, depression, anxiety, bipolar disease, affective disorders, ADHD/ADD, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, Alzheimer's disease, Parkinson's disease, migraine, senile dementia, and the like. Use of CON198 modulators, including CON198 ligands and anti-CON198 antibodies, to treat individuals having such disease states is intended as an aspect of the invention. Such modulators are administered by any means effective to safely deliver the modulators to the CON198-expressing cells, including but not limited to oral administration, inhalation, or injection of compositions comprising the modulators in a pharmaceutically acceptable diluent, adjuvant, or carrier. Efficacy of treatment can initially be determined in any accepted animal model that provides a biochemical or behavioral marker that correlates with disease severity or treatment efficacy.

- 107 -

**F. CON197**

A 261 bp fragment of CON197 containing *Eco*RI and *Bam*HI restriction sites was amplified from the full-length clone by PCR, using the primers LW1306: 5'-GCATGAATTCTTCTACTTCATCATCCTCC-3' (SEQ ID NO: 50) and  
5 LW1307: 5'-GCATGGATCCAAAGGCCATCACAACAAG-3' (SEQ ID NO: 51).  
Primer LW1306 includes sequence identical to nucleotides 100-118 of SEQ ID NO: 11 (underlined), preceded by an *Eco*RI site. Primer LW1307 includes sequence complementary to nucleotides 361-343 of SEQ ID NO: 11 (underlined), preceded by a *Bam*HI restriction site. The amplification product was digested with *Eco*RI and  
10 *Bam*HI, and then subcloned into an *Eco*RI- and *Bam*HI-digested pBluescript II vector (Stratagene). The 261 amplified and subcloned basepairs correspond to nucleotides 100 to 361 of SEQ ID NO: 11.

The subcloned CON197-Bluescript construct was used to generate strand-specific probes for the *in situ* hybridization experiments. The construct was  
15 linearized with *Bam*HI, for labeling with T7 polymerase (sense), or *Eco*RI, for T3 polymerase (antisense), and used as a template for *in vitro* transcription of sense and antisense cRNA riboprobes. The riboprobes were labeled with <sup>35</sup>S-UTP to yield a specific activity of 0.51 x 10<sup>6</sup> cpm/pmol for antisense and 0.432 x 10<sup>6</sup> cpm/pmol for sense probe. Hybridization with the riboprobes and subsequent washing of the slides  
20 was carried out as described above for CON193 in Example 3A.

Specific labeling with the antisense probe showed wide spread distribution of CON197 mRNA in the rat brain. Labelled regions included neo and allo cortex, piriform cortex, neostriatum, thalamic nuclei, hypothalamic nuclei, hippocampus, amygdala, cerebellum, and the olfactory bulb. The specificity of  
25 labeling was confirmed by microscopic analysis of emulsion coated sections. These sections revealed that the autoradiographic grains generated by the antisense probe were distributed over cell bodies rather than trapped between cell bodies. Sense probe did not generate specific labeling.

The observed regional distribution of CON197 mRNA provides a  
30 therapeutic indication for natural ligands for CON197 as well as modulators of CON197 activity, such as anti-CON197 antibody substances or small molecules that

- 108 -

agonize or antagonize ligand-mediated CON197 signalling. In particular, the expression pattern provides an indication that such molecules will have utility for treating neurological and/or psychiatric diseases, including but not limited to dementia, schizophrenia, depression, anxiety, bipolar disease, migraine, Parkinson's disease, affective disorders, Alzheimer's disease, senile dementia, attention deficit hyperactivity disorder/attention deficit disorder (ADHD/ADD), epilepsy, neuritis, neurasthenia, neuropathy, neuroses, and the like. Use of CON197 modulators, including CON197 ligands and anti-CON197 antibodies, to treat individuals having such disease states is intended as an aspect of the invention. Such modulators are administered by any means effective to safely deliver the modulators to the CON197-expressing cells, including but not limited to oral administration, inhalation, or injection of compositions comprising the modulators in a pharmaceutically acceptable diluent, adjuvant, or carrier. Efficacy of treatment can initially be determined in any accepted animal model that provides a biochemical or behavioral marker that correlates with disease severity or treatment efficacy.

**G. CON202**

A 272 bp fragment of CON202 containing EcoRI and BamHI restriction sites was amplified from the full-length clone by PCR, using the primers LW1310 GCATGAATTTCGCAGAAGAAGGCTATTGG (SEQ ID NO: 56) and LW1311 GCATGGATCCGCAGTAAAGAAGGGTTGTG (SEQ ID NO: 57). The amplification product was digested with EcoRI and BamHI, and then subcloned into a pBluescript II vector (Stratagene) that was digested with EcoRI and BamHI. The 272 amplified and subcloned basepairs correspond to nucleotides 1065 to 1336 of SEQ ID NO: 13.

The subcloned CON202-Bluescript construct was used to generate strand-specific probes for the *in situ* hybridization experiments. The construct was linearized with BamHI, for labeling with T7 polymerase (sense), or EcoRI, for T3 polymerase (antisense), and used as a template for *in vitro* transcription of sense and antisense cRNA riboprobes. The riboprobes were labeled with <sup>35</sup>S-UTP to yield a specific activity of 4.7 x 10<sup>5</sup> cpm/pmol for antisense and 4.3 x 10<sup>5</sup> cpm/pmol for sense

- 109 -

probe. Hybridization with the riboprobes and subsequent washing of the slides was carried out as described above for CON193 in Example 3A.

Specific labeling with the antisense probe showed wide spread distribution of CON202 mRNA in the rat brain. Labelled regions included the cortical regions, lateral olfactory nuclei, hippocampus, subthalamic nucleus, and at a lower level, the nigra-pars compacta.

The observed regional distribution of CON202 mRNA provides a therapeutic indication for natural ligands for CON202 as well as modulators of CON202 activity, such as anti-CON202 antibody substances or small molecules that agonize or antagonize ligand-mediated CON202 signaling. In particular, the expression pattern provides an indication that such molecules will have utility for treating neurological and/or psychiatric diseases, including but not limited to schizophrenia, affective disorders, attention deficit hyperactivity disorder/attention deficit disorder, depression, anxiety, bipolar disease, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, Alzheimer's disease, Parkinson's disease, migraine, senile dementia and the like. Use of CON202 modulators, including CON202 ligands and anti-CON202 antibodies, to treat individuals having such disease states is intended as an aspect of the invention. Such modulators are administered by any means effective to safely deliver the modulators to the CON202-expressing cells, including but not limited to oral administration, inhalation, or injection of compositions comprising the modulators in a pharmaceutically acceptable diluent, adjuvant, or carrier. Efficacy of treatment can initially be determined in any accepted animal model that provides a biochemical or behavioral marker that correlates with disease severity or treatment efficacy.

#### H. CON222

A 264 bp fragment of CON222 containing EcoRI and BamHI restriction sites was amplified from the full-length clone by PCR, using the primers LW1472 (5'GCATGAATTCTGCCATGTCAATCATTTCTCTC3'; SEQ ID NO: 62, EcoRI site is underlined) and LW1473 (5'GCATGGATCCGTTCTGCATTTTCC-AGGTCTC3'; SEQ ID NO: 63, BamHI site is underlined). The amplification product

- 110 -

was digested with EcoRI and BamHI, and then subcloned into a predigested pBluescript II vector (Stratagene). The 264 amplified and subcloned basepairs correspond to nucleotides 237 to 500 of SEQ ID NO: 15.

5 The subcloned CON222-Bluescript construct was used to generate strand-specific probes for the *in situ* hybridization experiments. The construct was linearized with BamHI, for labeling with T7 polymerase (sense), or EcoRI, for T3 polymerase (antisense), and used as a template for *in vitro* transcription of sense and antisense cRNA riboprobes. The riboprobes were labeled with <sup>35</sup>S-UTP to yield a specific activity of  $4.25 \times 10^5$  cpm/pmol for antisense and  $3.9 \times 10^5$  cpm/pmol for sense probe. Hybridization with the riboprobes and subsequent washing of the slides was carried out as described above for CON193 in Example 3A.

15 Specific labeling with the antisense probe showed wide spread distribution of CON222 mRNA in the rat brain. Labelled regions included the cortical regions, piriform cortex, striatum, hippocampus, thalamus, hypothalamus, dorsal raphe, and habenula.

The observed regional distribution of CON222 mRNA provides a therapeutic indication for natural ligands for CON222 as well as modulators of CON222 activity, such as anti-CON222 antibody substances or small molecules that agonize or antagonize ligand-mediated CON222 signaling. In particular, the expression pattern provides an indication that such molecules will have utility for treating neurological and/or psychiatric diseases, including but not limited to schizophrenia, affective disorders, attention deficit hyperactivity disorder/attention deficit disorder, depression, anxiety, bipolar disease, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, Alzheimer's disease, Parkinson's Disease, migraine, senile dementia, and the like. Use of CON222 modulators, including CON222 ligands and anti-CON222 antibodies, to treat individuals having such disease states is intended as an aspect of the invention. Such modulators are administered by any means effective to safely deliver the modulators to the CON222-expressing cells, including but not limited to oral administration, inhalation, or injection of compositions comprising the modulators in a pharmaceutically acceptable diluent, adjuvant, or carrier. Efficacy of treatment can initially be determined in any accepted animal model that provides a

20  
25  
30

- 111 -

biochemical or behavioral marker that correlates with disease severity or treatment efficacy.

# I. CON215

5 A 261 bp fragment of CON215 containing *Eco*RI and *Bam*HI restriction sites was amplified from the full-length clone by PCR, using the primers LW1411: 5'-GCATGAATTCTGCCAAACATCATCCTGAC-3' (SEQ ID NO: 64) and LW1412: 5'-GCATGGATCCTACACAGCCACAACAACCC-3' (SEQ ID NO: 65). Primer LW1411 contains an *Eco*RI site (underlined) followed by sequence identical to CON215 coding nucleotides 521-537, which correspond to positions 533-10 549 of SEQ ID NO: 17. Primer LW1412 contain a *Bam*HI site (underlined) followed by sequence complementary to CON215 coding nucleotides 764-781, which correspond to positions 776-793 of SEQ ID NO: 17. The amplification product was digested with *Eco*RI and *Bam*HI, and then subcloned into an *Eco*RI- and *Bam*HI-15 digested pBluescript II vector (*Stratagene*). The 261 amplified and subcloned basepairs correspond to nucleotides 521 to 781 of SEQ ID NO: 17.

The subcloned CON215-Bluescript construct was used to generate strand-specific probes for the *in situ* hybridization experiments. The construct was linearized with *Bam*HI, for labeling with T7 polymerase (sense), or *Eco*RI, for T320 polymerase (antisense), and used as a template for *in vitro* transcription of sense and antisense cRNA riboprobes. The riboprobes were labeled with <sup>35</sup>S-UTP to yield a specific activity of 48.03 x 10<sup>6</sup> cpm/pmol for antisense and 48.09 x 10<sup>6</sup> cpm/pmol for sense probe. Hybridization with the riboprobes and subsequent washing of the slides was carried out as described above for CON193 in Example 3A.

25 Subsequently, the slides were exposed to Kodak BioMax MR-1 film. After 9 days of exposure, the film was developed. Slides containing sections that showed a hybridization signal on film autoradiograms were coated with Kodak NTB-2 nuclear track emulsion and stored in the dark for 25 days. The slides were then developed as described above for CON193 in Example 3A.

30 Specific labeling with the antisense probe showed distribution of CON215 mRNA in the rat brain in limbic endocrine and motor circuits. Specifically,

- 112 -

CON215 mRNA was present in the cortex, hippocampus, and red nucleus. The specificity of labeling was confirmed by microscopic analysis of emulsion coated sections. These sections revealed that the autoradiographic grains generated by the antisense probe were distributed over cell bodies rather than trapped between cell  
5 bodies. Sense probe did not generate specific labeling.

The observed regional distribution of CON215 mRNA provides a therapeutic indication for natural ligands for CON215 as well as modulators of CON215 activity, such as anti-CON215 antibody substances or small molecules that agonize or antagonize ligand-mediated CON215 signaling. In particular, the  
10 expression pattern provides an indication that such molecules will have utility for treating neurological and/or psychiatric diseases, including but not limited to schizophrenia, depression, anxiety, bipolar disease, epilepsy, migraine, attention deficit (with or without hyperactive disorder), neuritis, neuasthenia, neuropathy, neuroses, Parkinson's disease, dementia, obesity, and the like. Use of CON215  
15 modulators, including CON215 ligands and anti-CON215 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

Such modulators are administered by any means effective to safely deliver the modulators to the CON215-expressing cells, including but not limited to oral administration, inhalation, or injection of compositions comprising the  
20 modulators in a pharmaceutically acceptable diluent, adjuvant, or carrier. Efficacy of treatment can initially be determined in any accepted animal model that provides a biochemical or behavioral marker that correlates with disease severity or treatment efficacy.

#### 25 J. CON 217

Two oligonucleotides were designed based on SEQ ID NO: 19 and obtained from Sigma-Genosys (St. Louis, MO) to use as probes for *in situ* hybridization. The first oligonucleotide, designated 217A, has the sequence  
5'TAGGTCGGTAGTCAGGACACGGGAGAACAGAACTGTTGGTTGA3' (SEQ  
30 ID NO: 68) which is complementary to nucleotides 102 to 60 of SEQ ID NO: 19. The second oligonucleotide, designated 217B, has the sequence

- 113 -

5'GCCCCCTGTGGCGGTTTAGATCCAGAATGCCCATTTTCTGTTCCATCTAAC  
CA3' (SEQ ID NO: 69) which corresponds to the complement of nucleotides 1530 to  
1479 of SEQ ID NO: 17. Both oligonucleotides, 217A and 217B, were reconstituted  
with 1x TE buffer to a concentration of 20 pMol/ml and labeled with <sup>33</sup>P-dATP to  
5 yield a specific activity of 2.08 x 10<sup>6</sup> and 1.53 x 10<sup>6</sup> cpm/ml, respectively.

Hybridization was carried out at 37°C overnight as described above for  
CON193 in Example 3A. Following the hybridizations, the coverslips were washed  
off the slides with 1x SSC for 45 minutes. The slides were then washed for 20  
minutes at room temperature in 1x SSC followed by three high stringency washes in  
10 1x SSC at 65°C. After washing, the slides were dehydrated with 70%, 95%, and  
100% ethanol containing 0.3 mM NH<sub>4</sub>OAc, air-dried, and exposed to Kodak BioMax  
MR-1 film. After 21 days of exposure, the film was developed. Based on these  
results, sections that showed a hybridization signal on film autoradiography were  
coated with Kodak NTB-2 nuclear track emulsion and stored in the dark for 42 days.  
15 The slides were then developed and counterstained with hematoxylin. Emulsion-  
coated sections were analyzed microscopically to determine the specificity of labeling.  
The signal was judged to be specific if autoradiographic grains (generated by  
antisense probe hybridization) were associated clearly with crystal violet stained cell  
bodies. Autoradiographic grains found between cell bodies were deemed non-  
20 specific.

Specific labeling with the antisense probe showed wide spread  
distribution of CON217 mRNA in the rat brain. Labelled regions included the cortex,  
piriform cortex, hippocampus, cerebellum, medulla, spinal cord, temporal lobe,  
putamen, substantia nigra and thalamus.

25 The observed regional distribution of CON217 mRNAs provide a  
therapeutic indication for natural ligands for these G protein-coupled receptors as well  
as modulators of their activity, such as anti-CON217 antibody substances or small  
molecules that mimic, agonize or antagonize ligand-mediated CON217 signaling. In  
particular, the expression patterns provide an indication that such molecules will have  
30 utility for treating neurological and/or psychiatric diseases, including but not limited  
to schizophrenia, affective disorders, attention deficit hyperactivity disorder/attention

- 114 -

deficit disorder, depression, anxiety, bipolar disease, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, Alzheimer's disease, Parkinson's Disease, migraine, senile dementia, and the like. Use of CON217 polypeptide modulators, including CON217 ligands and anti-CON217 polypeptide antibodies, to treat individuals having such

5 disease states is intended as an aspect of the invention. Such modulators are administered by any means effective to safely deliver the modulators to the GPCR polypeptide-expressing cells, including but not limited to oral administration, inhalation, or injection of compositions comprising the modulators in a pharmaceutically acceptable diluent, adjuvant, or carrier. Efficacy of treatment can

10 initially be determined in any accepted animal model that provides a biochemical or behavioral marker that correlates with disease severity or treatment efficacy.

#### EXAMPLE 4

##### Recombinant Expression of GPCR Polypeptides in Eukaryotic Host Cells

15 To produce GPCR protein, a GPCR polypeptide-encoding polynucleotide is expressed in a suitable host cell using a suitable expression vector, using standard genetic engineering techniques. For example, one of the GPCR polypeptide-encoding sequences described in Example 1 (such as SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17 or 19) is subcloned into the commercial expression vector

20 pzeoSV2 (Invitrogen, San Diego, CA) and transfected into Chinese Hamster Ovary (CHO) cells (ATCC CRL-1781) using the transfection reagent fuGENE 6 (Boehringer-Mannheim) and the transfection protocol provided in the product insert. Additional eukaryotic cell lines, such as African Green Monkey Kidney cells (COS-7, ATCC CRL-1651) or Human Kidney cells (HEK 293, ATCC CRL-1573), may be

25 used as well. Cells stably expressing a GPCR polypeptide (e.g., CON193, CON166, CON103, CON203, CON198, CON197, CON202, CON222, CON215, or CON217) are selected by growth in the presence of 100 mg/ml zeocin (Stratagene, LaJolla, CA). Optionally, GPCR polypeptide is purified from the cells using standard chromatographic techniques. To facilitate purification, antisera is raised against one

30 or more synthetic peptide sequences that correspond to portions of the GPCR amino acid sequence, and the antisera is used to affinity purify GPCR polypeptides. The

- 115 -

GPCR gene also may be expressed in frame with a tag sequence (*e.g.*, polyhistidine, hemagglutinin, FLAG) to facilitate purification. Moreover, it will be appreciated that many of the uses for GPCR polypeptides, such as assays described below, do not require purification of GPCR polypeptides from the host cell.

5

## EXAMPLE 5

### Antibodies to GPCR Polypeptides

Standard techniques are employed to generate polyclonal or monoclonal antibodies to the GPCR receptors (*e.g.*, CON193, CON166, CON103, CON203, CON198, CON197, CON202, CON222, CON215, or CON217), and to generate useful antigen-binding fragments thereof or variants thereof, including “humanized” variants. Such protocols can be found, for example, in Sambrook *et al.*, *Molecular Cloning: a Laboratory Manual*. Second Edition, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory (1989); Harlow *et al.* (Eds), *Antibodies A Laboratory Manual*; Cold Spring Harbor Laboratory; Cold Spring Harbor, NY (1988); and other documents cited below. In one embodiment, recombinant GPCR polypeptides (or cells or cell membranes containing such polypeptides) of the invention are used as an antigen to generate the antibodies. In another embodiment, one or more peptides having amino acid sequences corresponding to an immunogenic portion of a GPCR polypeptide (*e.g.*, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more amino acids) are used as antigen. Peptides corresponding to extracellular portions of GPCR polypeptides, especially hydrophilic extracellular portions, are preferred. The antigen may be mixed with an adjuvant or linked to a hapten to increase antibody production.

25

#### A. Polyclonal or Monoclonal antibodies

As one exemplary protocol, a recombinant GPCR polypeptide or synthetic fragment thereof is used to immunize a mouse for generation of monoclonal antibodies (or larger mammal, such as a rabbit, for polyclonal antibodies). To increase antigenicity, peptides are conjugated to Keyhole Limpet Hemocyanine (Pierce), according to the manufacturer's recommendations. For an initial injection,

30

- 116 -

the antigen is emulsified with Freund's Complete Adjuvant and injected subcutaneously. At intervals of two to three weeks, additional aliquots of GPCR antigen are emulsified with Freund's Incomplete Adjuvant and injected subcutaneously. Prior to the final booster injection, a serum sample is taken from the immunized mice and assayed by Western blot to confirm the presence of antibodies that immunoreact with GPCR polypeptide. Serum from the immunized animals may be used as a polyclonal antisera or used to isolate polyclonal antibodies that recognize GPCR polypeptide. Alternatively, the mice are sacrificed and their spleen removed for generation of monoclonal antibodies.

To generate monoclonal antibodies, the spleens are placed in 10 ml serum-free RPMI 1640, and single cell suspensions are formed by grinding the spleens in serum-free RPMI 1640, supplemented with 2 mM L-glutamine, 1 mM sodium pyruvate, 100 units/ml penicillin, and 100 µg/ml streptomycin (RPMI) (Gibco, Canada). The cell suspensions are filtered and washed by centrifugation and resuspended in serum-free RPMI. Thymocytes taken from three naive Balb/c mice are prepared in a similar manner and used as a Feeder Layer. NS-1 myeloma cells, kept in log phase in RPMI with 10% fetal bovine serum (FBS) (Hyclone Laboratories, Inc., Logan, Utah) for three days prior to fusion, are centrifuged and washed as well.

To produce hybridoma fusions, spleen cells from the immunized mice are combined with NS-1 cells and centrifuged, and the supernatant is aspirated. The cell pellet is dislodged by tapping the tube, and 2 ml of 37°C PEG 1500 (50% in 75mM Hepes, pH 8.0) (Boehringer Mannheim) is stirred into the pellet, followed by the addition of serum-free RPMI. Thereafter, the cells are centrifuged and resuspended in RPMI containing 15% FBS, 100 µM sodium hypoxanthine, 0.4 µM aminopterin, 16 µM thymidine (HAT) (Gibco), 25 units/ml of IL-6 (Boehringer Mannheim) and  $1.5 \times 10^6$  thymocytes/ml and plated into 10 Corning flat-bottom 96-well tissue culture plates (Corning, Corning New York).

On days 2, 4, and 6, after the fusion, 100 µl of medium is removed from the wells of the fusion plates and replaced with fresh medium. On day 8, the fusions are screened by ELISA, testing for the presence of mouse IgG that binds to a

- 117 -

GPCR polypeptide. Selected fusion wells are further cloned by dilution until monoclonal cultures producing anti-GPCR polypeptide antibodies are obtained.

**B. Humanization of Anti-GPCR Monoclonal Antibodies**

5                   The expression patterns of GPCR polypeptides as reported herein and the proven track record of GPCR's as targets for therapeutic intervention suggest therapeutic indications for GPCR polypeptide inhibitors (antagonists). GPCR polypeptide-neutralizing antibodies comprise one class of therapeutics useful as antagonists. Following are protocols to improve the utility of anti-GPCR polypeptide  
10                   monoclonal antibodies as therapeutics in humans, by "humanizing" the monoclonal antibodies to improve their serum half-life and render them less immunogenic in human hosts (*i.e.*, to prevent human antibody response to non-human anti-GPCR polypeptide antibodies).

                  The principles of humanization have been described in the literature  
15                   and are facilitated by the modular arrangement of antibody proteins. To minimize the possibility of binding complement, a humanized antibody of the IgG4 isotype is preferred.

                  For example, a level of humanization is achieved by generating chimeric antibodies comprising the variable domains of non-human antibody proteins  
20                   of interest with the constant domains of human antibody molecules. (See, *e.g.*, Morrison and Oi, *Adv. Immunol.*, 44:65-92 (1989). The variable domains of GPCR-neutralizing anti-GPCR antibodies are cloned from the genomic DNA of a B-cell hybridoma or from cDNA generated from mRNA isolated from the hybridoma of interest. The V region gene fragments are linked to exons encoding human antibody  
25                   constant domains, and the resultant construct is expressed in suitable mammalian host cells (*e.g.*, myeloma or CHO cells).

                  To achieve an even greater level of humanization, only those portions of the variable region gene fragments that encode antigen-binding complementarity determining regions ("CDR") of the non-human monoclonal antibody genes are  
30                   cloned into human antibody sequences. [See, *e.g.*, Jones *et al.*, *Nature*, 321:522-525 (1986); Riechmann *et al.*, *Nature*, 332:323-327 (1988); Verhoeyen *et al.*, *Science*,

- 118 -

239:1534-36 (1988); and Tempest *et al.*, *Bio/Technology*, 9:266-71 (1991). If necessary, the  $\beta$ -sheet framework of the human antibody surrounding the CDR3 regions also is modified to more closely mirror the three dimensional structure of the antigen-binding domain of the original monoclonal antibody. (See Kettleborough  
5 *et al.*, *Protein Engin.*, 4:773-783 (1991); and Foote *et al.*, *J. Mol. Biol.*, 224:487-499 (1992).

In an alternative approach, the surface of a non-human monoclonal antibody of interest is humanized by altering selected surface residues of the non-human antibody, *e.g.*, by site-directed mutagenesis, while retaining all of the  
10 interior and contacting residues of the non-human antibody. See Padlan, *Molecular Immunol.*, 28(4/5):489-98 (1991).

The foregoing approaches are employed using GPCR-neutralizing anti-GPCR monoclonal antibodies and the hybridomas that produce them to generate humanized GPCR-neutralizing antibodies useful as therapeutics to treat or palliate  
15 conditions wherein GPCR expression or ligand-mediated GPCR signaling is detrimental.

### C. Human GPCR-Neutralizing Antibodies from Phage Display

Human GPCR-neutralizing antibodies are generated by phage display  
20 techniques such as those described in Aujame *et al.*, *Human Antibodies*, 8(4):155-168 (1997); Hoogenboom, *TIBTECH*, 15:62-70 (1997); and Rader *et al.*, *Curr. Opin. Biotechnol.*, 8:503-508 (1997), all of which are incorporated by reference. For example, antibody variable regions in the form of Fab fragments or linked single chain Fv fragments are fused to the amino terminus of filamentous phage minor coat  
25 protein pIII. Expression of the fusion protein and incorporation thereof into the mature phage coat results in phage particles that present an antibody on their surface and contain the genetic material encoding the antibody. A phage library comprising such constructs is expressed in bacteria, and the library is panned (screened) for GPCR-specific phage-antibodies using labelled or immobilized GPCR polypeptide as  
30 antigen-probe.

- 119 -

**D. Human GPCR-Neutralizing Antibodies from Transgenic Mice**

Human GPCR-neutralizing antibodies are generated in transgenic mice essentially as described in Bruggemann and Neuberger, *Immunol. Today*, 17(8):391-97 (1996) and Bruggemann and Taussig, *Curr. Opin. Biotechnol.*, 8:455-58 (1997). Transgenic mice carrying human V-gene segments in germline configuration and that express these transgenes in their lymphoid tissue are immunized with a GPCR composition using conventional immunization protocols. Hybridomas are generated using B cells from the immunized mice using conventional protocols and screened to identify hybridomas secreting anti-GPCR human antibodies (e.g., as described above).

**EXAMPLE 6****Assays to Identify Modulators of GPCR Polypeptide Activity**

Set forth below are assays for identifying modulators (agonists and antagonists) of GPCR polypeptide activity. Among the modulators that can be identified by these assays include natural ligand compounds of the receptor; synthetic analogs and derivatives of natural ligands; antibodies, antibody fragments, and/or antibody-like compounds derived from natural antibodies or from antibody-like combinatorial libraries; and/or synthetic compounds identified through high throughput screening of libraries; and the like. All modulators that bind GPCR polypeptide are useful for identifying GPCR polypeptide in tissue samples (e.g., for diagnostic purposes, pathological purposes, and the like). Agonist and antagonist modulators are useful for up-regulating and down-regulating GPCR polypeptide activity, respectively; to treat disease states characterized by abnormal levels of GPCR polypeptide activity. GPCR polypeptide binding molecules also may be used to deliver a therapeutic compound or a label to cells that express GPCR polypeptide (e.g., by attaching the compound or label to the binding molecule). The assays may be performed using single putative modulators, and/or may be performed using a known agonist in combination with candidate antagonists (or *visa versa*). Performance of the assays using any of the GPCR polypeptides of the invention described herein (e.g., CON193, CON166, CON103, CON203, CON198, CON197,

- 120 -

CON202, CON222, CON215, or CON217) is contemplated. It will be appreciated that co-transfecting cells with two or more of the receptors for simultaneous screening also is possible.

5       A.     cAMP Assays

              In one type of assay, levels of cyclic adenosine monophosphate (cAMP) are measured in GPCR-transfected cells that have been exposed to candidate modulator compounds. Protocols for cAMP assays have been described in the literature. [See, e.g., Sutherland *et al.*, *Circulation*, 37: 279 (1968); Frandsen, E.K. and Krishna, G, *Life Sciences*, 18: 529-541 (1976); Dooley *et al.*, *Journal of Pharmacology and Experimental Therapeutics*, 283 (2): 735-41 (1997); and George *et al.*, *Journal of Biomolecular Screening*, 2 (4): 235-40 (1997).] An exemplary protocol for such an assay, using an Adenylyl Cyclase Activation FlashPlate® Assay from NEN™ Life Science Products, is set forth below.

15             Briefly, the GPCR coding sequence (e.g., a cDNA or intronless genomic DNA) is subcloned into a commercial expression vector, such as pzeoSV2 (Invitrogen, San Diego, CA), and transiently transfected into Chinese Hamster Ovary (CHO) cells using known methods, such as the transfection reagent FuGENE 6 (Boehringer-Mannheim) and the transfection protocol provided in the product insert.

20             The transfected CHO cells are seeded into the 96 well microplates from the FlashPlate® assay kit, which are coated with solid scintillant to which antisera to cAMP has been bound. For a control, some wells are seeded with wild type (untransfected) CHO cells. Other wells on the plate receive various amounts of cAMP standard solution for use in creating a standard curve.

25             One or more test compounds are added to the cells in each well, with water and/or compound-free media/diluent serving as a control. After treatment, cAMP is allowed to accumulate in the cells for exactly 15 minutes at room temperature. The assay is terminated by the addition of lysis buffer containing [<sup>125</sup>I]-labelled cAMP, and the plate is counted using a Packard Topcount™ 96-well  
30             microplate scintillation counter. Unlabelled cAMP from the lysed cells (or from standards) competes with the fixed amounts of [<sup>125</sup>I]-cAMP for antibody bound to the

- 121 -

plate. A standard curve is constructed, and cAMP values for the unknowns are obtained by interpolation. Changes in intracellular cAMP level of the cells in response to exposure to a test compound are indicative of GPCR polypeptide modulating activity. Modulators that act as agonists at receptors which couple to the Gs subtype of G-proteins will stimulate production of cAMP, leading to a measurable 3-10 fold increase. Receptor agonists which couple to the Gi/o subtype of G-proteins will inhibit forskolin-stimulated cAMP production, leading to a measurable decrease of 50-100%. Modulators that act as inverse agonists will reverse these effects at receptors that are either constitutively active or activated by known agonists.

#### B. Aequorin Assays

In another assay cells (*e.g.*, CHO cells) are transiently co-transfected with both a GPCR expression construct and a construct that encodes the photoprotein apoaequorin. In the presence of the cofactor coelenterazine, apoaequorin will emit a measurable luminescence that is proportional to the amount of intracellular (cytoplasmic) free calcium. [See generally Cobbold P.H. and Lee, J.A.C. "Aequorin measurements of cytoplasmic free calcium. In: McCormack J.G. and Cobbold P.H., eds., *Cellular Calcium: A Practical Approach*. Oxford:IRL Press (1991); Stables *et al.*, *Analytical Biochemistry*, 252: 115-26 (1997); and Haugland, R.P. *Handbook of Fluorescent Probes and Research Chemicals*. Sixth edition. Eugene OR: Molecular Probes (1996).]

In one exemplary assay, a GPCR-encoding polynucleotide is subcloned into the commercial expression vector pzeoSV2 (Invitrogen, San Diego, CA) and transiently co-transfected along with a construct that encodes the photoprotein apoaequorin (Molecular Probes, Eugene, OR) into CHO cells using the transfection reagent FuGENE 6 (Boehringer-Mannheim) and the transfection protocol provided in the product insert.

The cells are cultured for 24 hours at 37°C in  $\alpha$ MEM (Gibco/BRL, Gaithersburg, MD) supplemented with 10% FBS, 2 mM glutamine, 10 U/ml of penicillin and 10  $\mu$ g/ml of streptomycin. Subsequently, the media is changed to serum-free  $\alpha$ MEM containing 5  $\mu$ M coelenterazine (Molecular Probes, Eugene, OR),

- 122 -

and the cells are cultured for two additional hours at 37°C. Cells are then detached from the plate using VERSEN (Gibco/BRL), washed and resuspended at  $2 \times 10^5$  cells/ml in serum-free  $\alpha$ MEM.

Dilutions of candidate GPCR modulator drugs are prepared in serum-free  $\alpha$ MEM and dispensed into wells of an opaque 96-well assay plate, 50  $\mu$ l/well. Plates are loaded onto an MLX microtiter plate luminometer (Dynex Technologies, Inc., Chantilly, VA). The instrument is programmed to dispense 50  $\mu$ l of cell suspension into each well, one well at a time, and immediately read luminescence for 15 seconds. Dose-response curves for the modulator candidates are constructed using the area under the curve for each light signal peak. Data are analyzed with SlideWrite, using the equation for 1-site ligand, and  $EC_{50}$  values are obtained. Changes in luminescence caused by the drugs are considered indicative of modulatory activity. Modulators that act as receptor agonists which couple to the Gq subtype of G-proteins give an increase in luminescence of up to 100 fold. Modulators that act as inverse agonists will reverse this effect at receptors that are either constitutively active or activated by known agonists.

#### C. Luciferase Reporter Gene Assay

The photoprotein luciferase provides another useful tool for assaying for modulators of GPCR activity. Cells (*e.g.*, CHO cells or COS 7 cells) are transiently co-transfected with both a GPCR expression construct (*e.g.*, GPCR-encoding sequence in pzeoSV2 (Invitrogen, San Diego, CA)) and a reporter construct which includes a gene for the luciferase protein downstream from a transcription factor, either cAMP-response element (CRE), AP-1, or NF kappa B. Agonist binding to receptors coupled to the Gs subtype of G-proteins leads to increases in cAMP, activating the CRE transcription factor and resulting in expression of the luciferase gene. Agonist binding to receptors coupled to the Gq subtype of G-protein leads to production of diacylglycerol that activates protein kinase C. As a result, the AP-1 or NF kappa B transcription factors are activated which stimulate expression of the luciferase gene. Expression levels of luciferase reflect the activation status of the signaling events. [See generally George *et al.*, *Journal of Biomolecular Screening*,

- 123 -

2(4): 235-40 (1997); and Stratowa *et al.*, *Current Opinion in Biotechnology*, 6: 574-81 (1995).] Luciferase activity may be quantitatively measured using, *e.g.*, luciferase assay reagents that are commercially available from Promega (Madison, WI).

In one exemplary assay, CHO cells are plated in 24-well culture dishes at a density of 100,000 cells/well one day prior to transfection and cultured at 37°C in  $\alpha$ MEM (Gibco/BRL, Gaithersburg, MD) supplemented with 10% FBS, 2 mM glutamine, 10 U/ml penicillin and 10  $\mu$ g/ml streptomycin. Cells are transiently co-transfected with both a GPCR expression construct and a reporter construct containing the luciferase gene. The reporter plasmids CRE-luciferase, AP-1-luciferase and NF kappa B-luciferase may be purchased from Stratagene (LaJolla, CA).

Transfections are performed using FuGENE 6 transfection reagent (Boehringer-Mannheim), and the protocol provided in the product insert. Cells transfected with the reporter construct alone are used as a control. Twenty-four hours after transfection, cells are washed once with phosphate buffered saline (PBS) pre-warmed to 37°C. Serum-free  $\alpha$ MEM is then added to the cells either alone (control) or with one or more candidate modulators and the cells are incubated at 37°C for five hours. Thereafter, cells are washed once with ice cold PBS and lysed by the addition of 100  $\mu$ l of lysis buffer/well (from luciferase assay kit, Promega, Madison, WI). After incubation for 15 minutes at room temperature, 15  $\mu$ l of the lysate is mixed with 50  $\mu$ l substrate solution (Promega) in an opaque white 96-well plate, and the luminescence is read immediately on a Wallace model 1450 MicroBeta scintillation and luminescence counter (Wallace Instruments, Gaithersburg, MD).

Differences in luminescence in the presence versus the absence of a candidate modulator compound are indicative of modulatory activity. Receptors that are either constitutively active or activated by agonists give a 3-20 fold stimulation of luminescence compared to cells transfected with the reporter gene alone. Modulators that act as inverse agonists will reverse this effect.

#### **D. Intracellular Calcium Measurement using FLIPR**

Changes in intracellular calcium levels are another recognized indicator of G protein-coupled receptor activity, and such assays can be employed to

- 124 -

evaluate modulators of GPCR activity. For example, CHO cells stably transfected with a GPCR expression vector are plated at a density of  $4 \times 10^4$  cells/well in Packard black-walled 96-well plates specially designed to isolate fluorescent signal to individual wells. The cells are incubated for 60 minutes at 37°C in modified  
5 Dulbecco's PBS (D-PBS) containing 36 mg/L of pyruvate and 1 g/L of glucose with the addition of 1% FBS and one of four calcium indicator dyes (Fluo-3™ AM, Fluo-4™ AM, Calcium Green™-1 AM, or Oregon Green™ 488 BAPTA-1 AM) at a concentration of 4 µM. Plates are washed once with modified D-PBS without 1% FBS and incubated for 10 minutes at 37°C to remove residual dye from the cellular  
10 membrane. In addition, a series of washes with modified D-PBS without 1% FBS is performed immediately prior to activation of the calcium response.

Calcium response is initiated by the addition of one or more candidate receptor agonist compounds, calcium ionophore A23187 (10 µM), or ATP (4 µM). Fluorescence is measured by Molecular Device's FLIPR with an argon laser,  
15 excitation at 488 nm. [See, e.g., Kuntzweiler *et al.*, *Drug Development Research*, 44(1): 14-20 (1998).] The F-stop for the detector camera was set at 2.5 and the length of exposure was 0.4 milliseconds. Basal fluorescence of cells was measured for 20 seconds prior to addition of agonist, ATP, or A23187, and was subtracted from the response signal. The calcium signal is measured for approximately 200 seconds,  
20 taking readings every two seconds. Calcium ionophore and ATP increase the calcium signal 200% above baseline levels. In general, activated orphan GPCRs increase the calcium signal approximately 10-15% above baseline signal.

#### E. Mitogenesis Assay

25 In mitogenesis assays, the ability of candidate modulators to induce or inhibit GPCR-mediated cell growth is determined. [See, e.g., Lajiness *et al.*, *Journal of Pharmacology and Experimental Therapeutics*, 267(3): 1573-81 (1993).]

For example, CHO cells stably expressing a GPCR are seeded into 96-well plates at a density of 5000 cells/well and grown at 37°C in αMEM supplemented with 10% fetal calf serum. After 48 hours, the cells are rinsed twice with serum-free  
30 αMEM and 80 µl of fresh αMEM, or αMEM containing a known mitogen, is added

- 125 -

along with 20  $\mu$ l  $\alpha$ MEM containing varying concentrations of one or more test compounds diluted in serum free media. As controls, some wells on each plate receive serum-free media alone, and some receive media containing 10% FBS. Untransfected cells or cells transfected with vector alone also may serve as controls.

5                   After culture for 16-18 hours, 1  $\mu$ Ci/well of [ $^3$ H]-thymidine (2 Ci/mmol; cpm) is added to the wells and cells are incubated for an additional 2 hours at 37°C. The cells are trypsinized and harvested onto filter mats with a cell harvester (Tomtec) and the filters are counted in a Betaplate counter. The incorporation of  $^3$ H-thymidine in serum-free test wells is compared to the results achieved in cells  
10                   stimulated with serum. Use of multiple concentrations of test compounds permits creation and analysis of dose-response curves using the non-linear, least squares fit equation:  $A = B \times [C / (D + C)] + G$  where A is the percent of serum stimulation; B is the maximal effect minus baseline; C is the  $EC_{50}$ ; D is the concentration of the compound; and G is the maximal effect. Parameters B, C and G are determined by  
15                   Simplex optimization.

                  Agonists that bind to the receptor are expected to increase [ $^3$ H]-thymidine incorporation into cells, showing up to 80% of the response to serum. Antagonists that bind to the receptor will inhibit the stimulation seen with a known agonist by up to 100%.

20

#### F.     [ $^{35}$ S]GTP $\gamma$ S Binding Assay

                  Because G protein-coupled receptors signal through intracellular "G proteins" whose activity involves GTP/GDP binding and hydrolysis. Another indicator of GPCR modulator activity is measuring binding of the non-hydrolyzable  
25                   GTP analog [ $^{35}$ S]GTP $\gamma$ S in the presence and absence of putative modulators. [See, e.g., Kowal, *et al.*, *Neuropharmacology*, 37: 179-87 (1998).]

                  In one exemplary assay, cells stably transfected with a GPCR expression vector are grown in 10 cm dishes to subconfluence, rinsed once with 5 ml of ice cold  $Ca^{2+}$ /Mg $^{2+}$  free PBS, and scraped into 5 ml of the same buffer. Cells are  
30                   pelleted by centrifugation (500 x g, 5 minutes), resuspended in TEE buffer (25 mM Tris, 5 mM EDTA, 5 mM EGTA, pH 7.5) and frozen in liquid nitrogen. After

- 126 -

thawing, the cells are homogenized using a dounce (one ml TEE per plate of cells), and centrifuged at 1,000 x g for 5 minutes to remove nuclei and unbroken cells.

The homogenate supernatant is centrifuged at 20,000 x g for 20 minutes to isolate the membrane fraction. The membrane pellet is then washed once with TEE and resuspended in binding buffer (20 mM HEPES, pH 7.5, 150 mM NaCl, 10 mM MgCl<sub>2</sub>, 1 mM EDTA). The resuspended membranes can be frozen in liquid nitrogen and stored at -70°C until use.

Aliquots of cell membranes prepared as described above and stored at -70°C are thawed, homogenized, and diluted to a concentration of 10-50 µg/ml in buffer containing 20 mM HEPES, 10 mM MgCl<sub>2</sub>, 1 mM EDTA, 120 mM NaCl, 10 µM GDP, and 0.2 mM ascorbate. In a final volume of 90 µl, homogenates are incubated with varying concentrations of putative modulator compounds or 100 µM GTP for 30 minutes at 30°C and then placed on ice. To each sample, 10 µl guanosine 5'-O-(3[<sup>35</sup>S]thio) triphosphate (NEN, 1200 Ci/mmol), ([<sup>35</sup>S]-GTPγS), was added to a final concentration of 100-200 pM. Samples are incubated at 30°C for an additional 30 minutes. The reaction is then stopped by the addition of 1 ml of 10 mM HEPES, and 10 mM MgCl<sub>2</sub> (pH 7.4), at 4°C, and filtration.

Samples are filtered over Whatman GF/B filters. These filters are washed with 20 ml ice-cold 10 mM HEPES (pH 7.4) and 10 mM MgCl<sub>2</sub> and counted by liquid scintillation spectroscopy. Nonspecific binding of [<sup>35</sup>S]-GTPγS is measured in the presence of 100 µM GTP and subtracted from the total. Compounds are selected that modulate the amount of [<sup>35</sup>S]-GTPγS binding in the cells, compared to untransfected control cells. Activation of receptors by agonists gives up to a five-fold increase in [<sup>35</sup>S]GTPγS binding. This response is blocked by antagonists.

#### G. MAP Kinase Activity Assay

Evaluation of MAP Kinase activity in cells expressing a GPCR provide another assay to identify modulators of GPCR activity. [See, *e.g.*, Lajiness *et al.*, *Journal of Pharmacology and Experimental Therapeutics*, 267(3): 1573-81 (1993); and Boulton *et al.*, *Cell*, 65: 663-75 (1991).]

- 127 -

In one embodiment, CHO cells stably transfected with a GPCR-encoding polynucleotide are seeded into 6 well plates at a density of 70,000 cells/well 48 hours prior to the assay. During this time, the cells are cultured at 37°C in  $\alpha$ MEM media supplemented with 10% FBS, 2 mM glutamine, 10 U/ml penicillin and 10  $\mu$ g/ml streptomycin. The cells are serum starved for 1-2 hours prior to the addition of  
5 stimulants.

For the assay, the cells are treated with media alone or media containing a putative agonist or phorbol ester-myristoyl acetate (PMA) as a positive control. After treatment, cells are incubated at 37°C for varying times. To stop the  
10 reaction, the plates are placed on ice, the media is aspirated, and the cells are rinsed with 1 ml of ice-cold PBS containing 1 mM EDTA. Thereafter, 200  $\mu$ l cell lysis buffer (12.5 mM MOPS (pH 7.3), 12.5 mM  $\beta$ -glycerophosphate, 7.5 mM MgCl<sub>2</sub>, 0.5 mM EGTA, 0.5 mM sodium vanadate, 1 mM benzamidine, 1 mM dithiothreitol, 10  $\mu$ g/ml leupeptin, 10  $\mu$ g/ml aprotinin, 2  $\mu$ g/ml pepstatin A, and 1  $\mu$ M okadaic acid) is  
15 added to the cells. The cells are scraped from the plates and homogenized by 10 passages through a 23 3/4 gauge needle. The cytosol fraction is prepared by centrifugation at 20,000 x g for 15 minutes.

Aliquots (5-10  $\mu$ l containing 1-5  $\mu$ g protein) of cytosols are mixed with 1 mM MAPK Substrate Peptide (APRTPGGRR; SEQ ID NO: 25); Upstate  
20 Biotechnology, Inc., N.Y.) and 50  $\mu$ M [ $\gamma$ -<sup>32</sup>P]ATP, (NEN, 3000 Ci/mmol) diluted to a final specific activity of ~2000 cpm/pmol in a total volume of 25  $\mu$ l. The samples are incubated for 5 minutes at 30°C, and reactions are stopped by spotting 20  $\mu$ l on 2 cm<sup>2</sup> of Whatman P81 phosphocellulose paper. The filter squares are washed in 4 changes of 1% H<sub>3</sub>PO<sub>4</sub>, and the squares are counted by liquid scintillation spectroscopy.  
25 Equivalent cytosolic extracts are incubated without MAPK substrate peptide, and the cpm from these samples are subtracted from the matched samples with the substrate peptide. The cytosolic extract from each well is used as a separate point. Protein concentrations are determined by a dye binding protein assay (Bio-Rad). Agonist activation of the receptor is expected to result in up to a five fold increase in MAPK  
30 enzyme activity. This increase is blocked by antagonists.

- 128 -

#### H. <sup>3</sup>H]Arachidonic Acid Release

The activation of GPCR's also has been observed to potentiate arachidonic acid release in cells, providing yet another useful assay for modulators of the activity of GPCR's of the present invention. [See, e.g., Kanterman *et al.*,  
5 *Molecular Pharmacology*, 39: 364-9 (1991).] For example, CHO cells that are stably transfected with a GPCR expression vector are plated in 24-well plates at a density of 15,000 cells/well and grown in  $\alpha$ MEM media supplemented with 10% FBS, 2 mM glutamine, 10 U/ml penicillin and 10  $\mu$ g/ml streptomycin for 48 hours at 37°C before use. Cells of each well are labeled by incubation with [<sup>3</sup>H]arachidonic acid  
10 (Amersham Corp., 210 Ci/mmol) at 0.5  $\mu$ Ci/ml in 1 ml  $\alpha$ MEM supplemented with 10 mM HEPES (pH 7.5), and 0.5% fatty-acid-free bovine serum albumin for 2 hours at 37°C. The cells are then washed twice with 1 ml of the same buffer.

Candidate modulator compounds are added in 1 ml of the same buffer, either alone or containing 10  $\mu$ M ATP (Adenosine 5'-triphosphate) and the cells are  
15 incubated at 37°C for 30 minutes. Buffer alone and mock transfected cells are used as controls. Samples (0.5 ml) from each well are counted by liquid scintillation spectroscopy. Agonists which activate the receptor will lead to potentiation of the ATP-stimulated release of [<sup>3</sup>H]-arachidonic acid. This potentiation is blocked by antagonists.

#### I. Extracellular Acidification Rate

In yet another assay, the effects of putative modulators of GPCR activity are assayed by monitoring extracellular changes in pH induced by the putative modulators. [See, e.g., Dunlop *et al.*, *Journal of Pharmacological and Toxicological*  
25 *Methods*, 40(1): 47-55 (1998).]

CHO cells transfected with a GPCR expression vector are seeded into 12-mm capsule cups (Molecular Devices Corp.) at  $4 \times 10^5$  cells/cup in  $\alpha$ MEM supplemented with 10% FBS, 2 mM l-glutamine, 10 units/ml penicillin, and 10  $\mu$ g/ml streptomycin. The cells are incubated in this media at 37°C in 5% CO<sub>2</sub> for 24 hours.

30 Extracellular acidification rates are measured using a Cytosensor microphysiometer (Molecular Devices Corp.). The capsule cups are loaded into the

- 129 -

sensor chambers of the microphysiometer and the chambers are perfused with running buffer (bicarbonate free  $\alpha$ MEM supplemented with 4 mM L-glutamine, 10 units/ml penicillin, 10  $\mu$ g/ml streptomycin, 26 mM NaCl) at a flow rate of 100  $\mu$ l/min.

Agonists or other agents are diluted into the running buffer and perfused through a second fluid path. During each 60 second pump cycle, the pump is run for 38 seconds and is off for the remaining 22 seconds. The pH of the running buffer in the sensor chamber is recorded during the cycle from 43-58 seconds, and the pump is re-started at 60 seconds to start the next cycle. The rate of acidification of the running buffer during the recording time is calculated by the Cytosoft program. Changes in the rates of acidification are calculated by subtracting the baseline value (the average of 4 rate measurements immediately before addition of modulator candidates) from the highest rate measurement obtained after addition of a modulator candidate. The selected instrument detects 61 mV/pH unit. Modulators that act as agonists at the receptor result in an increase in the rate of extracellular acidification as compared to the rate in the absence of agonist. This response is blocked by modulators which act as antagonists at the receptor.

## EXAMPLE 7

### Luciferase Reporter Gene Assays

Luciferase reporter gene assays (essentially as described in Example 6) were carried out to measure signaling activity of the GPCR receptors when coupled to Gs, Gi or Gq G-proteins. Activation of Gs coupled receptors results in stimulation of intracellular cAMP production which leads to activation of the transcription factor cyclic AMP response element (CRE). Therefore activation of Gs coupled receptors can be detected by measuring transcription and translation of the reporter gene CRE-luciferase. The level of expression of the CRE reporter gene is dependent on the intracellular level of cAMP. Similarly, activation of Gs, Gi or Gq coupled receptors will result in activation of the AP-1 transcription factor. Expression of the AP-1 transcription factor can be attributed to changes in cAMP levels and/or increases in the levels of intracellular calcium and therefore can be an indication of G-protein coupled receptor activation.

- 130 -

CHO 10001A cells (Gottesman *et al.*, *Somatic Cell Genetics* 6: 45-61, 1980) were maintained in Minimal Essential Medium (MEM) supplemented with 10% FBS (Hyclone Laboratories, Inc., Logan, Utah) at 37°C in an atmosphere of 5% CO<sub>2</sub>. The cells were split 1:5 twice a week for maintenance. Plasmids used in the experiments were propagated in *E.coli* strain DH5 (Gibco BRL) and purified using the Qiagen Maxi-prep plasmid purification system according to the manufacturer's instructions.

One day prior to transfection,  $1 \times 10^5$  CHO cells/well were plated on 24 well culture plates and allowed to adhere overnight. Each well on the plate was transfected with 0.5 µg of either AP-1 luciferase (Stratagene, LaJolla, CA) or CRE luciferase plasmid alone or in combination with 0.125 µg of a GPCR plasmid (GPCR DNA inserted into the pCDNA3 vector from Invitrogen). Cells were transiently transfected with the commercially available transfection reagent FUGENE-6 according to the manufacturer's instructions (Boehringer Mannheim, Indianapolis, IN).

Twenty-four hours after transfection, the cells were washed in PBS pre-warmed to 37°C. Agonists and antagonists were diluted in pre-warmed serum-free MEM, added to the transfected cells and incubated at 37°C, 5% CO<sub>2</sub> for 5 hours. Subsequently, the cells were washed once in ice cold PBS and lysed with the addition of 100 µl of lysis buffer (Promega) to each well. After a 15 minute incubation at room temperature, luciferase reporter gene activation was analyzed with the Luciferase Assay Reagents commercially available from Promega (Madison, WI). An aliquot of lysate (15 µl) was mixed with 50 µl of substrate solution in an opaque white 96 well plate. The luminescence from the plate was read in a Wallance 1450 MicroBeta scintillation and luminescence counter (Wallac Instruments, Gaithersburg, MD). Constitutive GPCR activity was calculated as activity measured in GPCR transfected cells divided by activity measured in control cells (control cells= luciferase-transfected cells in the absence of GPCR plasmid). The measurements of GPCR constitutive activity (as a percentage of control measurements) are summarized in the table below:

- 131 -

	<u>GPCR</u>	<u>CRE Activity</u>	<u>AP-1 Activity</u>
	CON193	128%	100%
	CON197	165%	100%
	CON198	178%	146%
5	CON203	100%	468%
	CON215	173%	307%
	CON222	100%	100%
	CON202	135%	336%
	CON166	115%	100%
10	CON217	211%	100%

These results provide useful information for designing screening assays to identify molecules (natural or artificial) that activate or inhibit the GPCR's of the invention. For example, compound libraries can be screened using the AP-1 luciferase (for CON198, CON203, CON215, or CON202) or the CRE-luciferase assay (for CON193, CON197, CON198, CON215, CON202, and CON166) to identify compounds which increase the signaling activity in GPCR polypeptide expressing cells as compared to receptor negative cells. The identified compounds may be useful for predicting endogenous ligands for the GPCR polypeptides, for measuring the physiological effects of GPCR activation in animal models, and for designing therapeutics to modulate GPCR activity to treat disease states.

#### EXAMPLE 8

##### Chromosomal Localization of GPCR

The following example pertains to chromosomal localization of GPCR genes of the present invention (e.g., CON193, CON166, CON103, CON203, CON198, CON197, CON202, CON222, CON215, or CON217). The chromosomal localization permits use of the GPCR polynucleotide sequences (including fragments thereof) as chromosomal markers to assist with genome mapping and to provide markers for disease states. Chromosomal localization also permits correlation of the

- 132 -

GPCR's of the invention with disease states in which aberrant activity of the GPCR is implicated, especially disease states that have previously linked (or will be linked) with mutations, polymorphisms, chromosomal rearrangements, and other chromosomal changes near the locus of the GPCR gene.

5

**A. CON197**

Chomosomal localization of the gene encoding CON197 (SEQ ID NO: 11) was determined using the Standford G3 Radiation Hybrid Panel (Research Genetics, Inc. Huntsville, AL). This panel contains 83 radiation hybrid clones of the entire human genome as created by the Stanford Human Gemone Center (Stanford, California). PCR was carried out with each clone within the Hybrid Panel and the results were submitted to the Standford Human Genomic Center via e-mail for analysis (<http://www.shgc.stanford.edu/RH/rhserverformnew.html>).

PCR reactions were carried out with the Expand Hi-Fi PCR System™ according the manufacturer's instructions (Roche Molecular Biochemicals, Indianapolis, IN). Primers, synthesized by Genosys Corp. (The Woodlands, TX), were designed to generate a 10 base pair fragment of CON197-encoding DNA in the presence of the appropriate genomic DNA. The forward primer, denoted as LW1332 (TCCTACTGTCATGAACCC; SEQ ID NO: 74), corresponded to nuceotides 396 through 413 of SEQ ID NO: 11. The reverse primer, denoted as LW1333 (CAGAAGAAGTTGTCCAGC; SEQ ID NO: 75), corresponded to the complement of nucleotides 519 through 536 of SEQ ID NO: 11. Each reaction contained 25 ng of DNA from a hybrid clone, 60 ng of Primer LW1332, and 60 ng of Primer LW1333 resulting in a final volume of 15 µl. The PCR reactions were carried our in a GeneAmp 9700 PCR thermocycler (Perkin Elmer Applied Biosystems) under the following conditions: 94°C for 3 minutes followed by 35 cycles of 94°C for 30 seconds, 52°C for 1 minute, and 72°C for 2 minutes. The PCR reactions were then analyzed on a 2.0% agarose gel and stained with ethidium bromide. The lanes were scored for the presence of the 140 base pair PCR product.

The G3 Hybrid Panal analysis revealed that the CON197 gene (SEQ ID NO: 11) was localized to chromosome 14, most nearly linked to Standford marker

30

- 133 -

SHGC-10764 with a LOD score of 9.10. The SHGC-10764 marker lies at position 1q11.1.

**B. CON202**

5                   Chomosomal localization of the gene encoding CON202 (SEQ ID NO: 13) was determined using the Stanford G3 Radiation Hybrid Panel (Research Genetics, Inc. Huntsville, AL). This panel contains 83 radiation hybrid clones of the entire human genome as created by the Stanford Human Genome Center (Stanford, California). PCR was carried out with each clone within the Hybrid Panel and the results were submitted to the Stanford Human Genomic Center via e-mail for analysis (<http://www.shgc.stanford.edu/RH/rhserverformnew.html>).

                  PCR reactions were carried out with the Expand Hi-Fi PCR System™ according to the manufacturer's instructions (Roche Molecular Biochemicals, Indianapolis, IN). Primers, synthesized by Genosys Corp. (The Woodlands, TX), were designed to generate a 250 base pair fragment of CON202-encoding DNA in the presence of the appropriate genomic DNA. The forward primer, denoted as LW1480 (GGTTCTACCTGGACTTATGG; SEQ ID NO: 70), corresponded to nucleotides 515 through 534 of SEQ ID NO: 13. The reverse primer, denoted as LW1481 (TAATGAATGAGTAAGTGCCC; SEQ ID NO: 71), corresponded to the complement of nucleotides 745 through 764 of SEQ ID NO: 13. Each reaction contained 25 ng of DNA from a hybrid clone, 60 ng of Primer LW1480, and 60 ng of Primer LW1481 resulting in a final volume of 15 µl. The PCR reactions were carried out in a GeneAmp 9700 PCR thermocycler (Perkin Elmer Applied Biosystems) under the following conditions: 94°C for 3 minutes followed by 35 cycles of 94°C for 30 seconds, 52°C for 1 minute, and 72°C for 2 minutes. The PCR reactions were then analyzed on a 2.0% agarose gel and stained with ethidium bromide. The lanes were scored for the presence of the 250 base pair PCR product.

                  The G3 Hybrid Panel analysis revealed that the CON202 gene (SEQ ID NO: 13) was localized to chromosome 7, most nearly linked to Stanford marker SHGC-12021 with a LOD score of 10.36. The SHGC-12021 marker lies at position 7q21. There is evidence that schizophrenia is linked to chromosome 7q22, and

- 134 -

therefor any genes localized to this region are candidates for disease involvement or susceptibility. [See Ekelund *et al.*, *Human Mol. Genetics* 9(7): 1049-1057 (2000); Faraone *et al.*, *Am. J. Med. Genet.* 81: 290-295 (September, 1998); and Blouin *et al.*, *Nat. Genet.* 20: 70-73 (1998)]. The SHGC-12021 marker is proximal to 7q22 (~1 cM) and therefore may be associated with schizophrenia susceptibility.

In particular, G protein-coupled receptors, such as CON202 polypeptide, have the biochemical and functional potential to play a role in the disease process of schizophrenia. CON202 is an attractive target for screening for ligands (natural and synthetic) that are useful in modulating cellular processes involved in schizophrenia. In addition, the chromosomal localization data (especially coupled with CON202 expression patterns in the brain) identifies CON202 as a candidate for screening healthy and affected (schizophrenia) individuals for CON202 allelic variants, mutations, duplications, rearrangements, and other chromosomal variations that correlate with the disease state. Variations that correlate with disease state are useful for diagnosis of disease or disease susceptibility. CON202 constructs containing the variations are useful for designing targeted therapeutics for treatment of the disease (e.g., by using the assays for modulators described in preceding examples.

### C. High throughput Analysis

The EMBL High Throughput Genome database (provided by the European Bioinformatics Institute) was searched with GPCR nucleotide sequences to determine chromosomal localization for CON193, CON166, CON103, CON203, CON198, and CON215 genes. The results are summarized in the table below:

- 135 -

	<u>GPCR</u>	<u>SEQ ID NO:</u>	<u>Chromosome Localization</u>	<u>Based on Genbank Accession No.</u>
	CON193	1	11	AC026090
	CON166	3	X	AC021992
	CON103	5	2	AC013396
5	CON203	7	3	AC024886
	CON198	9	11	AC025249
	CON215	17	3	AC024886

While the present invention has been described in terms of specific  
 10 embodiments, it is understood that variations and modifications will occur to those in  
 the art, all of which are intended as aspects of the present invention. Accordingly,  
 only such limitations as appear in the claims should be placed on the invention.

#### Summary of Sequences:

15	<u>SEQ ID NO.</u>	<u>Description</u>
	1	CON 193 DNA
	2	CON 193 protein
	3	CON 166 DNA
	4	CON 166 protein
20	5	CON 103 DNA
	6	CON 103 protein
	7	CON 203 DNA
	8	CON 203 protein
	9	CON 198 DNA
25	10	CON 198 protein
	11	CON 197 DNA
	12	CON 197 protein
	13	CON 202 DNA
	14	CON 202 protein
30	15	CON 222 DNA
	16	CON 222 protein
	17	CON 215 DNA

- 136 -

	<u>SEQ ID NO.</u>	<u>Description</u>
	18	CON 215 protein
	19	CON 217 DNA
	20	CON 217 protein
	21	PCR primer LW 1282 for CON 193
5	22	PCR primer LW 1283 for CON 193
	23	PCR primer LW 1372 for CON 193
	24	PCR primer LW 1374 for CON 193
	25	MAPK Substrate Peptide
	26	Primer LW 1248 for CON 193 to generate insitu hybridization probe
10	27	Primer LW 1249 for CON 193 to generate insitu hybridization probe
	28	PCR primer LW 1278 for CON 166
	29	PCR primer LW 1279 for CON 166
	30	PCR primer LW 1405 for CON 166
	31	PCR primer LW 1406 for CON 166
15	32	PCR primer LW 1280 for CON 103
	33	PCR primer LW 1281 for CON 103
	34	PCR primer LW 1385 for CON 103
	35	PCR primer LW 1386 for CON 103
	36	PCR primer LW 1329 for CON 203
20	37	PCR primer LW 1377 for CON 203
	38	PCR primer LW 1387 for CON 203
	39	PCR primer LW 1388 for CON 203
	40	Primer LW 1314 for CON 203 to generate insitu hybridization probe
	41	Primer LW 1315 for CON 203 to generate insitu hybridization probe
25	42	PCR primer LW 1326 for CON 198
	43	PCR primer LW 1327 for CON 198
	44	PCR primer LW 1415 for CON 198
	45	PCR primer LW 1416 for CON 198
	46	Primer LW 1308 for CON 198 to generate insitu hybridization probe
30	47	Primer LW 1309 for CON 198 to generate insitu hybridization probe
	48	PCR primer LW 1324 for CON 197
	49	PCR primer LW 1325 for CON 197
	50	Primer LW 1306 for CON 197 to generate insitu hybridization probe
	51	Primer LW 1307 for CON 197 to generate insitu hybridization probe

- 137 -

	<u>SEQ ID NO.</u>	<u>Description</u>
	52	PCR primer GV 599 for CON 202
	53	PCR primer GV 600 for CON 202
	54	PCR primer LW 1482 for CON 202
	55	PCR primer LW 148 for CON 202
5	56	Primer LW 1310 for CON 202 to generate insitu hybridization probe
	57	Primer LW 1311 for CON 202 to generate insitu hybridization probe
	58	PCR primer LW 1442 for CON 222
	59	PCR primer LW 1443 for CON 222
	60	PCR primer LW 1440 for CON 222
10	61	PCR primer LW 1441 for CON 222
	62	Primer LW 1472 for CON 222 to generate insitu hybridization probe
	63	Primer LW 1473 for CON 222 to generate insitu hybridization probe
	64	Primer LW 1411 for CON 215 to generate insitu hybridization probe
	65	Primer LW 1412 for CON 215 to generate insitu hybridization probe
15	66	PCR primer LW 1448 for CON 217
	67	PCR primer LW 1449 for CON 217
	68	Primer LW 217A for CON 217 to generate insitu hybridization probe
	69	Primer LW 218B for CON 217 to generate insitu hybridization probe
	70	Primer LW 1480 for CON 202 chromosomal localization
20	71	Primer LW 1481 for CON 202 chromosomal localization
	72	Primer CON103a for CON 103 to generate insitu hybridization probe
	73	Primer CON103b for CON 103 to generate insitu hybridization probe
	74	Primer LW 1332 for CON 197 chromosomal localization
	75	Primer LW 1333 for CON 197 chromosomal localization

25


30

Applicant's or agent's file reference number 28341/uz ruP	International application No. To Be Determined
--	---

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>90; 98</u> , line <u>10-14; 15</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S. A.	
Date of deposit 18 January 2000	Accession Number B-30250
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
When designating Australia, in accordance with regulation 3.25 of the Patents Regulations (Australia Statutory Rules 1991 No. 71), samples of materials deposited in accordance with the Budapest Treaty in relation to this Patent Request are only to be provided before: the patent is granted on the application; or the application has lapsed or been withdrawn or refused; to a person who is: a skilled addressee without an interest in the invention; and nominated by a person who makes a request for the furnishing of those samples.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 


For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number 28341/6276P	International application No To Be <del>Deposited</del>
---	--

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>90; 98</u> , line <u>10-14; 15</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S.A.	
Date of deposit 18 January 2000	Accession Number B-30250
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
In respect of those designations in which a European patent or a patent in Norway is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or the corresponding information concerning the patent in Norway or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC and the corresponding regulations in Norway).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<input checked="" type="checkbox"/> For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 


<input type="checkbox"/> For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number	28341/bZ/bP	International application No. To Be Determined
---	-------------	---

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91; 98</u> , line <u>10-14; 16</u>	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S. A.	
Date of deposit 18 January 2000	Accession Number B-30248
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
When designating Australia, in accordance with regulation 3.25 of the Patents Regulations (Australia Statutory Rules 1991 No. 71), samples of materials deposited in accordance with the Budapest Treaty in relation to this Patent Request are only to be provided before: the patent is granted on the application; or the application has lapsed or been withdrawn or refused; to a person who is: a skilled addressee without an interest in the invention; and nominated by a person who makes a request for the furnishing of those samples.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 


For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number	28341/bZ/bP	International application No. To Be Determined
--	-------------	--

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91; 98</u> , line <u>10-14; 16</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S.A.	
Date of deposit 18 January 2000	Accession Number B-30248
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
In respect of those designations in which a European patent or a patent in Norway is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or the corresponding information concerning the patent in Norway or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC and the corresponding regulations in Norway).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 


For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number 28341.02101	International application No. To Be Determined
---	---

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>92; 98</u> , line <u>10-14; 17</u>	
B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input checked="" type="checkbox"/></span>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S. A.	
Date of deposit 18 January 2000	Accession Number B-30247
C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input checked="" type="checkbox"/></span>	
When designating Australia, in accordance with regulation 3.25 of the Patents Regulations (Australia Statutory Rules 1991 No. 71), samples of materials deposited in accordance with the Budapest Treaty in relation to this Patent Request are only to be provided before: the patent is granted on the application; or the application has lapsed or been withdrawn or refused; to a person who is: a skilled addressee without an interest in the invention; and nominated by a person who makes a request for the furnishing of those samples.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number 28341/0210P	International application N° To Be Date:
---	---

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>92; 98</u> , line <u>10-14; 17</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S.A.	
Date of deposit 18 January 2000	Accession Number B-30247
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
In respect of those designations in which a European patent or a patent in Norway is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or the corresponding information concerning the patent in Norway or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC and the corresponding regulations in Norway).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer <i>Shaul Lipk</i>


For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number	28341/bz/bp	International application N To Be Determined
---	-------------	---

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>93; 98</u> , line <u>10-14; 18</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S. A.	
Date of deposit 18 January 2000	Accession Number B-30254
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
When designating Australia, in accordance with regulation 3.25 of the Patents Regulations (Australia Statutory Rules 1991 No. 71), samples of materials deposited in accordance with the Budapest Treaty in relation to this Patent Request are only to be provided before: the patent is granted on the application; or the application has lapsed or been withdrawn or refused; to a person who is: a skilled addressee without an interest in the invention; and nominated by a person who makes a request for the furnishing of those samples.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	


<input checked="" type="checkbox"/> For receiving Office use only This sheet was received with the international application Authorized officer 	<input type="checkbox"/> For International Bureau use only This sheet was received by the International Bureau on: Authorized officer
--	---

Applicant's or agent's file reference number 28341/6276P	International application No. To Be Determined
---	---

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>93; 98</u> , line <u>10-14; 18</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S.A.	
Date of deposit 18 January 2000	Accession Number B-30254
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
In respect of those designations in which a European patent or a patent in Norway is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or the corresponding information concerning the patent in Norway or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC and the corresponding regulations in Norway).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	


<input checked="" type="checkbox"/> For receiving Office use only <input checked="" type="checkbox"/> This sheet was received with the international application  Authorized officer 	<input type="checkbox"/> For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on:  Authorized officer
--	--

Applicant's or agent's file reference number	28341/62/6P	International application I To Be I
---	-------------	--

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>94; 98</u> , line <u>3-7; 19</u>	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S. A.	
Date of deposit 18 January 2000	Accession Number B-30252
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
When designating Australia, in accordance with regulation 3.25 of the Patents Regulations (Australia Statutory Rules 1991 No. 71), samples of materials deposited in accordance with the Budapest Treaty in relation to this Patent Request are only to be provided before: the patent is granted on the application; or the application has lapsed or been withdrawn or refused; to a person who is: a skilled addressee without an interest in the invention; and nominated by a person who makes a request for the furnishing of those samples.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	


<input checked="" type="checkbox"/> For receiving Office use only <input checked="" type="checkbox"/> This sheet was received with the international application  Authorized officer 	<input type="checkbox"/> For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on:  Authorized officer
--	--

Applicant's or agent's file reference number 28341/62/61	International application No. To Be Determined
---	---

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>94; 98</u> , line <u>3-7; 19</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S.A.	
Date of deposit 18 January 2000	Accession Number B-30252
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
In respect of those designations in which a European patent or a patent in Norway is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or the corresponding information concerning the patent in Norway or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC and the corresponding regulations in Norway).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 


For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number 283416Z/bP	International application No. To E
--	---------------------------------------

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>94; 98</u> , line <u>27-31; 20</u>	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S. A.	
Date of deposit 18 January 2000	Accession Number B-30251
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
When designating Australia, in accordance with regulation 3.25 of the Patents Regulations (Australia Statutory Rules 1991 No. 71), samples of materials deposited in accordance with the Budapest Treaty in relation to this Patent Request are only to be provided before: the patent is granted on the application; or the application has lapsed or been withdrawn or refused; to a person who is: a skilled addressee without an interest in the invention; and nominated by a person who makes a request for the furnishing of those samples.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 


For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number	28341/b2/bP	International Application No. To Be Del
---	-------------	--

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13b/s)

A. The indications made below relate to the microorganism referred to in the description on page <u>94; 98</u> , line <u>27-31; 20</u>	
B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input checked="" type="checkbox"/></span>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S.A.	
Date of deposit 18 January 2000	Accession Number B-30251
C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input checked="" type="checkbox"/></span>	
In respect of those designations in which a European patent or a patent in Norway is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or the corresponding information concerning the patent in Norway or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC and the corresponding regulations in Norway).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 


For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number 2834	International application To Be Determined
--	---

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>95; 98</u> , line <u>19-23; 21</u>	
B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input checked="" type="checkbox"/></span>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S. A.	
Date of deposit 18 January 2000	Accession Number B-30253
C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input checked="" type="checkbox"/></span>	
When designating Australia, in accordance with regulation 3.25 of the Patents Regulations (Australia Statutory Rules 1991 No. 71), samples of materials deposited in accordance with the Budapest Treaty in relation to this Patent Request are only to be provided before: the patent is granted on the application; or the application has lapsed or been withdrawn or refused; to a person who is: a skilled addressee without an interest in the invention; and nominated by a person who makes a request for the furnishing of those samples.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<input checked="" type="checkbox"/> For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 


<input type="checkbox"/> For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number 28341.02.00	International a To Be Determined
---	-------------------------------------

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>95; 98</u> , line <u>19-23; 21</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S.A.	
Date of deposit 18 January 2000	Accession Number B-30253
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
In respect of those designations in which a European patent or a patent in Norway is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or the corresponding information concerning the patent in Norway or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC and the corresponding regulations in Norway).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	


For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer 	Authorized officer

Applicant's or agent's file reference number	28341/62/bP	International application No. To Be Determined
---	-------------	---

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>96; 98</u> , line <u>11-15; 22</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S. A.	
Date of deposit 18 January 2000	Accession Number B-30257
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
When designating Australia, in accordance with regulation 3.25 of the Patents Regulations (Australia Statutory Rules 1991 No. 71), samples of materials deposited in accordance with the Budapest Treaty in relation to this Patent Request are only to be provided before: the patent is granted on the application; or the application has lapsed or been withdrawn or refused; to a person who is: a skilled addressee without an interest in the invention; and nominated by a person who makes a request for the furnishing of those samples.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<input checked="" type="checkbox"/> For receiving Office use only <input checked="" type="checkbox"/> This sheet was received with the international application  Authorized officer 	<input type="checkbox"/> For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on:  Authorized officer
--	--

Applicant's or agent's file reference number 28341102100P	Information To Be De
--	-------------------------

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>96; 98</u> , line <u>11-15; 22</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S.A.	
Date of deposit 18 January 2000	Accession Number B-30257
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
In respect of those designations in which a European patent or a patent in Norway is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or the corresponding information concerning the patent in Norway or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC and the corresponding regulations in Norway).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer <i>Heidi Lih</i>


For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number 28341/02101	International application No. To Be Determined
---	---

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>97; 98</u> , line <u>4-8; 23</u>	
B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input checked="" type="checkbox"/></span>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S. A.	
Date of deposit 18 January 2000	Accession Number B-30255
C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input checked="" type="checkbox"/></span>	
When designating Australia, in accordance with regulation 3.25 of the Patents Regulations (Australia Statutory Rules 1991 No. 71), samples of materials deposited in accordance with the Budapest Treaty in relation to this Patent Request are only to be provided before: the patent is granted on the application; or the application has lapsed or been withdrawn or refused; to a person who is: a skilled addressee without an interest in the invention; and nominated by a person who makes a request for the furnishing of those samples.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	


<p>For receiving Office use only</p> <p><input checked="" type="checkbox"/> This sheet was received with the international application</p> <p>Authorized officer </p>	<p>For International Bureau use only</p> <p><input type="checkbox"/> This sheet was received by the International Bureau on:</p> <p>Authorized officer</p>
--	--

Applicant's or agent's file reference number 28341/6276P	International: To Be Det
---	-----------------------------

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>97; 98</u> , line <u>4-8; 23</u>	
B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input checked="" type="checkbox"/></span>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S.A.	
Date of deposit 18 January 2000	Accession Number B-30255
C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input checked="" type="checkbox"/></span>	
In respect of those designations in which a European patent or a patent in Norway is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or the corresponding information concerning the patent in Norway or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC and the corresponding regulations in Norway).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<input checked="" type="checkbox"/> For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 


<input type="checkbox"/> For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number	2834102101	International apj To Be Determined
---	------------	---------------------------------------

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>98</u> , line <u>1-3; 24</u>	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S. A.	
Date of deposit 18 January 2000	Accession Number B-30256
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
When designating Australia, in accordance with regulation 3.25 of the Patents Regulations (Australia Statutory Rules 1991 No. 71), samples of materials deposited in accordance with the Budapest Treaty in relation to this Patent Request are only to be provided before: the patent is granted on the application; or the application has lapsed or been withdrawn or refused; to a person who is: a skilled addressee without an interest in the invention; and nominated by a person who makes a request for the furnishing of those samples.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	


For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer 	Authorized officer

Applicant's or agent's file reference number	28341, 02, 01	International a To Be Determined
---	---------------	-------------------------------------

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>98</u> line <u>1-3; 24</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution Agricultural Research Service Culture Collection	
Address of depositary institution (including postal code and country) National Center for Agricultural Utilization Research Agricultural Research Service, U.S. Department of Agriculture 1815 North University Street, Peoria, Illinois 61604 U.S.A.	
Date of deposit 18 January 2000	Accession Number B-30256
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
In respect of those designations in which a European patent or a patent in Norway is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or the corresponding information concerning the patent in Norway or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC and the corresponding regulations in Norway).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

## CLAIMS

What is claimed is:

1. A purified and isolated seven transmembrane receptor polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence set forth in any one of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18 or 20, or a fragment thereof comprising an epitope specific to said seven transmembrane receptor polypeptide.  
5
2. A purified and isolated seven transmembrane receptor polypeptide according to claim 1 comprising an amino acid sequence at least 90% identical to the amino acid sequence set forth in SEQ ID NO: 2, or a fragment thereof comprising an epitope specific to said seven transmembrane receptor polypeptide.  
10
3. A purified and isolated seven transmembrane receptor polypeptide according to claim 1 comprising an amino acid sequence at least 90% identical to the amino acid sequence set forth in SEQ ID NO: 4, or a fragment thereof comprising an epitope specific to said seven transmembrane receptor polypeptide.  
15
4. A purified and isolated seven transmembrane receptor polypeptide according to claim 1 comprising an amino acid sequence at least 90% identical to the amino acid sequence set forth in SEQ ID NO: 6, or a fragment thereof comprising an epitope specific to said seven transmembrane receptor polypeptide.  
20
5. A purified and isolated seven transmembrane receptor polypeptide according to claim 1 comprising an amino acid sequence at least 90% identical to the amino acid sequence set forth in SEQ ID NO: 8, or a fragment thereof comprising an epitope specific to said seven transmembrane receptor polypeptide.  
25

6. A purified and isolated seven transmembrane receptor polypeptide according to claim 1 comprising an amino acid sequence at least 90% identical to the amino acid sequence set forth in SEQ ID NO: 10, or a fragment thereof comprising an epitope specific to said seven transmembrane receptor polypeptide.

5

7. A purified and isolated seven transmembrane receptor polypeptide according to claim 1 comprising an amino acid sequence at least 90% identical to the amino acid sequence set forth in SEQ ID NO: 12, or a fragment thereof comprising an epitope specific to said seven transmembrane receptor polypeptide.

10

8. A purified and isolated seven transmembrane receptor polypeptide according to claim 1 comprising an amino acid sequence at least 90% identical to the amino acid sequence set forth in SEQ ID NO: 14, or a fragment thereof comprising an epitope specific to said seven transmembrane receptor polypeptide.

15

9. A purified and isolated seven transmembrane receptor polypeptide according to claim 1 comprising an amino acid sequence at least 90% identical to the amino acid sequence set forth in SEQ ID NO: 16, or a fragment thereof comprising an epitope specific to said seven transmembrane receptor polypeptide.

20

10. A purified and isolated seven transmembrane receptor polypeptide according to claim 1 comprising an amino acid sequence at least 90% identical to the amino acid sequence set forth in SEQ ID NO: 18, or a fragment thereof comprising an epitope specific to said seven transmembrane receptor polypeptide.

25

11. A purified and isolated seven transmembrane receptor polypeptide according to claim 1 comprising an amino acid sequence at least 90% identical to the amino acid sequence set forth in SEQ ID NO: 20, or a fragment thereof comprising an epitope specific to said seven transmembrane receptor polypeptide.

30

12. A purified and isolated seven transmembrane receptor polypeptide according to any one of claims 1-11.

5 13. A purified and isolated polypeptide according to any one of claims 1-11 comprising at least one extracellular domain of the seven transmembrane receptor polypeptide.

10 14. A purified and isolated polypeptide according to any one of claims 1-11 comprising the N-terminal extracellular domain of the seven transmembrane receptor polypeptide.

15 15. A purified and isolated polypeptide according to any one of claims 1-11 comprising a seven transmembrane receptor fragment selected from the group consisting of an N-terminal extracellular domain transmembrane domains, extracellular loops connecting transmembrane domains, intracellular loops connecting transmembrane domains, a C-terminal cytoplasmic domain, and fusions thereof.

20 16. A polypeptide according to any one of claims 1-15, wherein the polypeptide further includes a heterologous tag amino acid sequence.

17. A purified and isolated polynucleotide comprising a nucleotide sequence that encodes the polypeptide of claim 16.

25 18. A purified and isolated polynucleotide comprising a nucleotide sequence that encodes a polypeptide according to any one of claims 2, 3, 4, 8 or 9.

30 19. A purified and isolated polynucleotide comprising a heterologous expression control sequence operatively linked to a nucleotide sequence that encodes a polypeptide according to any one of claims 1-16.

20. The polynucleotide according to claim 19, wherein the expression control sequence is a promoter sequence that promotes expression of said polynucleotide in an eukaryotic cell.

5                   21. The polynucleotide according to claim 19, wherein the promoter is a heterologous promoter that promotes expression of the polynucleotide in a human cell.

22. A purified and isolated polynucleotide comprising a nucleotide  
10                   sequence that encodes a mammalian seven transmembrane receptor, wherein said polynucleotide hybridizes to any one of the nucleotide sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, or 19 or the non-coding strand complementary thereto, under the following hybridization conditions:

                  (a) hybridization for 16 hours at 42°C in a hybridization solution  
15                   comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate and  
                  (b) washing 2 times for 30 minutes at 60°C in a wash solution comprising 0.1x SSC and 1% SDS,  
with the proviso that the nucleotide sequence of the polynucleotide differs from the coding sequence set forth in any one of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, or  
20                   19 and from its complementary strand by at least one nucleotide.

23. A polynucleotide according to claim 22 that encodes a human seven transmembrane receptor.

25                   24. A vector comprising a polynucleotide according to any one of claims 17-23.

25. A vector according to claim 24 that is an expression vector for expressing the polynucleotide in a mammalian cell.

30

26. A host cell stably transformed or transfected with a polynucleotide according to any one of claims 17-23 in a manner allowing the expression in said host cell of the polypeptide or fragment thereof encoded by the polynucleotide.

5                   27. A host cell stably transformed or transfected with a vector according to claim 24 or 25 in a manner allowing the expression in said host cell of the polypeptide or fragment thereof encoded by the polynucleotide.

10                   28. A method for producing a seven transmembrane receptor polypeptide comprising the steps of growing a host cell according to claim 26 or 27 in a nutrient medium under conditions in which the host cell expresses a seven transmembrane receptor encoded by the polynucleotide.

15                   29. A method according to claim 28, further comprising a step of isolating said polypeptide from said cell or said medium.

20                   30. A method according to claim 29, further comprising a step of isolating cell membranes from the host cell, wherein the cell membrane comprises the seven transmembrane receptor.

                    31. An antibody specific for a polypeptide according to any one of claims 1-15.

25                   32. The antibody of claim 31 which is a monoclonal antibody.

                    33. A hybridoma that produces an antibody according to claim 32.

                    34. An antibody according to claim 31 that is a humanized antibody.

35. An antibody according to claim 31 that specifically binds an extracellular epitope of a seven transmembrane receptor having an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18 or 20.

5

36. An antibody according to claim 35 that specifically binds to the amino-terminal extracellular domain of the seven transmembrane receptors.

10

37. A cell-free composition comprising polyclonal antibodies, wherein at least one of said antibodies is an antibody according to claim 31.

38. An anti-idiotypic antibody specific for an antibody according to claim 31.

15

39. A polypeptide comprising a fragment of an antibody according to claim 31, wherein said fragment and said polypeptide specifically bind to a seven transmembrane receptor having an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18 or 20.

20

40. A polypeptide according to claim 39 that is selected from the group consisting of single chain antibodies and CDR-grafted antibodies.

41. A composition comprising a polypeptide according to any one of claims 1-16 in a pharmaceutically acceptable carrier.

25

42. A composition comprising an antibody according to any one of claims 31, 32, 34, 35, or 36, or a polypeptide according to claim 39 or 40, in a pharmaceutically acceptable carrier.

43. A method for modulating ligand binding of a seven  
transmembrane receptor polypeptide according to any one of claims 1-15, comprising  
the step of contacting said seven transmembrane receptor polypeptide with an  
antibody specific for said seven transmembrane receptor, under conditions wherein  
5 the antibody binds the receptor.

44. A method for modulating ligand binding of a seven  
transmembrane receptor polypeptide comprising the step of contacting said seven  
transmembrane receptor polypeptide with a polypeptide according to claim 39 or 40.  
10

45. An assay to identify compounds that bind a seven transmembrane  
receptor polypeptide, said assay comprising the steps of:

(a) contacting a composition comprising a seven transmembrane  
receptor polypeptide according to any of claims 1-15 with a compound suspected of  
15 binding the seven transmembrane receptor polypeptide; and

(b) measuring binding between the compound and the seven  
transmembrane receptor polypeptide.

46. A method for identifying a modulator of binding between a seven  
transmembrane receptor polypeptide and a binding partner of the seven  
transmembrane receptor polypeptide, comprising the steps of:  
20

(a) contacting the binding partner and a composition comprising  
the seven transmembrane receptor polypeptide in the presence and in the absence of a  
putative modulator compound, where the seven transmembrane receptor polypeptide  
25 is a polypeptide according to any one of claims 1-15;

(b) measuring binding between the binding partner and said seven  
transmembrane receptor polypeptide; and

(c) identifying a putative modulator compound in view of  
decreased or increased binding between the binding partner and seven transmembrane  
receptor polypeptide in the presence of the putative modulator, as compared to  
30 binding in the absence of the putative modulator.

47. An assay according to claim 45 or 46 wherein the composition comprises a cell expressing the seven transmembrane receptor polypeptide on its surface.

5                   48. An assay according to claim 47 wherein the measuring step comprises measuring intracellular signaling of the seven transmembrane receptor polypeptide induced by the compound.

10                   49. A method for treating a neurological disorder comprising the step of administering to a mammal in need of such treatment a pharmaceutical composition comprising a compound in an amount effective to modulate biological activity of a seven transmembrane receptor in neurons of said mammal, wherein the compound is selected from the group consisting of:

- 15                   (a) an antibody according to any one of claims 31, 32, 34, 35, or 36;  
                    (b) an anti-idiotypic antibody according to claim 38;  
                    (c) a polypeptide according to claim 39 or 40;  
                    (d) a compound identified according to the method of claim 45; and  
                    (e) a modulator identified according to claim 46.

20                   50. The method of claim 49 wherein the neurological disorder is schizophrenia.

25                   51. A method according to claim 50, wherein the seven transmembrane receptor comprises a polypeptide according to claim 8.

30                   52. A method of treating schizophrenia comprising the step of administering to a human diagnosed with schizophrenia an amount of a modulator of CON202 receptor activity sufficient to modulate CON202 receptor activity or CON202 ligand binding in said human.

53. A method of diagnosing schizophrenia or a susceptibility to schizophrenia comprising the steps of:

(a) measuring the presence or amount of expression or activity of a polypeptide according to claim 8 in a cell of a human patient; and

5 (b) comparing the measurement of step (a) to a measurement of expression or activity of the polypeptide in a cell from a normal subject or the patient at an earlier time, wherein the diagnosis of schizophrenia or susceptibility to schizophrenia is based on the presence or amount of CON202 polypeptide expression or activity.

10 54. A method of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor, comprising the steps of:

(a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering the amino acid sequence, expression, or biological activity of at least one seven transmembrane receptor that is expressed in the brain,  
15 wherein the seven transmembrane receptor comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, and 20, or an allelic variant thereof, and wherein the nucleic acid corresponds to the gene encoding the seven transmembrane receptor; and

(b) diagnosing the disorder or predisposition from the presence or absence of  
20 said mutation, wherein the presence of a mutation altering the amino acid sequence, expression, or biological activity of allele in the nucleic acid correlates with an increased risk of developing the disorder.

55. A method according to claim 54, wherein the seven  
25 transmembrane receptor is CON202 comprising an amino acid sequence set forth in SEQ ID NO: 14, or an allelic variant thereof.

56. A method according to claim 55, wherein the disease is schizophrenia.

30

57. A method according to claim 56, wherein the assaying step comprises at least one procedure selected from the group consisting of:

(a) determining a nucleotide sequence of at least one codon of at least one CON202 allele of the human subject;

5 (b) performing a hybridization assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences;

(c) performing a polynucleotide migration assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different  
10 from one or more reference sequences; and

(d) performing a restriction endonuclease digestion to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences.

15 58. A method according to claim 56 wherein the assaying step comprises: performing a polymerase chain reaction (PCR) to amplify nucleic acid comprising CON202 coding sequence, and determining nucleotide sequence of the amplified nucleic acid.

20 59. A method of screening for a CON202 hereditary schizophrenia genotype in a human patient, comprising the steps of:

(a) providing a biological sample comprising nucleic acid from said patient, said nucleic acid including sequences corresponding to said patient's CON202 alleles;

25 (b) analyzing said nucleic acid for the presence of a mutation or mutations;

(c) determining a CON202 genotype from said analyzing step; and

(d) correlating the presence of a mutation in a CON202 allele with a hereditary schizophrenia genotype.

30

60. The method according to claim 59 wherein said biological sample is a cell sample.

61. The method according to claim 59 wherein said analyzing comprises sequencing a portion of said nucleic acid, said portion comprising at least one codon of said CON202 alleles.

62. The method according to claim 59 wherein said nucleic acid is DNA.

63. The method according to claim 59 wherein said nucleic acid is RNA.

64. A kit for screening a human subject to diagnose schizophrenia or a genetic predisposition therefor, comprising, in association:

(a) an oligonucleotide useful as a probe for identifying polymorphisms in a human CON202 seven transmembrane receptor gene, the oligonucleotide comprising 6-50 nucleotides that have a sequence that is identical or exactly complementary to a portion of a wild type human CON202 gene sequence or CON202 coding sequence, except for one sequence difference selected from the group consisting of a nucleotide addition, a nucleotide deletion, or nucleotide substitution; and

(b) a media packaged with the oligonucleotide containing information identifying polymorphisms identifiable with the probe that correlate with schizophrenia or a genetic predisposition therefor.

65. A method of identifying a seven transmembrane allelic variant that correlates with a mental disorder, comprising steps of:

(a) providing a biological sample comprising nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny;

(b) analyzing said nucleic acid for the presence of a mutation or mutations in at least one seven transmembrane receptor that is expressed in the brain, wherein the at least one seven transmembrane receptor comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, and 20, or an allelic variant thereof, and wherein the nucleic acid includes sequence corresponding to the gene or genes encoding the at least one seven transmembrane receptor;

(c) determining a genotype for the patient for the at least one seven transmembrane receptor from said analyzing step; and

(d) identifying an allelic variant that correlates with the mental disorder from the determining step.

66. A method according to claim 65, wherein the disorder is schizophrenia, and wherein the at least one seven transmembrane receptor comprises CON202 having an amino acid sequence set forth in SEQ ID NO: 14, or an allelic variant thereof.

67. A purified and isolated polynucleotide comprising a nucleotide sequence encoding a CON202 receptor allelic variant identified according to claim 66.

68. A host cell transformed or transfected with a polynucleotide according to claim 67 or with a vector comprising the polynucleotide.

69. A purified polynucleotide comprising a nucleotide sequence encoding a CON202 seven transmembrane receptor protein of a human that is affected with schizophrenia;

wherein said polynucleotide hybridizes to the complement of SEQ ID

5 NO: 13 under the following hybridization conditions:

(a) hybridization for 16 hours at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate and

(b) washing 2 times for 30 minutes at 60°C in a wash solution comprising 0.1x SSC and 1% SDS; and

10 wherein the polynucleotide encodes a CON202 amino acid sequence that differs from SEQ ID NO: 14 at at least one residue.

70. A vector comprising a polynucleotide according to claim 69.

15 71. A host cell that has been transformed or transfected with a polynucleotide according to claim 70 and that expresses the CON202 protein encoded by the polynucleotide.

20 72. A host cell according to claim 71 that has been co-transfected with a polynucleotide encoding the CON202 amino acid sequence set forth in SEQ ID NO: 14 and that expresses the con202 protein having the amino acid sequence set forth in SEQ ID NO: 14.

25 73. A method for identifying a modulator of CON202 biological activity, comprising the steps of:

- a) contacting a cell according to claim 71 in the presence and in the absence of a putative modulator compound;
  - b) measuring CON202 biological activity in the cell; and
  - c) identifying a putative modulator compound in view of
- 30 decreased or increased CON202 biological activity in the presence versus absence of the putative modulator.

74. An assay to identify compounds useful for the treatment of schizophrenia, said assay comprising steps of:

(a) contacting a composition comprising a seven transmembrane receptor polypeptide according to claim 8 with a compound suspected of binding the seven transmembrane receptor polypeptide;

(b) measuring binding between the compound and the seven transmembrane receptor polypeptide; and

(c) identifying molecules that bind the seven transmembrane receptor as candidate compounds useful for the treatment of schizophrenia.

75. A method for identifying compound useful for a modulator of binding between a seven transmembrane receptor polypeptide and a binding partner of the seven transmembrane receptor polypeptide, which modulator is useful for treatment of schizophrenia, comprising the steps of:

(a) contacting the binding partner and a composition comprising the seven transmembrane receptor polypeptide in the presence and in the absence of a putative modulator compound, where the seven transmembrane receptor polypeptide is a polypeptide according to claim 8;

(b) measuring binding between the binding partner and the seven transmembrane receptor polypeptide;

(c) identifying a modulator compound useful for the treatment of schizophrenia in view of decreased or increased binding between the binding partner and seven transmembrane receptor polypeptide in the presence of the putative modulator, as compared to binding in the absence of the putative modulator.

76. An assay according to claim 74 or 75 wherein the composition comprises a cell expressing the seven transmembrane receptor polypeptide on its surface.

77. An assay according to claim 76 wherein the composition comprises a cell transformed or transfected with a polynucleotide encoding the seven transmembrane polypeptide and expressing the seven transmembrane receptor polypeptide on its surface.

- 1 -

## SEQUENCE LISTING

<110> Pharmacia & Upjohn Company  
<120> G PROTEIN-COUPLED RECEPTORS EXPRESSED IN BRAIN  
<130> 28341/6276P  
<140>  
<141>  
<150> US 09/481,794  
<151> 2000-01-12  
<150> US 09/454,399  
<151> 1999-12-03  
<150> US 09/429,517  
<151> 1999-10-28  
<150> US 09/429,555  
<151> 1999-10-28  
<150> US 09/429,676  
<151> 1999-10-28  
<150> US 09/429,695  
<151> 1999-10-28  
<150> US 09/428,114  
<151> 1999-10-27  
<150> US 09/428,020  
<151> 1999-10-27  
<150> US 09/427,859  
<151> 1999-10-27  
<150> US 09/427,653  
<151> 1999-10-27  
<160> 75  
<170> PatentIn Ver. 2.0  
<210> 1  
<211> 1308  
<212> DNA  
<213> Homo sapiens  
<220>  
<221> CDS  
<222> (157)..(1122)  
<220>  
<221> misc\_feature  
<222> (1)  
<223> N = A or C or G or T  
  
<220>  
<221> misc\_feature

- 2 -

&lt;222&gt; (1237)

&lt;223&gt; N = A or C or G or T

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1274)

&lt;223&gt; N = A or C or G or T

&lt;400&gt; 1

ntggttggtg gaccattaaa atgcattatg gaatttttaa aagttggggg agagggagac 60

agtaaaaata acctatattt tctcttggtt tttttttttt aactctagga aagcccagac 120

aaattttgag	ctatttcata	acctaccaga	cttatc	atg	cta	aca	ctg	aat	aaa	174
				Met	Leu	Thr	Leu	Asn	Lys	
				1				5		

aca	gac	cta	ata	cca	gct	tca	ttt	att	ctg	aat	gga	gtc	cca	gga	ctg	222
Thr	Asp	Leu	Ile	Pro	Ala	Ser	Phe	Ile	Leu	Asn	Gly	Val	Pro	Gly	Leu	
			10					15					20			

gaa	gac	aca	caa	ctc	tgg	att	tcc	ttc	cca	ttc	tgc	tct	atg	tat	gtt	270
Glu	Asp	Thr	Gln	Leu	Trp	Ile	Ser	Phe	Pro	Phe	Cys	Ser	Met	Tyr	Val	
		25				30						35				

gtg	gct	atg	gta	ggg	aat	tgt	gga	ctc	ctc	tac	ctc	att	cac	tat	gag	318
Val	Ala	Met	Val	Gly	Asn	Cys	Gly	Leu	Leu	Tyr	Leu	Ile	His	Tyr	Glu	
	40				45					50						

gat	gcc	ctg	cac	aaa	ccc	atg	tac	tac	ttc	ttg	gcc	atg	ctt	tcc	ttt	366
Asp	Ala	Leu	His	Lys	Pro	Met	Tyr	Tyr	Phe	Leu	Ala	Met	Leu	Ser	Phe	
	55			60					65						70	

act	gac	ctt	gtt	atg	tgc	tct	agt	aca	atc	cct	aaa	gcc	ctc	tgc	atc	414
Thr	Asp	Leu	Val	Met	Cys	Ser	Ser	Thr	Ile	Pro	Lys	Ala	Leu	Cys	Ile	
			75					80						85		

ttc	tgg	ttt	cat	ctc	aag	gac	att	gga	ttt	gat	gaa	tgc	ctt	gtc	cag	462
Phe	Trp	Phe	His	Leu	Lys	Asp	Ile	Gly	Phe	Asp	Glu	Cys	Leu	Val	Gln	
		90					95						100			

atg	ttc	ttc	atc	cac	acc	ttc	aca	ggg	atg	gag	tct	ggg	gtg	ctt	atg	510
Met	Phe	Phe	Ile	His	Thr	Phe	Thr	Gly	Met	Glu	Ser	Gly	Val	Leu	Met	
		105				110						115				

ctt	atg	gcc	ctg	gat	cgc	tat	gtg	gcc	atc	tgc	tac	ccc	tta	cgc	tat	558
Leu	Met	Ala	Leu	Asp	Arg	Tyr	Val	Ala	Ile	Cys	Tyr	Pro	Leu	Arg	Tyr	
	120				125					130						

tca	act	atc	ctc	acc	aat	cct	gta	att	gca	aag	gtt	ggg	act	gcc	acc	606
Ser	Thr	Ile	Leu	Thr	Asn	Pro	Val	Ile	Ala	Lys	Val	Gly	Thr	Ala	Thr	
	135			140						145				150		

ttc	ctg	aga	ggg	gta	tta	ctc	att	att	ccc	ttt	act	ttc	ctc	acc	aag	654
Phe	Leu	Arg	Gly	Val	Leu	Leu	Ile	Ile	Pro	Phe	Thr	Phe	Leu	Thr	Lys	
			155					160						165		

cgc	ctg	ccc	tcc	tgc	aga	ggc	aat	ata	ctt	ccc	cat	acc	tac	tgt	gac	702
Arg	Leu	Pro	Ser	Cys	Arg	Gly	Asn	Ile	Leu	Pro	His	Thr	Tyr	Cys	Asp	
			170					175						180		

- 3 -

cac atg tct gta gcc aaa ttg tcc tgt ggt aat gtc aag gtc aat gcc 750  
 His Met Ser Val Ala Lys Leu Ser Cys Gly Asn Val Lys Val Asn Ala  
 185 190 195

atc tat ggt ctg atg gtt gcc ctc ctg att ggg ggc ttt gac ata ctg 798  
 Ile Tyr Gly Leu Met Val Ala Leu Leu Ile Gly Gly Phe Asp Ile Leu  
 200 205 210

tgt atc acc atc tcc tat acc atg att ctc cgg gca gtg gtc agc ctc 846  
 Cys Ile Thr Ile Ser Tyr Thr Met Ile Leu Arg Ala Val Val Ser Leu  
 215 220 225 230

tcc tca gca gat gct cgg cag aag gcc ttt aat acc tgc act gcc cac 894  
 Ser Ser Ala Asp Arg Gln Lys Ala Phe Asn Thr Cys Thr Ala His  
 235 240 245

att tgt gcc att gtt ttc tcc tat act cca gct ttc ttc tcc ttc ttt 942  
 Ile Cys Ala Ile Val Phe Ser Tyr Thr Pro Ala Phe Phe Ser Phe Phe  
 250 255 260

tcc cac cgc ttt ggg gaa cac ata atc ccc cct tct tgc cac atc att 990  
 Ser His Arg Phe Gly Glu His Ile Ile Pro Pro Ser Cys His Ile Ile  
 265 270 275

gta gcc aat att tat ctg ctc cta cca ccc act atg aac cct att gtc 1038  
 Val Ala Asn Ile Tyr Leu Leu Leu Pro Pro Thr Met Asn Pro Ile Val  
 280 285 290

tat ggg gtg aaa acc aaa cag ata cga gac tgt gtc ata agg atc ctt 1086  
 Tyr Gly Val Lys Thr Lys Gln Ile Arg Asp Cys Val Ile Arg Ile Leu  
 295 300 305 310

tca ggt tct aag gat acc aaa tcc tac agc atg tga atgaacactt 1132  
 Ser Gly Ser Lys Asp Thr Lys Ser Tyr Ser Met  
 315 320

gccaggagtg agaagagaag gaaagaatta cttctatttg cctcttatgc aggagtcat 1192

aaaatcttcc tggaagtact gtattgatca caaaatggag tttgntgact ggtgcattct 1252

caataagtac cttgggaatc tnacatcact ggaaggccca ccacatttct ataaat 1308

<210> 2  
 <211> 321  
 <212> PRT  
 <213> Homo sapiens

<400> 2  
 Met Leu Thr Leu Asn Lys Thr Asp Leu Ile Pro Ala Ser Phe Ile Leu  
 1 5 10 15

Asn Gly Val Pro Gly Leu Glu Asp Thr Gln Leu Trp Ile Ser Phe Pro  
 20 25 30

Phe Cys Ser Met Tyr Val Val Ala Met Val Gly Asn Cys Gly Leu Leu  
 35 40 45

Tyr Leu Ile His Tyr Glu Asp Ala Leu His Lys Pro Met Tyr Tyr Phe  
 50 55 60

Leu Ala Met Leu Ser Phe Thr Asp Leu Val Met Cys Ser Ser Thr Ile  
 65 70 75 80

Pro	Lys	Ala	Leu	Cys	Ile	Phe	Trp	Phe	His	Leu	Lys	Asp	Ile	Gly	Phe
				85					90					95	
Asp	Glu	Cys	Leu	Val	Gln	Met	Phe	Phe	Ile	His	Thr	Phe	Thr	Gly	Met
			100					105					110		
Glu	Ser	Gly	Val	Leu	Met	Leu	Met	Ala	Leu	Asp	Arg	Tyr	Val	Ala	Ile
		115					120					125			
Cys	Tyr	Pro	Leu	Arg	Tyr	Ser	Thr	Ile	Leu	Thr	Asn	Pro	Val	Ile	Ala
	130					135					140				
Lys	Val	Gly	Thr	Ala	Thr	Phe	Leu	Arg	Gly	Val	Leu	Leu	Ile	Ile	Pro
145					150					155					160
Phe	Thr	Phe	Leu	Thr	Lys	Arg	Leu	Pro	Ser	Cys	Arg	Gly	Asn	Ile	Leu
				165					170					175	
Pro	His	Thr	Tyr	Cys	Asp	His	Met	Ser	Val	Ala	Lys	Leu	Ser	Cys	Gly
			180					185						190	
Asn	Val	Lys	Val	Asn	Ala	Ile	Tyr	Gly	Leu	Met	Val	Ala	Leu	Leu	Ile
		195					200					205			
Gly	Gly	Phe	Asp	Ile	Leu	Cys	Ile	Thr	Ile	Ser	Tyr	Thr	Met	Ile	Leu
	210					215					220				
Arg	Ala	Val	Val	Ser	Leu	Ser	Ser	Ala	Asp	Ala	Arg	Gln	Lys	Ala	Phe
225					230					235					240
Asn	Thr	Cys	Thr	Ala	His	Ile	Cys	Ala	Ile	Val	Phe	Ser	Tyr	Thr	Pro
				245					250					255	
Ala	Phe	Phe	Ser	Phe	Phe	Ser	His	Arg	Phe	Gly	Glu	His	Ile	Ile	Pro
			260					265					270		
Pro	Ser	Cys	His	Ile	Ile	Val	Ala	Asn	Ile	Tyr	Leu	Leu	Leu	Pro	Pro
		275					280					285			
Thr	Met	Asn	Pro	Ile	Val	Tyr	Gly	Val	Lys	Thr	Lys	Gln	Ile	Arg	Asp
	290					295						300			
Cys	Val	Ile	Arg	Ile	Leu	Ser	Gly	Ser	Lys	Asp	Thr	Lys	Ser	Tyr	Ser
305					310					315					320

```
<400> 3
atg gat gaa aca gga aat ctg aca gta tct tct gcc aca tgc cat gac 48
Met Asp Glu Thr Gly Asn Leu Thr Val Ser Ser Ala Thr Cys His Asp
      1             5             10             15
```

- 5 -

act att gat gac ttc cgc aat caa gtg tat tcc acc ttg tac tct atg	96
Thr Ile Asp Asp Phe Arg Asn Gln Val Tyr Ser Thr Leu Tyr Ser Met	
20 25 30	
atc tct gtt gta ggc ttc ttt ggc aat ggc ttt gtg ctc tat gtc ctc	144
Ile Ser Val Val Gly Phe Phe Gly Asn Gly Phe Val Leu Tyr Val Leu	
35 40 45	
ata aaa acc tat cac aag aag tca gcc ttc caa gta tac atg att aat	192
Ile Lys Thr Tyr His Lys Lys Ser Ala Phe Gln Val Tyr Met Ile Asn	
50 55 60	
tta gca gta gca gat cta ctt tgt gtg tgc aca ctg cct ctc cgt gtg	240
Leu Ala Val Ala Asp Leu Cys Val Cys Thr Leu Pro Leu Arg Val	
65 70 75 80	
gtc tat tat gtt cac aaa ggc att tgg ctc ttt ggt gac ttc ttg tgc	288
Val Tyr Tyr Val His Lys Gly Ile Trp Leu Phe Gly Asp Phe Leu Cys	
85 90 95	
cgc ctc agc acc tat gct ttg tat gtc aac ctc tat tgt agc atc ttc	336
Arg Leu Ser Thr Tyr Ala Leu Tyr Val Asn Leu Tyr Cys Ser Ile Phe	
100 105 110	
ttt atg aca gcc atg agc ttt ttc cgg tgc att gca att gtt ttt cca	384
Phe Met Thr Ala Met Ser Phe Phe Arg Cys Ile Ala Ile Val Phe Pro	
115 120 125	
gtc cag aac att aat ttg gtt aca cag aaa aaa gcc agg ttt gtg tgt	432
Val Gln Asn Ile Asn Leu Val Thr Gln Lys Lys Ala Arg Phe Val Cys	
130 135 140	
gta ggt att tgg att ttt gtg att ttg acc agt tct cca ttt cta atg	480
Val Gly Ile Trp Ile Phe Val Ile Leu Thr Ser Ser Pro Phe Leu Met	
145 150 155 160	
gcc aaa cca caa aaa gat gag aaa aat aat acc aag tgc ttt gag ccc	528
Ala Lys Pro Gln Lys Asp Glu Lys Asn Asn Thr Lys Cys Phe Glu Pro	
165 170 175	
cca caa gac aat caa act aaa aat cat gtt ttg gtc ttg cat tat gtg	576
Pro Gln Asp Asn Gln Thr Lys Asn His Val Leu Val Leu His Tyr Val	
180 185 190	
tca ttg ttt gtt ggc ttt atc atc cct ttt gtt att ata att gtc tgt	624
Ser Leu Phe Val Gly Phe Ile Ile Pro Phe Val Ile Ile Ile Val Cys	
195 200 205	
tac aca atg atc att ttg acc tta cta aaa aaa tca atg aaa aaa aat	672
Tyr Thr Met Ile Ile Leu Thr Leu Leu Lys Lys Ser Met Lys Lys Asn	
210 215 220	
ctg tca agt cat aaa aag gct ata gga atg atc atg gtc gtg acc gct	720
Leu Ser Ser His Lys Lys Ala Ile Gly Met Ile Met Val Val Thr Ala	
225 230 235 240	
gcc ttt tta gtc agt ttc atg cca tat cat att caa cgt acc att cac	768
Ala Phe Leu Val Ser Phe Met Pro Tyr His Ile Gln Arg Thr Ile His	
245 250 255	
ctt cat ttt tta cac aat gaa act aaa ccc tgt gat tct gtc ctt aga	816
Leu His Phe Leu His Asn Glu Thr Lys Pro Cys Asp Ser Val Leu Arg	
260 265 270	

- 6 -

```

atg cag aag tcc gtg gtc ata acc ttg tct ctg gct gca tcc aat tgt 864
Met Gln Lys Ser Val Val Ile Thr Leu Ser Leu Ala Ala Ser Asn Cys
      275                280                285

tgc ttt gac cct ctc cta tat ttc ttt tct ggg ggt aac ttt agg aaa 912
Cys Phe Asp Pro Leu Leu Tyr Phe Phe Ser Gly Gly Asn Phe Arg Lys
      290                295                300

agg ctg tct aca ttt aga aag cat tct ttg tcc agc gtg act tat gta 960
Arg Leu Ser Thr Phe Arg Lys His Ser Leu Ser Ser Val Thr Tyr Val
      305                310                315                320

ccc aga aag aag gcc tct ttg cca gaa aaa gga gaa gaa ata tgt aaa 1008
Pro Arg Lys Lys Ala Ser Leu Pro Glu Lys Gly Glu Glu Ile Cys Lys
      325                330                335

gta tag
Val                                           1014

```

<210> 4  
 <211> 337  
 <212> PRT  
 <213> Homo sapiens

```

<400> 2
Met Asp Glu Thr Gly Asn Leu Thr Val Ser Ser Ala Thr Cys His Asp
 1          5          10          15

Thr Ile Asp Asp Phe Arg Asn Gln Val Tyr Ser Thr Leu Tyr Ser Met
      20          25          30

Ile Ser Val Val Gly Phe Phe Gly Asn Gly Phe Val Leu Tyr Val Leu
      35          40          45

Ile Lys Thr Tyr His Lys Lys Ser Ala Phe Gln Val Tyr Met Ile Asn
      50          55          60

Leu Ala Val Ala Asp Leu Leu Cys Val Cys Thr Leu Pro Leu Arg Val
      65          70          75          80

Val Tyr Tyr Val His Lys Gly Ile Trp Leu Phe Gly Asp Phe Leu Cys
      85          90          95

Arg Leu Ser Thr Tyr Ala Leu Tyr Val Asn Leu Tyr Cys Ser Ile Phe
      100         105         110

Phe Met Thr Ala Met Ser Phe Phe Arg Cys Ile Ala Ile Val Phe Pro
      115         120         125

Val Gln Asn Ile Asn Leu Val Thr Gln Lys Lys Ala Arg Phe Val Cys
      130         135         140

Val Gly Ile Trp Ile Phe Val Ile Leu Thr Ser Ser Pro Phe Leu Met
      145         150         155         160

Ala Lys Pro Gln Lys Asp Glu Lys Asn Asn Thr Lys Cys Phe Glu Pro
      165         170         175

Pro Gln Asp Asn Gln Thr Lys Asn His Val Leu Val Leu His Tyr Val
      180         185         190

```

- 7 -

Ser Leu Phe Val Gly Phe Ile Ile Pro Phe Val Ile Ile Ile Val Cys  
 195 200 205  
 Tyr Thr Met Ile Ile Leu Thr Leu Leu Lys Lys Ser Met Lys Lys Asn  
 210 215 220  
 Leu Ser Ser His Lys Lys Ala Ile Gly Met Ile Met Val Val Thr Ala  
 225 230 235 240  
 Ala Phe Leu Val Ser Phe Met Pro Tyr His Ile Gln Arg Thr Ile His  
 245 250 255  
 Leu His Phe Leu His Asn Glu Thr Lys Pro Cys Asp Ser Val Leu Arg  
 260 265 270  
 Met Gln Lys Ser Val Val Ile Thr Leu Ser Leu Ala Ala Ser Asn Cys  
 275 280 285  
 Cys Phe Asp Pro Leu Leu Tyr Phe Phe Ser Gly Gly Asn Phe Arg Lys  
 290 295 300  
 Arg Leu Ser Thr Phe Arg Lys His Ser Leu Ser Ser Val Thr Tyr Val  
 305 310 315 320  
 Pro Arg Lys Lys Ala Ser Leu Pro Glu Lys Gly Glu Glu Ile Cys Lys  
 325 330 335  
 Val

<210> 5  
 <211> 2429  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (691)..(1845)

<400> 5  
 ggggcctact tcaccgtgta cccggacttg ggaccatcac agacttcaga accatcagga 60  
 acctgggagc aactgaaagc tgaactacag tgggctttca gacacacagc aggctgcgga 120  
 gcacaaatag gactgggttc ctccaggcca ccagcagggc ggtggagggtc ttcactgact 180  
 ccctgcctac ctctcaggac aatgtccttt tggctccaca gtccctgaag ccagagctgg 240  
 tgggggcagg gaggcagcca ccagcctcta tatgtagtgg aggaggggggt gtccagggag 300  
 ggctgcatga tcctgagagc cccacactca cccggctgga ctatcctccc acttcagggt 360  
 ttctctgggc ttccatcttg cccctgctga gccctgcttc ctctctacc agcagcacia 420  
 cccccaggct gggctcagag acctcatgtg gtgggatcac tcagtacccc gaggcggagg 480  
 gaaggagggga gggctgcagg gtcccccttg gcctgcaaac aggaacacag ggtgtttctc 540  
 agtggctgcg agaatgctga tgaaaacccc aggatgttgt gtcaccgtgg tggccagctg 600  
 atagtgccaa tcatcccaact ttgccctgag cactcctgca ggggtagaag actccagaac 660

- 8 -

```

cttctctcag gcccatggcc caagcagccc atg gaa ctt cat aac ctg agc tct 714
                               Met Glu Leu His Asn Leu Ser Ser
                               1                               5

cca tct ccc tct ctc tcc tcc tct gtt ctc cct ccc tcc ttc tct ccc 762
Pro Ser Pro Ser Leu Ser Ser Ser Val Leu Pro Pro Ser Phe Ser Pro
   10                               15                               20

tca ccc tcc tct gct ccc tct gcc ttt acc act gtg ggg ggg tcc tct 810
Ser Pro Ser Ser Ala Pro Ser Ala Phe Thr Thr Val Gly Gly Ser Ser
   25                               30                               35                               40

gga ggg ccc tgc cac ccc acc tct tcc tcg ctg gtg tct gcc ttc ctg 858
Gly Gly Pro Cys His Pro Thr Ser Ser Ser Leu Val Ser Ala Phe Leu
                               45                               50                               55

gca cca atc ctg gcc ctg gag ttt gtc ctg ggc ctg gtg ggg aac agt 906
Ala Pro Ile Leu Ala Leu Glu Phe Val Leu Gly Leu Val Gly Asn Ser
                               60                               65                               70

ttg gcc ctc ttc atc ttc tgc atc cac acg cgg ccc tgg acc tcc aac 954
Leu Ala Leu Phe Ile Phe Cys Ile His Thr Arg Pro Trp Thr Ser Asn
   75                               80                               85

acg gtg ttc ctg gtc agc ctg gtg gcc gct gac ttc ctc ctg atc agc 1002
Thr Val Phe Leu Val Ser Leu Val Ala Ala Asp Phe Leu Leu Ile Ser
   90                               95                               100

aac ctg ccc ctc cgc gtg gac tac tac ctc ctc cat gag acc tgg cgc 1050
Asn Leu Pro Leu Arg Val Asp Tyr Tyr Leu Leu His Glu Thr Trp Arg
  105                               110                               115                               120

ttt ggg gct gct gcc tgc aaa gtc aac ctc ttc atg ctg tcc acc aac 1098
Phe Gly Ala Ala Ala Cys Lys Val Asn Leu Phe Met Leu Ser Thr Asn
                               125                               130                               135

cgc acg gcc agc gtt gtc ttc ctc aca gcc atc gca ctc aac cgc tac 1146
Arg Thr Ala Ser Val Val Phe Leu Thr Ala Ile Ala Leu Asn Arg Tyr
                               140                               145                               150

ctg aag gtg gtg cag ccc cac cac gtg ctg agc cgt gct tcc gtg ggg 1194
Leu Lys Val Val Gln Pro His His Val Leu Ser Arg Ala Ser Val Gly
   155                               160                               165

gca gct gcc cgg gtg gcc ggg gga ctc tgg gtg ggc atc ctg ctc ctc 1242
Ala Ala Ala Arg Val Ala Gly Gly Leu Trp Val Gly Ile Leu Leu Leu
   170                               175                               180

aac ggg cac ctg ctc ctg agc acc ttc tcc ggc ccc tcc tgc ctc agc 1290
Asn Gly His Leu Leu Leu Ser Thr Phe Ser Gly Pro Ser Cys Leu Ser
  185                               190                               195                               200

tac agg gtg ggc acg aag ccc tcg gcc tcg ctc cgc tgg cac cag gca 1338
Tyr Arg Val Gly Thr Lys Pro Ser Ala Ser Leu Arg Trp His Gln Ala
                               205                               210                               215

ctg tac ctg ctg gag ttc ttc ctg cca ctg gcg ctc atc ctc ttt gct 1386
Leu Tyr Leu Leu Glu Phe Phe Leu Pro Leu Ala Leu Ile Leu Phe Ala
   220                               225                               230

```

att	gtg	agc	att	ggg	ctc	acc	atc	cgg	aac	cgt	ggc	ctg	ggc	ggg	cag	1434
Ile	Val	Ser	Ile	Gly	Leu	Thr	Ile	Arg	Asn	Arg	Gly	Leu	Gly	Gly	Gln	
		235					240					245				
gca	ggc	ccg	cag	agg	gcc	atg	cgt	gtg	ctg	gcc	atg	gtg	gtg	gcc	gtc	1482
Ala	Gly	Pro	Gln	Arg	Ala	Met	Arg	Val	Leu	Ala	Met	Val	Val	Ala	Val	
		250				255					260					
tac	acc	atc	tgc	ttc	ttg	ccc	agc	atc	atc	ttt	ggc	atg	gct	tcc	atg	1530
Tyr	Thr	Ile	Cys	Phe	Leu	Pro	Ser	Ile	Ile	Phe	Gly	Met	Ala	Ser	Met	
					270					275					280	
gtg	gct	ttc	tgg	ctg	tcc	gcc	tgc	cga	tcc	ctg	gac	ctc	tgc	aca	cag	1578
Val	Ala	Phe	Trp	Leu	Ser	Ala	Cys	Arg	Ser	Leu	Asp	Leu	Cys	Thr	Gln	
				285					290					295		
ctc	ttc	cat	ggc	tcc	ctg	gcc	ttc	acc	tac	ctc	aac	agt	gtc	ctg	gac	1626
Leu	Phe	His	Gly	Ser	Leu	Ala	Phe	Thr	Tyr	Leu	Asn	Ser	Val	Leu	Asp	
			300					305					310			
ccc	gtg	ctc	tac	tgc	ttc	tct	agc	ccc	aac	ttc	ctc	cac	cag	agc	cgg	1674
Pro	Val	Leu	Tyr	Cys	Phe	Ser	Ser	Pro	Asn	Phe	Leu	His	Gln	Ser	Arg	
		315					320					325				
gcc	ttg	ctg	ggc	ctc	acg	cgg	ggc	cgg	cag	ggc	cca	gtg	agc	gac	gag	1722
Ala	Leu	Leu	Gly	Leu	Thr	Arg	Gly	Arg	Gln	Gly	Pro	Val	Ser	Asp	Glu	
		330				335					340					
agc	tcc	tac	caa	ccc	tcc	agg	cag	tgg	cgc	tac	cgg	gag	gcc	tct	agg	1770
Ser	Ser	Tyr	Gln	Pro	Ser	Arg	Gln	Trp	Arg	Tyr	Arg	Glu	Ala	Ser	Arg	
		345			350					355					360	
aag	gcg	gag	gcc	ata	ggg	aag	ctg	aaa	gtg	cag	ggc	gag	gtc	tct	ctg	1818
Lys	Ala	Glu	Ala	Ile	Gly	Lys	Leu	Lys	Val	Gln	Gly	Glu	Val	Ser	Leu	
				365					370					375		
gaa	aag	gaa	ggc	tcc	tcc	cag	ggc	tga	gggccagctg	cagggctgca						1865
Glu	Lys	Glu	Gly	Ser	Ser	Gln	Gly									
			380					385								
gcgctgtggg	ggtaagggct	gccgcgctct	ggcctggagg	gacaaggcca	gcacacggtg											1925
cctcaaccaa	ctggacaagg	gatggcggca	gaccaggggc	caggccaaag	cactggcagg											1985
actcatgttg	gtggcaggga	gagaaacca	cctaggcctc	tcagtgtgtc	caggatggca											2045
ttcccagaat	gcagggggaga	gcaggatgcc	gggtggagga	gacaggcaag	gtgccgttgg											2105
cacaccagct	cagacagggg	cctgcgcagc	tgcaggggac	agacgccaat	cactgtcaca											2165
gcagagtcac	cttagaaatt	ggacagctgc	atgttctgtg	ctctccagtt	tgtcccttcc											2225
aatattaata	aacttccctt	ttaaataatat	ttatttgcag	accaatatct	gtctttaatt											2285
ctaacctggg	actgtcagta	ggcgtcaaag	tgagcgcccc	agtgaaggaa	ccttggagag											2345
agtgggagca	ttcccagcct	tccaggggga	ctcgtcttcc	agacttttga	gcccgc											

- 10 -

<210> 6  
 <211> 384  
 <212> PRT  
 <213> Homo sapiens

<400> 6

Met Glu Leu His Asn Leu Ser Ser Pro Ser Pro Ser Leu Ser Ser Ser  
 1 5 10 15  
 Val Leu Pro Pro Ser Phe Ser Pro Ser Pro Ser Ser Ala Pro Ser Ala  
 20 25 30  
 Phe Thr Thr Val Gly Gly Ser Ser Gly Gly Pro Cys His Pro Thr Ser  
 35 40 45  
 Ser Ser Leu Val Ser Ala Phe Leu Ala Pro Ile Leu Ala Leu Glu Phe  
 50 55 60  
 Val Leu Gly Leu Val Gly Asn Ser Leu Ala Leu Phe Ile Phe Cys Ile  
 65 70 75 80  
 His Thr Arg Pro Trp Thr Ser Asn Thr Val Phe Leu Val Ser Leu Val  
 85 90 95  
 Ala Ala Asp Phe Leu Leu Ile Ser Asn Leu Pro Leu Arg Val Asp Tyr  
 100 105 110  
 Tyr Leu Leu His Glu Thr Trp Arg Phe Gly Ala Ala Ala Cys Lys Val  
 115 120 125  
 Asn Leu Phe Met Leu Ser Thr Asn Arg Thr Ala Ser Val Val Phe Leu  
 130 135 140  
 Thr Ala Ile Ala Leu Asn Arg Tyr Leu Lys Val Val Gln Pro His His  
 145 150 155 160  
 Val Leu Ser Arg Ala Ser Val Gly Ala Ala Ala Arg Val Ala Gly Gly  
 165 170 175  
 Leu Trp Val Gly Ile Leu Leu Leu Asn Gly His Leu Leu Leu Ser Thr  
 180 185 190  
 Phe Ser Gly Pro Ser Cys Leu Ser Tyr Arg Val Gly Thr Lys Pro Ser  
 195 200 205  
 Ala Ser Leu Arg Trp His Gln Ala Leu Tyr Leu Leu Glu Phe Phe Leu  
 210 215 220  
 Pro Leu Ala Leu Ile Leu Phe Ala Ile Val Ser Ile Gly Leu Thr Ile  
 225 230 235 240  
 Arg Asn Arg Gly Leu Gly Gly Gln Ala Gly Pro Gln Arg Ala Met Arg  
 245 250 255  
 Val Leu Ala Met Val Val Ala Val Tyr Thr Ile Cys Phe Leu Pro Ser  
 260 265 270  
 Ile Ile Phe Gly Met Ala Ser Met Val Ala Phe Trp Leu Ser Ala Cys  
 275 280 285  
 Arg Ser Leu Asp Leu Cys Thr Gln Leu Phe His Gly Ser Leu Ala Phe  
 290 295 300

- 11 -

Thr Tyr Leu Asn Ser Val Leu Asp Pro Val Leu Tyr Cys Phe Ser Ser  
305 310 315 320

Pro Asn Phe Leu His Gln Ser Arg Ala Leu Leu Gly Leu Thr Arg Gly  
325 330 335

Arg Gln Gly Pro Val Ser Asp Glu Ser Ser Tyr Gln Pro Ser Arg Gln  
340 345 350

Trp Arg Tyr Arg Glu Ala Ser Arg Lys Ala Glu Ala Ile Gly Lys Leu  
355 360 365

Lys Val Gln Gly Glu Val Ser Leu Glu Lys Glu Gly Ser Ser Gln Gly  
370 375 380

<210> 7

<211> 1484

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (146)..(1147)

<400> 7

ttgaatttag gtgacactat agaagagcta tgacgtcgca tgcacgcgta cgtaagctcg 60

gaattcggtc cgagctgaac taatgactgc cgccataaga agacagagag aactgagtat 120

cctcccaaag gtgacactgg aagca atg aac acc aca gtg atg caa ggc ttc 172  
Met Asn Thr Thr Val Met Gln Gly Phe  
1 5

aac aga tct gag cgg tgc ccc aga gac act cgg ata gta cag ctg gta 220  
Asn Arg Ser Glu Arg Cys Pro Arg Asp Thr Arg Ile Val Gln Leu Val  
10 15 20 25

ttc cca gcc ctc tac aca gtg gtt ttc ttg acc ggc atc ctg ctg aat 268  
Phe Pro Ala Leu Tyr Thr Val Val Phe Leu Thr Gly Ile Leu Leu Asn  
30 35 40

act ttg gct ctg tgg gtg ttt gtt cac atc ccc agc tcc tcc acc ttc 316  
Thr Leu Ala Leu Trp Val Phe Val His Ile Pro Ser Ser Ser Thr Phe  
45 50 55

atc atc tac ctc aaa aac act ttg gtg gcc gac ttg ata atg aca ctc 364  
Ile Ile Tyr Leu Lys Asn Thr Leu Val Ala Asp Leu Ile Met Thr Leu  
60 65 70

atg ctt cct ttc aaa atc ctc tct gac tca cac ctg gca ccc tgg cag 412  
Met Leu Pro Phe Lys Ile Leu Ser Asp Ser His Leu Ala Pro Trp Gln  
75 80 85

ctc aga gct ttt gtg tgt cgt ttt tct tcg gtg ata ttt tat gag acc 460  
Leu Arg Ala Phe Val Cys Arg Phe Ser Ser Val Ile Phe Tyr Glu Thr  
90 95 100 105

atg tat gtg ggc atc gtg ctg tta ggg ctc ata gcc ttt gac aga ttc 508  
Met Tyr Val Gly Ile Val Leu Leu Gly Leu Ile Ala Phe Asp Arg Phe  
110 115 120

- 12 -

ctc aag atc atc aga cct ttg aga aat att ttt cta aaa aaa cct gtt 556  
 Leu Lys Ile Ile Arg Pro Leu Arg Asn Ile Phe Leu Lys Lys Pro Val  
 125 130 135

ttt gca aaa acg gtc tca atc ttc atc tgg gtc ttt ttg gtc ttc atc 604  
 Phe Ala Lys Thr Val Ser Ile Phe Ile Trp Val Phe Leu Val Phe Ile  
 140 145 150

tcc ctg cca aat atg atc ttg agc aac aag gaa gca aca cca tcg tct 652  
 Ser Leu Pro Asn Met Ile Leu Ser Asn Lys Glu Ala Thr Pro Ser Ser  
 155 160 165

gtg aaa aag tgt gct tcc tta aag ggg cct ctg ggg ctg aaa tgg cat 700  
 Val Lys Lys Cys Ala Ser Leu Lys Gly Pro Leu Gly Leu Lys Trp His  
 170 175 180 185

caa atg gta aat aac ata tgc cag ttt att ttc tgg act ggt ttt atc 748  
 Gln Met Val Asn Ile Cys Gln Phe Ile Phe Trp Thr Gly Phe Ile  
 190 195 200

cta atg ctt gtg ttt tat gtg gtt att gca aaa aaa gta tat gat tct 796  
 Leu Met Leu Val Phe Tyr Val Val Ile Ala Lys Lys Val Tyr Asp Ser  
 205 210 215

tat aga aag tcc aaa agt aag gac aga aaa aac aac aaa aag ctg gaa 844  
 Tyr Arg Lys Ser Lys Ser Lys Asp Arg Lys Asn Asn Lys Lys Leu Glu  
 220 225 230

ggc aaa gta ttt gtt gtc gtg gct gtc ttc ttt gtg tgt ttt gct cca 892  
 Gly Lys Val Phe Val Val Val Ala Val Phe Phe Val Cys Phe Ala Pro  
 235 240 245

ttt cat ttt gcc aga gtt cca tat act cac agt caa acc aac aat aag 940  
 Phe His Phe Ala Arg Val Pro Tyr Thr His Ser Gln Thr Asn Asn Lys  
 250 255 260 265

act gac tgt aga ctg caa aat caa ctg ttt att gct aaa gaa aca act 988  
 Thr Asp Cys Arg Leu Gln Asn Gln Leu Phe Ile Ala Lys Glu Thr Thr  
 270 275 280

ctc ttt ttg gca gca act aac att tgt atg gat ccc tta ata tac ata 1036  
 Leu Phe Leu Ala Ala Thr Asn Ile Cys Met Asp Pro Leu Ile Tyr Ile  
 285 290 295

ttc tta tgt aaa aaa ttc aca gaa aag cta cca tgt atg caa ggg aga 1084  
 Phe Leu Cys Lys Lys Phe Thr Glu Lys Leu Pro Cys Met Gln Gly Arg  
 300 305 310

aag acc aca gca tca agc caa gaa aat cat agc agt cag aca gac aac 1132  
 Lys Thr Thr Ala Ser Ser Gln Glu Asn His Ser Ser Gln Thr Asp Asn  
 315 320 325

ata acc tta ggc tga caactgtaca taggggtaac ttctatttat tgatgagact 1187  
 Ile Thr Leu Gly  
 330

tccgtagata atgtggaaat caaatttaac caagaaaaaa agattggaac aaatgctctc 1247

ttacatttta ttatcctgg tgtccaggaa aagattatat taaatttaaa tccacataga 1307

tctattcata agctgaatga accattacct aagagaatgc aacaggatac caatggccac 1367

tagaggcata ttccctcttc tttttttttt gttaaatttc aagagcatte actttacatt 1427

- 13 -

tggaagact aaggggaacg gttatcctac aaacctcct tcaacacctt ttacatt 1484

&lt;210&gt; 8

&lt;211&gt; 333

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 8

Met	Asn	Thr	Thr	Val	Met	Gln	Gly	Phe	Asn	Arg	Ser	Glu	Arg	Cys	Pro
1				5					10					15	

Arg	Asp	Thr	Arg	Ile	Val	Gln	Leu	Val	Phe	Pro	Ala	Leu	Tyr	Thr	Val
		20						25					30		

Val	Phe	Leu	Thr	Gly	Ile	Leu	Leu	Asn	Thr	Leu	Ala	Leu	Trp	Val	Phe
	35						40					45			

Val	His	Ile	Pro	Ser	Ser	Ser	Thr	Phe	Ile	Ile	Tyr	Leu	Lys	Asn	Thr
	50					55					60				

Leu	Val	Ala	Asp	Leu	Ile	Met	Thr	Leu	Met	Leu	Pro	Phe	Lys	Ile	Leu
65					70					75					80

Ser	Asp	Ser	His	Leu	Ala	Pro	Trp	Gln	Leu	Arg	Ala	Phe	Val	Cys	Arg
				85					90					95	

Phe	Ser	Ser	Val	Ile	Phe	Tyr	Glu	Thr	Met	Tyr	Val	Gly	Ile	Val	Leu
			100					105					110		

Leu	Gly	Leu	Ile	Ala	Phe	Asp	Arg	Phe	Leu	Lys	Ile	Ile	Arg	Pro	Leu
	115						120					125			

Arg	Asn	Ile	Phe	Leu	Lys	Lys	Pro	Val	Phe	Ala	Lys	Thr	Val	Ser	Ile
	130					135					140				

Phe	Ile	Trp	Val	Phe	Leu	Val	Phe	Ile	Ser	Leu	Pro	Asn	Met	Ile	Leu
145				150						155					160

Ser	Asn	Lys	Glu	Ala	Thr	Pro	Ser	Ser	Val	Lys	Lys	Cys	Ala	Ser	Leu
			165						170					175	

Lys	Gly	Pro	Leu	Gly	Leu	Lys	Trp	His	Gln	Met	Val	Asn	Asn	Ile	Cys
		180					185						190		

Gln	Phe	Ile	Phe	Trp	Thr	Gly	Phe	Ile	Leu	Met	Leu	Val	Phe	Tyr	Val
	195					200						205			

Val	Ile	Ala	Lys	Lys	Val	Tyr	Asp	Ser	Tyr	Arg	Lys	Ser	Lys	Ser	Lys
	210					215					220				

Asp	Arg	Lys	Asn	Asn	Lys	Lys	Leu	Glu	Gly	Lys	Val	Phe	Val	Val	Val
225				230						235					240

Ala	Val	Phe	Phe	Val	Cys	Phe	Ala	Pro	Phe	His	Phe	Ala	Arg	Val	Pro
			245						250				255		

Tyr	Thr	His	Ser	Gln	Thr	Asn	Asn	Lys	Thr	Asp	Cys	Arg	Leu	Gln	Asn
			260					265					270		

Gln	Leu	Phe	Ile	Ala	Lys	Glu	Thr	Thr	Leu	Phe	Leu	Ala	Ala	Thr	Asn
	275						280					285			

- 14 -

Ile Cys Met Asp Pro Leu Ile Tyr Ile Phe Leu Cys Lys Lys Phe Thr  
 290 295 300

Glu Lys Leu Pro Cys Met Gln Gly Arg Lys Thr Thr Ala Ser Ser Gln  
 305 310 315 320

Glu Asn His Ser Ser Gln Thr Asp Asn Ile Thr Leu Gly  
 325 330

&lt;210&gt; 9

&lt;211&gt; 957

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(954)

&lt;400&gt; 9

```

atg atg gtg gat ccc aat ggc aat gaa tcc agt gct aca tac ttc atc   48
Met Met Val Asp Pro Asn Gly Asn Glu Ser Ser Ala Thr Tyr Phe Ile
  1             5             10             15

cta ata ggc ctc cct ggt tta gaa gag gct cag ttc tgg ttg gcc ttc   96
Leu Ile Gly Leu Pro Gly Leu Glu Glu Ala Gln Phe Trp Leu Ala Phe
          20             25             30

cca ttg tgc tcc ctc tac ctt att gct gtg cta ggt aac ttg aca atc   144
Pro Leu Cys Ser Leu Tyr Leu Ile Ala Val Leu Gly Asn Leu Thr Ile
          35             40             45

atc tac att gtg cgg act gag cac agc ctg cat gag ccc atg tat ata   192
Ile Tyr Ile Val Arg Thr Glu His Ser Leu His Glu Pro Met Tyr Ile
          50             55             60

ttt ctt tgc atg ctt tca ggc att gac atc ctc atc tcc acc tca tcc   240
Phe Leu Cys Met Leu Ser Gly Ile Asp Ile Leu Ile Ser Thr Ser Ser
  65             70             75             80

atg ccc aaa atg ctg gcc atc ttc tgg ttc aat tcc act acc atc cag   288
Met Pro Lys Met Leu Ala Ile Phe Trp Phe Asn Ser Thr Thr Ile Gln
          85             90             95

ttt gat gct tgt ctg cta cag atg ttt gcc atc cac tcc tta tct ggc   336
Phe Asp Ala Cys Leu Leu Gln Met Phe Ala Ile His Ser Leu Ser Gly
          100            105            110

atg gaa tcc aca gtg ctg ctg gcc atg gct ttt gac cgc tat gtg gcc   384
Met Glu Ser Thr Val Leu Leu Ala Met Ala Phe Asp Arg Tyr Val Ala
          115            120            125

atc tgt cac cca ctg cgc cat gcc aca gta ctt acg ttg cct cgt gtc   432
Ile Cys His Pro Leu Arg His Ala Thr Val Leu Thr Leu Pro Arg Val
          130            135            140

acc aaa att ggt gtg gct gct gtg gtg cgg ggg gct gca ctg atg gca   480
Thr Lys Ile Gly Val Ala Ala Val Val Arg Gly Ala Ala Leu Met Ala
          145            150            155            160

ccc ctt cct gtc ttc atc aag cag ctg ccc ttc tgc cgc tcc aat atc   528
Pro Leu Pro Val Phe Ile Lys Gln Leu Pro Phe Cys Arg Ser Asn Ile
          165            170            175

```

- 15 -

ctt tcc cat tcc tac tgc cta cac caa gat gtc atg aag ctg gcc tgt 576  
 Leu Ser His Ser Tyr Cys Leu His Gln Asp Val Met Lys Leu Ala Cys  
 180 185 190

gat gat atc cgg gtc aat gtc gtc tat ggc ctt atc gtc atc atc tcc 624  
 Asp Asp Ile Arg Val Asn Val Val Tyr Gly Leu Ile Val Ile Ile Ser  
 195 200 205

gcc att ggc ctg gac tca ctt ctc atc tcc ttc tca tat ctg ctt att 672  
 Ala Ile Gly Leu Asp Ser Leu Leu Ile Ser Phe Ser Tyr Leu Leu Ile  
 210 215 220

ctt aag act gtg ttg ggc ttg aca cgt gaa gcc cag gcc aag gca ttt 720  
 Leu Lys Thr Val Leu Gly Leu Thr Arg Glu Ala Gln Ala Lys Ala Phe  
 225 230 235 240

ggc act tgc gtc tct cat gtg tgt gct gtg ttc ata ttc tat gta cct 768  
 Gly Thr Cys Val Ser His Val Cys Ala Val Phe Ile Phe Tyr Val Pro  
 245 250 255

ttc att gga ttg tcc atg gtg cat cgc ttt agc aag cgg cgt gac tct 816  
 Phe Ile Gly Leu Ser Met Val His Arg Phe Ser Lys Arg Arg Asp Ser  
 260 265 270

ccg ctg ccc gtc atc ttg gcc aat atc tat ctg ctg gtt cct cct gtg 864  
 Pro Leu Pro Val Ile Leu Ala Asn Ile Tyr Leu Leu Val Pro Pro Val  
 275 280 285

ctc aac cca att gtc tat gga gtg aag aca aag gag att cga cag cgc 912  
 Leu Asn Pro Ile Val Tyr Gly Val Lys Thr Lys Glu Ile Arg Gln Arg  
 290 295 300

atc ctt cga ctt ttc cat gtg gcc aca cac gct tca gag ccc tag 957  
 Ile Leu Arg Leu Phe His Val Ala Thr His Ala Ser Glu Pro  
 305 310 315

<210> 10  
 <211> 318  
 <212> PRT  
 <213> Homo sapiens

<400> 10  
 Met Met Val Asp Pro Asn Gly Asn Glu Ser Ser Ala Thr Tyr Phe Ile  
 1 5 10 15

Leu Ile Gly Leu Pro Gly Leu Glu Glu Ala Gln Phe Trp Leu Ala Phe  
 20 25 30

Pro Leu Cys Ser Leu Tyr Leu Ile Ala Val Leu Gly Asn Leu Thr Ile  
 35 40 45

Ile Tyr Ile Val Arg Thr Glu His Ser Leu His Glu Pro Met Tyr Ile  
 50 55 60

Phe Leu Cys Met Leu Ser Gly Ile Asp Ile Leu Ile Ser Thr Ser Ser  
 65 70 75 80

Met Pro Lys Met Leu Ala Ile Phe Trp Phe Asn Ser Thr Thr Ile Gln  
 85 90 95

Phe Asp Ala Cys Leu Leu Gln Met Phe Ala Ile His Ser Leu Ser Gly  
 100 105 110

- 16 -

Met Glu Ser Thr Val Leu Leu Ala Met Ala Phe Asp Arg Tyr Val Ala  
 115 120 125

Ile Cys His Pro Leu Arg His Ala Thr Val Leu Thr Leu Pro Arg Val  
 130 135 140

Thr Lys Ile Gly Val Ala Ala Val Val Arg Gly Ala Ala Leu Met Ala  
 145 150 155 160

Pro Leu Pro Val Phe Ile Lys Gln Leu Pro Phe Cys Arg Ser Asn Ile  
 165 170 175

Leu Ser His Ser Tyr Cys Leu His Gln Asp Val Met Lys Leu Ala Cys  
 180 185 190

Asp Asp Ile Arg Val Asn Val Val Tyr Gly Leu Ile Val Ile Ile Ser  
 195 200 205

Ala Ile Gly Leu Asp Ser Leu Leu Ile Ser Phe Ser Tyr Leu Leu Ile  
 210 215 220

Leu Lys Thr Val Leu Gly Leu Thr Arg Glu Ala Gln Ala Lys Ala Phe  
 225 230 235 240

Gly Thr Cys Val Ser His Val Cys Ala Val Phe Ile Phe Tyr Val Pro  
 245 250 255

Phe Ile Gly Leu Ser Met Val His Arg Phe Ser Lys Arg Arg Asp Ser  
 260 265 270

Pro Leu Pro Val Ile Leu Ala Asn Ile Tyr Leu Leu Val Pro Pro Val  
 275 280 285

Leu Asn Pro Ile Val Tyr Gly Val Lys Thr Lys Glu Ile Arg Gln Arg  
 290 295 300

Ile Leu Arg Leu Phe His Val Ala Thr His Ala Ser Glu Pro  
 305 310 315

<210> 11  
 <211> 995  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (1)..(921)

<400> 11  
 atg gaa agc gag aac aga aga gtg ata aga gaa ttc atc ctc ctt ggt 48  
 Met Glu Ser Glu Asn Arg Arg Val Ile Arg Glu Phe Ile Leu Leu Gly  
 1 5 10 15

ctg acc cag tct caa gat att cag ctc ctg gtc ttt gtg cta gtt tta 96  
 Leu Thr Gln Ser Gln Asp Ile Gln Leu Leu Val Phe Val Leu Val Leu  
 20 25 30

ata ttc tac ttc atc atc ctc cct gga aat ttt ctc att att ttc acc 144  
 Ile Phe Tyr Phe Ile Ile Leu Pro Gly Asn Phe Leu Ile Ile Phe Thr  
 35 40 45

ata aag tca gac cct ggg ctc aca gcc ccc ctc tat ttc ttt ctg ggc 192  
 Ile Lys Ser Asp Pro Gly Leu Thr Ala Pro Leu Tyr Phe Phe Leu Gly  
 50 55 60

- 17 -

aac ttg gcc ttc ctg gat gca tcc tac tcc ttc att gtg gct ccc cgg	240
Asn Leu Ala Phe Leu Asp Ala Ser Tyr Ser Phe Ile Val Ala Pro Arg	
65 70 75 80	
atg ttg gtg gac ttc ctc tct gcg aag aag ata atc tcc tac aga ggc	288
Met Leu Val Asp Phe Leu Ser Ala Lys Lys Ile Ile Ser Tyr Arg Gly	
85 90 95	
tgc atc act cag ctc ttt ttc ttg cac ttc ctt gga gga ggg gag gga	336
Cys Ile Thr Gln Leu Phe Phe Leu His Phe Leu Gly Gly Gly Glu Gly	
100 105 110	
tta ctc ctt gtt gtg atg gcc ttt gac cgc tac atc gcc atc tgc cgg	384
Leu Leu Leu Val Val Met Ala Phe Asp Arg Tyr Ile Ala Ile Cys Arg	
115 120 125	
cct ctg cac tat cct act gtc atg aac cct aga acc tgc tat gca atg	432
Pro Leu His Tyr Pro Thr Val Met Asn Pro Arg Thr Cys Tyr Ala Met	
130 135 140	
atg ttg gct ctg tgg ctt ggg ggt ttt gtc cac tcc att atc cag gtg	480
Met Leu Ala Leu Trp Leu Gly Gly Phe Val His Ser Ile Ile Gln Val	
145 150 155 160	
gtc ctc atc ctc cgc ttg cct ttt tgt ggc cca aac cag ctg gac aac	528
Val Leu Ile Leu Arg Leu Pro Phe Cys Gly Pro Asn Gln Leu Asp Asn	
165 170 175	
ttc ttc tgt gat gtc cca cag gtc atc aag ctg gcc tgc acc gac aca	576
Phe Phe Cys Asp Val Pro Gln Val Ile Lys Leu Ala Cys Thr Asp Thr	
180 185 190	
ttt gtg gtg gag ctt ctg atg gtc ttc aac agt ggc ctg atg aca ctc	624
Phe Val Val Glu Leu Leu Met Val Phe Asn Ser Gly Leu Met Thr Leu	
195 200 205	
ctg tgc ttt ctg ggg ctt ctg gcc tcc tat gca gtc att ctt tgt cgc	672
Leu Cys Phe Leu Gly Leu Leu Ala Ser Tyr Ala Val Ile Leu Cys Arg	
210 215 220	
ata cga ggg tct tct tct gag gca aaa aac aag gcc atg tcc acg tgc	720
Ile Arg Gly Ser Ser Ser Glu Ala Lys Asn Lys Ala Met Ser Thr Cys	
225 230 235 240	
atc acc cat atc att gtt ata ttc ttc atg ttt gga cct ggc atc ttc	768
Ile Thr His Ile Ile Val Ile Phe Phe Met Phe Gly Pro Gly Ile Phe	
245 250 255	
atc tac acg cgc ccc ttc agg gct ttc cca gct gac aag gtg gtt tct	816
Ile Tyr Thr Arg Pro Phe Arg Ala Phe Pro Ala Asp Lys Val Val Ser	
260 265 270	
ctc ttc cac aca gtg att ttt cct ttg ttg aat cct gtc att tat acc	864
Leu Phe His Thr Val Ile Phe Pro Leu Leu Asn Pro Val Ile Tyr Thr	
275 280 285	
ctt cgc aac cag gaa gtg aaa gct tcc atg aaa aag gtg ttt aat aag	912
Leu Arg Asn Gln Glu Val Lys Ala Ser Met Lys Lys Val Phe Asn Lys	
290 295 300	
cac ata gcc tgaaaaaggg cgcaaaaaaa aaaagaataa aaatagactg	961
His Ile Ala	
305	

- 18 -

tagaattttt aaaaaaaaaa aaaaaaaaaa aaaa

995

<210> 12  
 <211> 307  
 <212> PRT  
 <213> Homo sapiens

<400> 12  
 Met Glu Ser Glu Asn Arg Arg Val Ile Arg Glu Phe Ile Leu Leu Gly  
     1                    5                    10                    15  
 Leu Thr Gln Ser Gln Asp Ile Gln Leu Leu Val Phe Val Leu Val Leu  
                     20                    25                    30  
 Ile Phe Tyr Phe Ile Ile Leu Pro Gly Asn Phe Leu Ile Ile Phe Thr  
             35                    40                    45  
 Ile Lys Ser Asp Pro Gly Leu Thr Ala Pro Leu Tyr Phe Phe Leu Gly  
             50                    55                    60  
 Asn Leu Ala Phe Leu Asp Ala Ser Tyr Ser Phe Ile Val Ala Pro Arg  
     65                    70                    75                    80  
 Met Leu Val Asp Phe Leu Ser Ala Lys Lys Ile Ile Ser Tyr Arg Gly  
                     85                    90                    95  
 Cys Ile Thr Gln Leu Phe Phe Leu His Phe Leu Gly Gly Gly Glu Gly  
                     100                    105                    110  
 Leu Leu Leu Val Val Met Ala Phe Asp Arg Tyr Ile Ala Ile Cys Arg  
             115                    120                    125  
 Pro Leu His Tyr Pro Thr Val Met Asn Pro Arg Thr Cys Tyr Ala Met  
     130                    135                    140  
 Met Leu Ala Leu Trp Leu Gly Gly Phe Val His Ser Ile Ile Gln Val  
     145                    150                    155                    160  
 Val Leu Ile Leu Arg Leu Pro Phe Cys Gly Pro Asn Gln Leu Asp Asn  
                     165                    170                    175  
 Phe Phe Cys Asp Val Pro Gln Val Ile Lys Leu Ala Cys Thr Asp Thr  
             180                    185                    190  
 Phe Val Val Glu Leu Leu Met Val Phe Asn Ser Gly Leu Met Thr Leu  
     195                    200                    205  
 Leu Cys Phe Leu Gly Leu Leu Ala Ser Tyr Ala Val Ile Leu Cys Arg  
     210                    215                    220  
 Ile Arg Gly Ser Ser Ser Glu Ala Lys Asn Lys Ala Met Ser Thr Cys  
     225                    230                    235                    240  
 Ile Thr His Ile Ile Val Ile Phe Phe Met Phe Gly Pro Gly Ile Phe  
             245                    250                    255  
 Ile Tyr Thr Arg Pro Phe Arg Ala Phe Pro Ala Asp Lys Val Val Ser  
             260                    265                    270  
 Leu Phe His Thr Val Ile Phe Pro Leu Leu Asn Pro Val Ile Tyr Thr  
     275                    280                    285

- 19 -

Leu Arg Asn Gln Glu Val Lys Ala Ser Met Lys Lys Val Phe Asn Lys  
 290 295 300

His Ile Ala  
 305

<210> 13  
 <211> 1380  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (266)..(1375)

<220>  
 <221> misc\_feature  
 <222> (32)  
 <223> n = A or C or G or T

<220>  
 <221> misc\_feature  
 <222> (55)  
 <223> n = A or C or G or T

<220>  
 <221> misc\_feature  
 <222> (74)  
 <223> n = A or C or G or T

<400> 13  
 tgcttcccca taaggtaaca gctttgttag cnetgtctga catcattgct tgttnactta 60  
 agaactgata ggtntttttt tttttttttt ttcagatatt ctgatggcaa aacaagtgga 120  
 agaaaagagg aagcatgact gcagatcaga tcagttctct ttgtggatta tattttcagt 180  
 aaaatgtatg gatctatctt ttccttggtc ttatatctag atcatgagac ttgactgagg 240  
 ctgtatcctt atcctccatc catct atg gcg aac tat agc cat gca gct gac 292  
 Met Ala Asn Tyr Ser His Ala Ala Asp  
 1 5

aac att ttg caa aat ctc tcg cct cta aca gcc ttt ctg aaa ctg act 340  
 Asn Ile Leu Gln Asn Leu Ser Pro Leu Thr Ala Phe Leu Lys Leu Thr  
 10 15 20 25

tcc ttg ggt ttc ata ata gga gtc agc gtg gtg ggc aac ctc ctg atc 388  
 Ser Leu Gly Phe Ile Ile Gly Val Ser Val Val Gly Asn Leu Leu Ile  
 30 35 40

tcc att ttg cta gtg aaa gat aag acc ttg cat aga gca cct tac tac 436  
 Ser Ile Leu Leu Val Lys Asp Lys Thr Leu His Arg Ala Pro Tyr Tyr  
 45 50 55

ttc ctg ttg gat ctt tgc tgt tca gat atc ctc aga tct gca att tgt 484  
 Phe Leu Leu Asp Leu Cys Cys Ser Asp Ile Leu Arg Ser Ala Ile Cys  
 60 65 70

ttc cca ttt gtg ttc aac tct gtc aaa aat ggt tct acc tgg act tat 532  
 Phe Pro Phe Val Phe Asn Ser Val Lys Asn Gly Ser Thr Trp Thr Tyr  
 75 80 85

- 20 -

ggg act ctg act tgc aaa gtg att gcc ttt ctg ggg gtt ttg tcc tgt	580
Gly Thr Leu Thr Cys Lys Val Ile Ala Phe Leu Gly Val Leu Ser Cys	
90 95 100 105	
ttc cac act gct ttc atg ctc ttc tgc atc agt gtc acc aga tat tta	628
Phe His Thr Ala Phe Met Leu Phe Cys Ile Ser Val Thr Arg Tyr Leu	
110 115 120	
gct atc gcc cat cac cgc ttc tat aca aag agg ctg acc ttt tgg acg	676
Ala Ile Ala His His Arg Phe Tyr Thr Lys Arg Leu Thr Phe Trp Thr	
125 130 135	
tgt ctg gct gtg atc tgt atg gtg tgg act ctg tct gtg gcc atg gca	724
Cys Leu Ala Val Ile Cys Met Val Trp Thr Leu Ser Val Ala Met Ala	
140 145 150	
ttt ccc ccg gtt tta gac gtg ggc act tac tca ttc att agg gag gaa	772
Phe Pro Pro Val Leu Asp Val Gly Thr Tyr Ser Phe Ile Arg Glu Glu	
155 160 165	
gat caa tgc acc ttc caa cac cgc tcc ttc agg gct aat gat tcc tta	820
Asp Gln Cys Thr Phe Gln His Arg Ser Phe Arg Ala Asn Asp Ser Leu	
170 175 180 185	
gga ttt atg ctg ctt ctt gct ctc atc ctc cta gcc aca cag ctt gtc	868
Gly Phe Met Leu Leu Leu Ala Leu Ile Leu Leu Ala Thr Gln Leu Val	
190 195 200	
tac ctc aag ctg ata ttt ttc gtc cac gat cga aga aaa atg aag cca	916
Tyr Leu Lys Leu Ile Phe Phe Val His Asp Arg Arg Lys Met Lys Pro	
205 210 215	
gtc cag ttt gta gca gca gtc agc cag aac tgg act ttt cat ggt cct	964
Val Gln Phe Val Ala Ala Val Ser Gln Asn Trp Thr Phe His Gly Pro	
220 225 230	
gga gcc agt ggc cag gca gct gcc aat tgg cta gca gga ttt gga agg	1012
Gly Ala Ser Gly Gln Ala Ala Ala Asn Trp Leu Ala Gly Phe Gly Arg	
235 240 245	
ggt ccc aca cca ccc acc ttg ctg ggc atc agg caa aat gca aac acc	1060
Gly Pro Thr Pro Pro Thr Leu Leu Gly Ile Arg Gln Asn Ala Asn Thr	
250 255 260 265	
aca ggc aga aga agg cta ttg gtc tta gac gag ttc aaa atg gag aaa	1108
Thr Gly Arg Arg Arg Leu Leu Val Leu Asp Glu Phe Lys Met Glu Lys	
270 275 280	
aga atc agc aga atg ttc tat ata atg act ttt ctg ttt cta acc ttg	1156
Arg Ile Ser Arg Met Phe Tyr Ile Met Thr Phe Leu Phe Leu Thr Leu	
285 290 295	
tgg ggc ccc tac ctg gtg gcc tgt tat tgg aga gtt ttt gca aga ggg	1204
Trp Gly Pro Tyr Leu Val Ala Cys Tyr Trp Arg Val Phe Ala Arg Gly	
300 305 310	
cct gta gta cca ggg gga ttt cta aca gct gct gtc tgg atg agt ttt	1252
Pro Val Val Pro Gly Gly Phe Leu Thr Ala Ala Val Trp Met Ser Phe	
315 320 325	

- 21 -

gcc caa gca gga atc aat cct ttt gtc tgc att ttc tca aac agg gag 1300  
 Ala Gln Ala Gly Ile Asn Pro Phe Val Cys Ile Phe Ser Asn Arg Glu  
 330 335 340 345

ctg agg cgc tgt ttc agc aca acc ctt ctt tac tgc aga aaa tcc agg 1348  
 Leu Arg Arg Cys Phe Ser Thr Thr Leu Leu Tyr Cys Arg Lys Ser Arg  
 350 355 360

tta cca agg gaa cct tac tgt gtt ata tgagg 1380  
 Leu Pro Arg Glu Pro Tyr Cys Val Ile  
 365 370

&lt;210&gt; 14

&lt;211&gt; 370

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 14

Met Ala Asn Tyr Ser His Ala Ala Asp Asn Ile Leu Gln Asn Leu Ser  
 1 5 10 15

Pro Leu Thr Ala Phe Leu Lys Leu Thr Ser Leu Gly Phe Ile Ile Gly  
 20 25 30

Val Ser Val Val Gly Asn Leu Leu Ile Ser Ile Leu Leu Val Lys Asp  
 35 40 45

Lys Thr Leu His Arg Ala Pro Tyr Tyr Phe Leu Leu Asp Leu Cys Cys  
 50 55 60

Ser Asp Ile Leu Arg Ser Ala Ile Cys Phe Pro Phe Val Phe Asn Ser  
 65 70 75 80

Val Lys Asn Gly Ser Thr Trp Thr Tyr Gly Thr Leu Thr Cys Lys Val  
 85 90 95

Ile Ala Phe Leu Gly Val Leu Ser Cys Phe His Thr Ala Phe Met Leu  
 100 105 110

Phe Cys Ile Ser Val Thr Arg Tyr Leu Ala Ile Ala His His Arg Phe  
 115 120 125

Tyr Thr Lys Arg Leu Thr Phe Trp Thr Cys Leu Ala Val Ile Cys Met  
 130 135 140

Val Trp Thr Leu Ser Val Ala Met Ala Phe Pro Pro Val Leu Asp Val  
 145 150 155 160

Gly Thr Tyr Ser Phe Ile Arg Glu Glu Asp Gln Cys Thr Phe Gln His  
 165 170 175

Arg Ser Phe Arg Ala Asn Asp Ser Leu Gly Phe Met Leu Leu Leu Ala  
 180 185 190

Leu Ile Leu Leu Ala Thr Gln Leu Val Tyr Leu Lys Leu Ile Phe Phe  
 195 200 205

Val His Asp Arg Arg Lys Met Lys Pro Val Gln Phe Val Ala Ala Val  
 210 215 220

Ser Gln Asn Trp Thr Phe His Gly Pro Gly Ala Ser Gly Gln Ala Ala  
 225 230 235 240

Ala	Asn	Trp	Leu	Ala	Gly	Phe	Gly	Arg	Gly	Pro	Thr	Pro	Pro	Thr	Leu	
				245					250						255	
Leu	Gly	Ile	Arg	Gln	Asn	Ala	Asn	Thr	Thr	Gly	Arg	Arg	Arg	Leu	Leu	
			260					265					270			
Val	Leu	Asp	Glu	Phe	Lys	Met	Glu	Lys	Arg	Ile	Ser	Arg	Met	Phe	Tyr	
		275					280					285				
Ile	Met	Thr	Phe	Leu	Phe	Leu	Thr	Leu	Trp	Gly	Pro	Tyr	Leu	Val	Ala	
		290				295					300					
Cys	Tyr	Trp	Arg	Val	Phe	Ala	Arg	Gly	Pro	Val	Val	Pro	Gly	Gly	Phe	
305					310					315					320	
Leu	Thr	Ala	Ala	Val	Trp	Met	Ser	Phe	Ala	Gln	Ala	Gly	Ile	Asn	Pro	
				325					330					335		
Phe	Val	Cys	Ile	Phe	Ser	Asn	Arg	Glu	Leu	Arg	Arg	Cys	Phe	Ser	Thr	
			340					345					350			
Thr	Leu	Leu	Tyr	Cys	Arg	Lys	Ser	Arg	Leu	Pro	Arg	Glu	Pro	Tyr	Cys	
		355					360					365				
Val	Ile															
	370															
<210> 15																
<211> 1191																
<212> DNA																
<213> Homo sapiens																
<220>																
<221> CDS																
<222> (1)..(1188)																
<400> 15																
atg	ttt	aga	cct	ctt	gtg	aat	ctc	tct	cac	ata	tat	ttt	aag	aaa	ttc	48
Met	Phe	Arg	Pro	Leu	Val	Asn	Leu	Ser	His	Ile	Tyr	Phe	Lys	Lys	Phe	
	1			5					10					15		
cag	tac	tgt	ggg	tat	gca	cca	cat	gtt	cgc	agc	tgt	aaa	cca	aac	act	96
Gln	Tyr	Cys	Gly	Tyr	Ala	Pro	His	Val	Arg	Ser	Cys	Lys	Pro	Asn	Thr	
			20					25					30			
gat	gga	att	tca	tct	cta	gag	aat	ctc	ttg	gca	agc	att	att	cag	aga	144
Asp	Gly	Ile	Ser	Ser	Leu	Glu	Asn	Leu	Leu	Ala	Ser	Ile	Ile	Gln	Arg	
		35					40					45				
gta	ttt	gtc	tgg	gtt	gta	tct	gca	gtt	acc	tgc	ttt	gga	aac	att	ttt	192
Val	Phe	Val	Trp	Val	Val	Ser	Ala	Val	Thr	Cys	Phe	Gly	Asn	Ile	Phe	
	50					55					60					
gtc	att	tgc	atg	cga	cct	tat	atc	agg	tct	gag	aac	aag	ctg	tat	gcc	240
Val	Ile	Cys	Met	Arg	Pro	Tyr	Ile	Arg	Ser	Glu	Asn	Lys	Leu	Tyr	Ala	
	65				70				75						80	
atg	tca	atc	att	tct	ctc	tgc	tgt	gcc	gac	tgc	tta	atg	gga	ata	tat	288
Met	Ser	Ile	Ile	Ser	Leu	Cys	Cys	Ala	Asp	Cys	Leu	Met	Gly	Ile	Tyr	
				85					90					95		

- 23 -

tta ttc gtg atc gga ggc ttt gac cta aag ttt cgt gga gaa tac aat	336
Leu Phe Val Ile Gly Gly Phe Asp Leu Lys Phe Arg Gly Glu Tyr Asn	
100 105 110	
aag cat gcg cag ctg tgg atg gag agt act cat tgt cag ctt gta gga	384
Lys His Ala Gln Leu Trp Met Glu Ser Thr His Cys Gln Leu Val Gly	
115 120 125	
tct ttg gcc att ctg tcc aca gaa gta tca gtt tta ctg tta aca ttt	432
Ser Leu Ala Ile Leu Ser Thr Glu Val Ser Val Leu Leu Leu Thr Phe	
130 135 140	
ctg aca ttg gaa aaa tac atc tgc att gtc tat cct ttt aga tgt gtg	480
Leu Thr Leu Glu Lys Tyr Ile Cys Ile Val Tyr Pro Phe Arg Cys Val	
145 150 155 160	
aga cct gga aaa tgc aga aca att aca gtt ctg att ctc att tgg att	528
Arg Pro Gly Lys Cys Arg Thr Ile Thr Val Leu Ile Leu Ile Trp Ile	
165 170 175	
act ggt ttt ata gtg gct ttc att cca ttg agc aat aag gaa ttt ttc	576
Thr Gly Phe Ile Val Ala Phe Ile Pro Leu Ser Asn Lys Glu Phe Phe	
180 185 190	
aaa aac tac tat ggc acc aat gga gta tgc ttc cct ctt cat tca gaa	624
Lys Asn Tyr Tyr Gly Thr Asn Gly Val Cys Phe Pro Leu His Ser Glu	
195 200 205	
gat aca gaa agt att gga gcc cag att tat tca gtg gca att ttt ctt	672
Asp Thr Glu Ser Ile Gly Ala Gln Ile Tyr Ser Val Ala Ile Phe Leu	
210 215 220	
ggt att aat ttg gcc gca ttt atc atc ata gtt ttt tcc tat gga agc	720
Gly Ile Asn Leu Ala Ala Phe Ile Ile Ile Val Phe Ser Tyr Gly Ser	
225 230 235 240	
atg ttt tat agt gtt cat caa agt gcc ata aca gca act gaa ata cgg	768
Met Phe Tyr Ser Val His Gln Ser Ala Ile Thr Ala Thr Glu Ile Arg	
245 250 255	
aat caa gtt aaa aaa gag atg atc ctt gcc aaa cgt ttt ttc ttt ata	816
Asn Gln Val Lys Lys Glu Met Ile Leu Ala Lys Arg Phe Phe Phe Ile	
260 265 270	
gta ttt act gat gca tta tgc tgg ata ccc att ttt gta gtg aaa ttt	864
Val Phe Thr Asp Ala Leu Cys Trp Ile Pro Ile Phe Val Val Lys Phe	
275 280 285	
ctt tca ctg ctt cag gta gaa ata cca ggt acc ata acc tct tgg gta	912
Leu Ser Leu Leu Gln Val Glu Ile Pro Gly Thr Ile Thr Ser Trp Val	
290 295 300	
gtg att ttt att ctg ccc att aac agt gct ttg aac cca att ctc tat	960
Val Ile Phe Ile Leu Pro Ile Asn Ser Ala Leu Asn Pro Ile Leu Tyr	
305 310 315 320	
act ctg acc aca aga cca ttt aaa gaa atg att cat cgg ttt tgg tat	1008
Thr Leu Thr Thr Arg Pro Phe Lys Glu Met Ile His Arg Phe Trp Tyr	
325 330 335	
aac tac aga caa aga aaa tct atg gac agc aaa ggt cag aaa aca tat	1056
Asn Tyr Arg Gln Arg Lys Ser Met Asp Ser Lys Gly Gln Lys Thr Tyr	
340 345 350	

gct	cca	tca	ttc	atc	tgg	gtg	gaa	atg	tgg	cca	ctg	cag	gag	atg	cca	1104
Ala	Pro	Ser	Phe	Ile	Trp	Val	Glu	Met	Trp	Pro	Leu	Gln	Glu	Met	Pro	
		355					360					365				
cct	gag	tta	atg	aag	ccg	gac	ctt	ttc	aca	tac	ccc	tgt	gaa	atg	tca	1152
Pro	Glu	Leu	Met	Lys	Pro	Asp	Leu	Phe	Thr	Tyr	Pro	Cys	Glu	Met	Ser	
		370				375					380					
ctg	att	tct	caa	tca	acg	aga	ctc	aat	tcc	tat	tca	tga				1191
Leu	Ile	Ser	Gln	Ser	Thr	Arg	Leu	Asn	Ser	Tyr	Ser					
385						390				395						

```
<210> 16
<211> 396
<212> PRT
<213> Homo sapiens
```

<400> 16																
Met	Phe	Arg	Pro	Leu	Val	Asn	Leu	Ser	His	Ile	Tyr	Phe	Lys	Lys	Phe	
1				5					10					15		
Gln	Tyr	Cys	Gly	Tyr	Ala	Pro	His	Val	Arg	Ser	Cys	Lys	Pro	Asn	Thr	
			20					25					30			
Asp	Gly	Ile	Ser	Ser	Leu	Glu	Asn	Leu	Leu	Ala	Ser	Ile	Ile	Gln	Arg	
		35					40					45				
Val	Phe	Val	Trp	Val	Val	Ser	Ala	Val	Thr	Cys	Phe	Gly	Asn	Ile	Phe	
	50					55					60					
Val	Ile	Cys	Met	Arg	Pro	Tyr	Ile	Arg	Ser	Glu	Asn	Lys	Leu	Tyr	Ala	
65					70					75					80	
Met	Ser	Ile	Ile	Ser	Leu	Cys	Cys	Ala	Asp	Cys	Leu	Met	Gly	Ile	Tyr	
				85					90					95		
Leu	Phe	Val	Ile	Gly	Gly	Phe	Asp	Leu	Lys	Phe	Arg	Gly	Glu	Tyr	Asn	
			100					105					110			
Lys	His	Ala	Gln	Leu	Trp	Met	Glu	Ser	Thr	His	Cys	Gln	Leu	Val	Gly	
		115					120					125				
Ser	Leu	Ala	Ile	Leu	Ser	Thr	Glu	Val	Ser	Val	Leu	Leu	Leu	Thr	Phe	
	130					135					140					
Leu	Thr	Leu	Glu	Lys	Tyr	Ile	Cys	Ile	Val	Tyr	Pro	Phe	Arg	Cys	Val	
145					150					155					160	
Arg	Pro	Gly	Lys	Cys	Arg	Thr	Ile	Thr	Val	Leu	Ile	Leu	Ile	Trp	Ile	
				165					170					175		
Thr	Gly	Phe	Ile	Val	Ala	Phe	Ile	Pro	Leu	Ser	Asn	Lys	Glu	Phe	Phe	
			180					185					190			
Lys	Asn	Tyr	Tyr	Gly	Thr	Asn	Gly	Val	Cys	Phe	Pro	Leu	His	Ser	Glu	
		195					200					205				
Asp	Thr	Glu	Ser	Ile	Gly	Ala	Gln	Ile	Tyr	Ser	Val	Ala	Ile	Phe	Leu	
	210					215					220					
Gly	Ile	Asn	Leu	Ala	Ala	Phe	Ile	Ile	Ile	Val	Phe	Ser	Tyr	Gly	Ser	
225					230					235					240	

[illegible]

- 26 -

ttt cca ttt cga ata gtc cat gat gca gga ttt gga cct tgg tac ttc	339
Phe Pro Phe Arg Ile Val His Asp Ala Gly Phe Gly Pro Trp Tyr Phe	
95 100 105	
aag ttt att ctc tgc aga tac act tca gtt ttg ttt tat gca aac atg	387
Lys Phe Ile Leu Cys Arg Tyr Thr Ser Val Leu Phe Tyr Ala Asn Met	
110 115 120 125	
tat act tcc atc gtg ttc ctt ggg ctg ata agc att gat cgc tat ctg	435
Tyr Thr Ser Ile Val Phe Leu Gly Leu Ile Ser Ile Asp Arg Tyr Leu	
130 135 140	
aag gtg gtc aag cca ttt ggg gac tct cgg atg tac agc ata acc ttc	483
Lys Val Val Lys Pro Phe Gly Asp Ser Arg Met Tyr Ser Ile Thr Phe	
145 150 155	
acg aag gtt tta tct gtt tgt gtt tgg gtg atc atg gct gtt ttg tct	531
Thr Lys Val Leu Ser Val Cys Val Trp Val Ile Met Ala Val Leu Ser	
160 165 170	
ttg cca aac atc atc ctg aca aat ggt cag cca aca gag gac aat atc	579
Leu Pro Asn Ile Ile Leu Thr Asn Gly Gln Pro Thr Glu Asp Asn Ile	
175 180 185	
cat gac tgc tca aaa ctt aaa agt cct ttg ggg gtc aaa tgg cat acg	627
His Asp Cys Ser Lys Leu Lys Ser Pro Leu Gly Val Lys Trp His Thr	
190 195 200 205	
gca gtc acc tat gtg aac agc tgc ttg ttt gtg gcc gtg ctg gtg att	675
Ala Val Thr Tyr Val Asn Ser Cys Leu Phe Val Ala Val Leu Val Ile	
210 215 220	
ctg atc gga tgt tac ata gcc ata tcc agg tac atc cac aaa tcc agc	723
Leu Ile Gly Cys Tyr Ile Ala Ile Ser Arg Tyr Ile His Lys Ser Ser	
225 230 235	
agg caa ttc ata agt cag tca agc cga aag cga aaa cat aac cag agc	771
Arg Gln Phe Ile Ser Gln Ser Ser Arg Lys Arg Lys His Asn Gln Ser	
240 245 250	
atc agg gtt gtt gtg gct gtg ttt ttt acc tgc ttt cta cca tat cac	819
Ile Arg Val Val Val Ala Val Phe Phe Thr Cys Phe Leu Pro Tyr His	
255 260 265	
ttg tgc aga att cct ttt act ttt agt cac tta gac agg ctt tta gat	867
Leu Cys Arg Ile Pro Phe Thr Phe Ser His Leu Asp Arg Leu Leu Asp	
270 275 280 285	
gaa tct gca caa aaa atc cta tat tac tgc aaa gaa att aca ctt ttc	915
Glu Ser Ala Gln Lys Ile Leu Tyr Tyr Cys Lys Glu Ile Thr Leu Phe	
290 295 300	
ttg tct gcg tgt aat gtt tgc ctg gat cca ata att tac ttt ttc atg	963
Leu Ser Ala Cys Asn Val Cys Leu Asp Pro Ile Ile Tyr Phe Phe Met	
305 310 315	
tgt agg tca ttt tca aga agg ctg ttc aaa aaa tca aat atc aga acc	1011
Cys Arg Ser Phe Ser Arg Arg Leu Phe Lys Lys Ser Asn Ile Arg Thr	
320 325 330	
agg agt gaa agc atc aga tca ctg caa agt gtg aga aga tcg gaa gtt	1059
Arg Ser Glu Ser Ile Arg Ser Leu Gln Ser Val Arg Arg Ser Glu Val	
335 340 345	

- 27 -

ctc ata tat tat gat tat act gat gtg tag gccttttatt gtttggttga 1109  
 Leu Ile Tyr Tyr Asp Tyr Thr Asp Val  
 350 355

atcgatatgt acaaagtgt aataaatgtt tcttttcatt aaaaaaaaaa aaaaa 1164

<210> 18  
 <211> 358  
 <212> PRT  
 <213> Homo sapiens

<400> 18  
 Met Gly Phe Asn Leu Thr Leu Ala Lys Leu Pro Asn Asn Glu Leu His  
 1 5 10 15  
 Gly Gln Glu Ser His Asn Ser Gly Asn Arg Ser Asp Gly Pro Gly Lys  
 20 25 30  
 Asn Thr Thr Leu His Asn Glu Phe Asp Thr Ile Val Leu Pro Val Leu  
 35 40 45  
 Tyr Leu Ile Ile Phe Val Ala Ser Ile Leu Leu Asn Gly Leu Ala Val  
 50 55 60  
 Trp Ile Phe Phe His Ile Arg Asn Lys Thr Ser Phe Ile Phe Tyr Leu  
 65 70 75 80  
 Lys Asn Ile Val Val Ala Asp Leu Ile Met Thr Leu Thr Phe Pro Phe  
 85 90 95  
 Arg Ile Val His Asp Ala Gly Phe Gly Pro Trp Tyr Phe Lys Phe Ile  
 100 105 110  
 Leu Cys Arg Tyr Thr Ser Val Leu Phe Tyr Ala Asn Met Tyr Thr Ser  
 115 120 125  
 Ile Val Phe Leu Gly Leu Ile Ser Ile Asp Arg Tyr Leu Lys Val Val  
 130 135 140  
 Lys Pro Phe Gly Asp Ser Arg Met Tyr Ser Ile Thr Phe Thr Lys Val  
 145 150 155 160  
 Leu Ser Val Cys Val Trp Val Ile Met Ala Val Leu Ser Leu Pro Asn  
 165 170 175  
 Ile Ile Leu Thr Asn Gly Gln Pro Thr Glu Asp Asn Ile His Asp Cys  
 180 185 190  
 Ser Lys Leu Lys Ser Pro Leu Gly Val Lys Trp His Thr Ala Val Thr  
 195 200 205  
 Tyr Val Asn Ser Cys Leu Phe Val Ala Val Leu Val Ile Leu Ile Gly  
 210 215 220  
 Cys Tyr Ile Ala Ile Ser Arg Tyr Ile His Lys Ser Ser Arg Gln Phe  
 225 230 235 240  
 Ile Ser Gln Ser Ser Arg Lys Arg Lys His Asn Gln Ser Ile Arg Val  
 245 250 255  
 Val Val Ala Val Phe Phe Thr Cys Phe Leu Pro Tyr His Leu Cys Arg  
 260 265 270

Ile	Pro	Phe	Thr	Phe	Ser	His	Leu	Asp	Arg	Leu	Leu	Asp	Glu	Ser	Ala				
		275					280					285							
Gln	Lys	Ile	Leu	Tyr	Tyr	Cys	Lys	Glu	Ile	Thr	Leu	Phe	Leu	Ser	Ala				
		290				295					300								
Cys	Asn	Val	Cys	Leu	Asp	Pro	Ile	Ile	Tyr	Phe	Phe	Met	Cys	Arg	Ser				
305					310					315					320				
Phe	Ser	Arg	Arg	Leu	Phe	Lys	Lys	Ser	Asn	Ile	Arg	Thr	Arg	Ser	Glu				
				325					330					335					
Ser	Ile	Arg	Ser	Leu	Gln	Ser	Val	Arg	Arg	Ser	Glu	Val	Leu	Ile	Tyr				
			340					345					350						
Tyr	Asp	Tyr	Thr	Asp	Val														
		355																	
<210> 19																			
<211> 2480																			
<212> DNA																			
<213> Homo sapiens																			
<220>																			
<221> CDS																			
<222> (42)..(1157)																			
<400> 19																			
catggcatcc ccagcctagc tcccaatccc actttggcac g atg tta gcc aac agc															56				
															Met Leu Ala Asn Ser				
															1 5				
tcc tca acc aac agt tct gtt ctc ccg tgt cct gac tac cga cct acc	104																		
Ser Ser Thr Asn Ser Ser Val Leu Pro Cys Pro Asp Tyr Arg Pro Thr																			
	10 15 20																		
cac cgc ctg cac ttg gtg gtc tac agc ttg gtg ctg gct gcc ggg ctc	152																		
His Arg Leu His Leu Val Val Tyr Ser Leu Val Leu Ala Ala Gly Leu																			
	25 30 35																		
ccc ctc aac gcg cta gcc ctc tgg gtc ttc ctg cgc gcg ctg cgc gtg	200																		
Pro Leu Asn Ala Leu Ala Leu Trp Val Phe Leu Arg Ala Leu Arg Val																			
	40 45 50																		
cac tcg gtg gtg agc gtg tac atg tgt aac ctg gcg gcc agc gac ctg	248																		
His Ser Val Val Ser Val Tyr Met Cys Asn Leu Ala Ala Ser Asp Leu																			
	55 60 65																		
ctc ttc acc ctc tcg ctg ccc gtt cgt ctc tcc tac tac gca ctg cac	296																		
Leu Phe Thr Leu Ser Leu Pro Val Arg Leu Ser Tyr Tyr Ala Leu His																			
	70 75 80 85																		
cac tgg ccc ttc ccc gac ctc ctg tgc cag acg acg ggc gcc atc ttc	344																		
His Trp Pro Phe Pro Asp Leu Leu Cys Gln Thr Thr Gly Ala Ile Phe																			
	90 95 100																		
cag atg aac atg tac ggc agc tgc atc ttc ctg atg ctc atc aac gtg	392																		
Gln Met Asn Met Tyr Gly Ser Cys Ile Phe Leu Met Leu Ile Asn Val																			
	105 110 115																		
gac cgc tac gcc gcc atc gtg cac ccg ctg cga ctg cgc cac ctg cgg	440																		
Asp Arg Tyr Ala Ala Ile Val His Pro Leu Arg Leu Arg His Leu Arg																			
	120 125 130																		

- 29 -

cgg ccc cgc gtg gcg cgg ctg ctc tgc ctg ggc gtg tgg gcg ctc atc	488
Arg Pro Arg Val Ala Arg Leu Leu Cys Leu Gly Val Trp Ala Leu Ile	
135 140 145	
ctg gtg ttt gcc gtg ccc gcc gcc cgc gtg cac agg ccc tcg cgt tgc	536
Leu Val Phe Ala Val Pro Ala Ala Arg Val His Arg Pro Ser Arg Cys	
150 155 160 165	
cgc tac cgg gac ctc gag gtg cgc cta tgc ttc gag agc ttc agc gac	584
Arg Tyr Arg Asp Leu Glu Val Arg Leu Cys Phe Glu Ser Phe Ser Asp	
170 175 180	
gag ctg tgg aaa ggc agg ctg ctg ccc ctc gtg ctg ctg gcc gag gcg	632
Glu Leu Trp Lys Gly Arg Leu Leu Pro Leu Val Leu Leu Ala Glu Ala	
185 190 195	
ctg ggc ttc ctg ctg ccc ctg gcg gcg gtg gtc tac tcg tcg ggc cga	680
Leu Gly Phe Leu Leu Pro Leu Ala Ala Val Val Tyr Ser Ser Gly Arg	
200 205 210	
gtc ttc tgg acg ctg gcg cgc ccc gac gcc acg cag agc cag cgg cgg	728
Val Phe Trp Thr Leu Ala Arg Pro Asp Ala Thr Gln Ser Gln Arg Arg	
215 220 225	
cgg aag acc gtg cgc ctc ctg ctg gct aac ctc gtc atc ttc ctg ctg	776
Arg Lys Thr Val Arg Leu Leu Leu Ala Asn Leu Val Ile Phe Leu Leu	
230 235 240 245	
tgc ttc gtg ccc tac aac agc acg ctg gcg gtc tac ggg ctg ctg cgg	824
Cys Phe Val Pro Tyr Asn Ser Thr Leu Ala Val Tyr Gly Leu Leu Arg	
250 255 260	
agc aag ctg gtg gcg gcc agc gtg cct gcc cgc gat cgc gtg cgc ggg	872
Ser Lys Leu Val Ala Ala Ser Val Pro Ala Arg Asp Arg Val Arg Gly	
265 270 275	
gtg ctg atg gtg atg gtg ctg ctg gcc ggc gcc aac tgc gtg ctg gac	920
Val Leu Met Val Met Val Leu Leu Ala Gly Ala Asn Cys Val Leu Asp	
280 285 290	
ccg ctg gtg tac tac ttt agc gcc gag ggc ttc cgc aac acc ctg cgc	968
Pro Leu Val Tyr Tyr Phe Ser Ala Glu Gly Phe Arg Asn Thr Leu Arg	
295 300 305	
ggc ctg ggc act ccg cac cgg gcc agg acc tcg gcc acc aac ggg acg	1016
Gly Leu Gly Thr Pro His Arg Ala Arg Thr Ser Ala Thr Asn Gly Thr	
310 315 320 325	
cgg gcg gcg ctc gcg caa tcc gaa agg tcc gcc gtc acc acc gac gcc	1064
Arg Ala Ala Leu Ala Gln Ser Glu Arg Ser Ala Val Thr Thr Asp Ala	
330 335 340	
acc agg ccg gat gcc gcc agt cag ggg ctg ctc cga ccc tcc gac tcc	1112
Thr Arg Pro Asp Ala Ala Ser Gln Gly Leu Leu Arg Pro Ser Asp Ser	
345 350 355	
cac tct ctg tct tcc ttc aca cag tgt ccc cag gat tcc gcc ctc	1157
His Ser Leu Ser Ser Phe Thr Gln Cys Pro Gln Asp Ser Ala Leu	
360 365 370	
tgaacacaca tgccattgcg ctgtccgtgc ccgactccca acgcctctcg ttctgggagg	1217
cttacagggt gtacacacaa gaagggtgggc tgggcacttg gacctttggg tggcaattcc	1277

- 30 -

```

agcttagcaa cgcagaagag tacaaagtgt ggaagccagg gcccagggaa ggcagtgctg 1337
ctggaaatgg cttcttttaa ctgtgagcac gcagagcacc ctttctccag cggatgggaag 1397
tgatgcagag agcccacccg tgcagagggc agaagaggac gaaatgcctt tgggtgggca 1457
gggcattaaa ctgctaaaag ctggttagat ggaacagaaa atgggcattc tggatctaaa 1517
ccgccacagg ggctgagag ctgaagagca ccaggtttg tggacaaagc tactgagatg 1577
cctgttcac tctgacttc tgtctaggct catggatgcc acccccttc atttcggcct 1637
aggcttcccc tctcaccac tgaggcctaa tacaagagtt cctatggaca gaactacatt 1697
ctttctcgca tagtgacttg tgacaattta gacttggcat ccagcatggg atagttgggg 1757
caaggcaaaa ctaacttaga gtttccccct caacaacatc caagtccaaa ccctttttag 1817
gttatccttt cttccatcac atcccccttt ccaggcctcc tccatttttag gtccttaata 1877
ttctttcttt ttctctctct ctcgtttctc tcttctctct cctctctctc cctctctctt 1937
ctcctcttct ctctctctcc ctctctctcc tttgtccaga gtaaggataa aattctttct 1997
actaaagcac tggttctcaa actttttggt ctgagacccc actcttagaa attgaggatc 2057
tcaaagagct ttgcttatat tttgttcttt tgatacttac catactagaa attaaagcga 2117
atacattttt aaaataaata cacatgcaca cattacatta gccatgggag caataatgtc 2177
accacacaca cttcatgaag cctctggaaa actctacagt atacttgtga gagaatgaga 2237
gtgaaagggg caaataacat ctgtgtagca gtattatgaa aatagcttga ccttgtggac 2297
ttctcagag ggttgggtccc tggatcacac ttgagaacc atacttgtcc tgaagtattg 2357
gagttcatgt ctaacttctt ccaggggcat tatgtacagt gctttttatt actgtgggga 2417
gagggcagtg ctaaataaat taatcactac tgataaaaaa aaaaaaaaaa aaaaaaaaaa 2477
aaa
2480

```

<210> 20  
 <211> 372  
 <212> PRT  
 <213> Homo sapiens

```

<400> 20
Met Leu Ala Asn Ser Ser Ser Thr Asn Ser Ser Val Leu Pro Cys Pro
 1             5             10             15
Asp Tyr Arg Pro Thr His Arg Leu His Leu Val Val Tyr Ser Leu Val
      20             25             30
Leu Ala Ala Gly Leu Pro Leu Asn Ala Leu Ala Leu Trp Val Phe Leu
      35             40             45
Arg Ala Leu Arg Val His Ser Val Val Ser Val Tyr Met Cys Asn Leu
      50             55             60
Ala Ala Ser Asp Leu Leu Phe Thr Leu Ser Leu Pro Val Arg Leu Ser
      65             70             75             80

```

[illegible]

```
<210> 21
<211> 19
<212> DNA
<213> Artificial Sequence
<220>
```

- 32 -

&lt;223&gt; Description of Artificial Sequence: Primer LW1282

&lt;400&gt; 21

taataacctgc actgcccac

19

&lt;210&gt; 22

&lt;211&gt; 22

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer LW 1283

&lt;400&gt; 22

tctttccttc tcttctcact cc

22

&lt;210&gt; 23

&lt;211&gt; 32

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer LW 1373

&lt;400&gt; 23

gcataagctt atgctaacac tgaataaaac ag

32

&lt;210&gt; 24

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer LW1374

&lt;400&gt; 24

gcattctcgag tcacatgctg taggatttg

30

&lt;210&gt; 25

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Peptide

&lt;400&gt; 25

Ala Pro Arg Thr Pro Gly Gly Arg Arg

1

5

&lt;210&gt; 26

&lt;211&gt; 32

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer LW1248

- 33 -

<400> 26  
gcatgaattc caatatactt ccccatacct ac 32

<210> 27  
<211> 30  
<212> DNA  
<213> Artificial Sequence  
<220>  
<223> Description of Artificial Sequence: Primer LW1249

<400> 27  
gcatggatcc ggaaaagaag gagaagaaag 30

<210> 28  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1278

<400> 28  
accgctgcct ttttagtc 18

<210> 29  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1279

<400> 29  
ccttctttct gggtacataa gtc 23

<210> 30  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1405

<400> 30  
aagcataaca tggatgaaac aggaaatctg 30

<210> 31  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1406

<400> 31  
aagcataact atactttaca tatttcttc 29

- 34 -

<210> 32  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1280

<400> 32  
tctgcacaca gctcttccat gg 22

<210> 33  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1281

<400> 33  
tcccttgccc agttggtga gg 22

<210> 34  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1385

<400> 34  
gcataagctt ccatggaact tcataacctg 30

<210> 35  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1386

<400> 35  
gcattctcgag ttacccccac agcgtgcag 30

<210> 36  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1329

<400> 36  
gcattctcgag tcagcctaag gttatgttg 29

<210> 37  
<211> 29  
<212> DNA

- 35 -

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer LW1377

&lt;400&gt; 37

gcataagctt atgaacacca cagtgatgc

29

&lt;210&gt; 38

&lt;211&gt; 41

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer LW1387

&lt;400&gt; 38

gagaaatatt ttcttaaaaa aacctgtttt tgcaaaaacg g

41

&lt;210&gt; 39

&lt;211&gt; 41

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer LW1388

&lt;400&gt; 39

ccgtttttgc aaaaacaggt ttttttagaa aaatatttct c

41

&lt;210&gt; 40

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer LW1314

&lt;400&gt; 40

gcatgaattc ccaccttcac catctacctc

30

&lt;210&gt; 41

&lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer LW1315

&lt;400&gt; 41

gcatggatcc gaagacaaaa aagacccag

29

&lt;210&gt; 42

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

- 36 -

<220>  
<223> Description of Artificial Sequence: Primer LW1326  
  
<400> 42  
gcatgaattc atgatggtgg atcccaatgg 30  
  
<210> 43  
<211> 27  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Primer LW1327  
  
<400> 43  
gcatctcgag cctagggctc tgaagcg 27  
  
<210> 44  
<211> 42  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Primer LW1415  
  
<400> 44  
ccatgtatat atttctttgc atgctttcag gcattgacat cc 42  
  
<210> 45  
<211> 42  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Primer LW1416  
  
<400> 45  
ggatgtcaat gcctgaaagc atgcaaagaa atatatacat gg 42  
  
<210> 46  
<211> 30  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Primer LW1308  
  
<400> 46  
gcatgaattc actcacttct catctccttc 30  
  
<210> 47  
<211> 30  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Primer LW1309  
  
<400> 47

- 37 -

gcatggatcc aatctccttt gtcttcactc 30

<210> 48  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1324

<400> 48  
gacgcgatcc atggaaagcg agaacag 27

<210> 49  
<211> 35  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1325

<400> 49  
gacctctcgag tcaggctatg tgcttattaa acacc 35

<210> 50  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1306

<400> 50  
gcatgaattc ttctacttca tcatcctcc 29

<210> 51  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1307

<400> 51  
gcatggatcc aaaggccatc acaacaag 28

<210> 52  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer GV599

<400> 52  
ggcagaagaa ggctattggc cttagacgag 30

- 38 -

<210> 53  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer GV600

<400> 53  
ctgaaacagc gcctcagctc cc 22

<210> 54  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1482

<400> 54  
agctatggcg aactatagcc atgcagc 27

<210> 55  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW148

<400> 55  
agtcctcata taacacagta aggttcc 27

<210> 56  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1310

<400> 56  
gcatgaattc gcagaagaag gctattgg 28

<210> 57  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1311

<400> 57  
gcatggatcc gcagtaaaga agggttgtg 29

<210> 58  
<211> 19

- 39 -

<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1442

<400> 58  
gccattctgt ccacagaag 19

<210> 59  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1443

<400> 59  
tcagttgctg ttatggcac 19

<210> 60  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1440

<400> 60  
aagcggatgt ttagacctct tgtg  
24

<210> 61  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1441

<400> 61  
aacagtcagc aataggaatt gag 23

<210> 62  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer LW1472

<400> 62  
gcatgaattc tgccatgtca atcatttctc tc 32

<210> 63  
<211> 31

- 40 -

<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Primer LW1473  
  
<400> 63  
gcatggatcc gttctgcatt ttccaggctc c 31  
  
<210> 64  
<211> 29  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Primer LW1411  
  
<400> 64  
gcatgaattc tgccaacat catcctgac 29  
  
<210> 65  
<211> 29  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Primer LW1412  
  
<400> 65  
gcatggatcc tacacagcca caacaaccc 29  
  
<210> 66  
<211> 30  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Primer LW1448  
  
<400> 66  
aagcggtagc atgttagcca acagctcctc 30  
  
<210> 67  
<211> 29  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Primer LW1449  
  
<400> 67  
aagctctaga tcagagggcg gaatcctgg 29  
  
<210> 68  
<211> 43  
<212> DNA

- 41 -

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer 217A

&lt;400&gt; 68

taggtcggta gtcaggacac gggagaacag aactgttggt tga

43

&lt;210&gt; 69

&lt;211&gt; 52

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer 217B

&lt;400&gt; 69

gcccctgtgg cggtttagat ccagaatgcc cattttctgt tccatctaac ca

52

&lt;210&gt; 70

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer LW1480

&lt;400&gt; 70

ggttctacct ggacttatgg

20

&lt;210&gt; 71

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer LW1481

&lt;400&gt; 71

taatgaatga gtaagtgcc

20

&lt;210&gt; 72

&lt;211&gt; 42

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer CON103a

&lt;400&gt; 72

tttattaata ttggaaggga caaactggag agcacagaac at

42

&lt;210&gt; 73

&lt;211&gt; 44

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Primer CON103b

&lt;400&gt; 73